



β FSH, β LH and Growth Hormone Gene Expression in Blue Gourami (*Trichogaster trichopterus*, Pallas 1770) During Spermatogenesis and Male Sexual Behavior

Authors: Degani, Gad, Jackson, Karen, Goldberg, Doron, Sarfati, Raheli, and Avtalion, Ramy R.

Source: Zoological Science, 20(6) : 737-743

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.20.737>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

β FSH, β LH and Growth Hormone Gene Expression in Blue Gourami (*Trichogaster trichopterus*, Pallas 1770) During Spermatogenesis and Male Sexual Behavior

Gad Degani^{1,2,*}, Karen Jackson^{1,2}, Doron Goldberg^{1,2}, Raheli Sarfati³
and Ramy R. Avtalion³

¹School of Science and Technology, Tel-Hai Academic College, Upper Galilee, Israel

²MIGAL, Galilee Technological Center, Kiryat Shmona 10200, Israel

³Faculty of Life Sciences, Bar Ilan University, Ramat-Gan 52900, Israel

ABSTRACT—The relationship between gonadal development (histological evidence for spermiogenesis and/or spermatogenesis), sexual behavior (nest-building) and mRNA levels of gonadotropins (β FSH and β LH) and growth hormone (GH) in the male pituitary was investigated. Amplification of β FSH cDNA showed a significantly higher mRNA level in mature males (whether sexually active or not) than in juveniles. However, following PCR amplification of β LH cDNA, a significantly higher mRNA level was found in the sexually active group compared to the sexually inactive group. These results suggest that FSH may participate in spermatogenesis, whereas LH is more involved in spermiogenesis. The GH mRNA level increased slightly during the maturation process but no significant differences were found between the groups studied.

Key words: teleost, RACE-PCR, cloning, gonadotropin, pituitary

INTRODUCTION

Spermatogenesis in teleosts, as in other vertebrates, involves a complex interaction of various hormones along the brain-pituitary-gonad axis (Degani *et al.*, 1998; Lee *et al.*, 1998). One of these hormones is gonadotropin (GtH), a glycoprotein consisting of two subunits, α and β , which are non-covalently bound. There are two distinct GtHs in the pituitary gland of teleosts: FSH (or GtH-I) and LH (or GtH-II). These proteins share a common α subunit but differ in their β subunits, which confer the immunological and biological specificity to each hormone.

The participation of FSH and LH in the female teleost reproductive cycle has been described thoroughly (Nagahama *et al.*, 1994). However, less information has been published on the effects of FSH and LH on gametogenesis in the male. Lee *et al.* (1998) described the relationship between steroids and GtH during spermatogenesis. Apparently, hormonal control is similar to that found in females, with FSH being responsible for the first stages of spermatogenesis and LH for the final stages. In the salmonid species, spermatogenesis is regulated by pituitary FSH and LH. FSH

levels rise in the plasma during the early stage of spermatogenesis (spermiogenesis) and decline during spermiation, whereas the FSH level is very low during spermiogenesis and increases during spermiation (Swanson, 1991; Prat *et al.*, 1996). The salmonids exhibit a difference in sensitivity to FSH and LH in the testis; the sensitivity to LH increases towards the final stages of spermatogenesis (Planas and Swanson, 1995; Swanson *et al.*, 1999).

The synthesis and secretion of gonadotropins are regulated by positive and negative factors that act at the brain, pituitary and gonad levels. Most studies describing changes in the levels of gonadotropins during the teleost reproductive cycle are related to the secretion of gonadotropin, but not to its synthesis. Molecular biology techniques have made it possible to examine changes in the mRNA levels of the gonadotropins during the gonadal cycle. Changes in the expression of FSH and LH during the gonadal cycle have been described recently (Jackson *et al.*, 1999; Gomez *et al.*, 1999; Weil *et al.*, 1995; Meiri *et al.*, 1995; Sohn *et al.*, 1999).

Pituitary growth hormone (GH), a single-chain polypeptide, plays an essential role in the regulation of growth and development by promoting the division, differentiation and enlargement of cells (Corin *et al.*, 1990). The GH gene has been cloned from several teleost species (Venkatesh and Brenner, 1997). Only one type of GH gene has been cloned

* Corresponding author: Tel. +972-4-6953540;
FAX. +972-4-6944980.
E-mail: gad@migal.co.il

in most teleosts, but two GH genes have been found in salmonids and tilapia (see reviews in Venkatesh and Brenner, 1997; Melamed *et al.*, 1998).

The biological activities of GH have been examined in many fish species (Sakata *et al.*, 1993). The metabolic effects of GH include increased protein synthesis, increased use of fat for energy production, and decreased glucose utilization throughout the body (Copeland and Nair, 1994; Salmon *et al.*, 1991). GH in fish also influences osmoregulation (Sakamoto and Hirano, 1993) and reproduction (Kraak *et al.*, 1990; Le Gac *et al.*, 1992).

Little information has been published on the GH level in the plasma of teleosts during growth and maturation, but accumulating data strongly point to the fact that GH is involved in the reproductive cycle of fish. Bjornsson *et al.* (1995) found that the GH level is higher in sexually mature Atlantic salmon (*Salmo salar*) than in immature fish. Le Gac *et al.* (1992) observed an increase in the GH level in male rainbow trout (*Salmo gairdneri*) at the beginning of sperm production, concomitant with an increase in the level of 17α , 20β -dihydroxy-progesterone ($17, 20\text{-P}$). Degani *et al.* (1996, 1998) found that the increase of GH in the plasma of both male and female carp (*Cyprinus carpio*) correlates to the gonadosomatic index.

Blue gourami (*Trichogaster trichopterus*) belongs to the Anabantidae family, which contains 16 genera and about 50 species distributed throughout most of Southern Asia, India and Central Africa. This family belongs to the Labyrinthici suborder, which is characterized by the presence of an air-filled breathing cavity (the labyrinth). The blue gourami shows a very complex social and spawning behavior, and has therefore become a common subject for ethological studies (Hodges and Behre, 1953; Cheal and Davis, 1974; McKinnon and Liley, 1986). The onset of the reproductive period in the male is marked by the development of nuptial coloration, initiation of nest-building behavior and increasing aggressiveness. *T. trichopterus* utilizes a floating bubble nest for egg deposition. This nest is composed of air bubbles enveloped in oral mucus that are released at the surface where they tend to stick together. Cheal and Davis (1974) found that the presence of a nest is not a prerequisite for spawning; in many cases, the female was able to spawn without a nest. However, in many cases when spawning took place without a nest, the eggs were found dispersed over the water surface and many did not develop (unpublished data).

Recently, cDNA sequences of blue gourami β FSH and β LH have been cloned and the mRNA of these hormones quantified in the female during the gonadal cycle (Jackson *et al.*, 1999). Mañanos *et al.* (1997) measured the LH levels in the plasma and the pituitary, as well as the steroidal testosterone (T) and progesterone (P) in the plasma and testis in male blue gouramis during sexual behavior.

The purpose of the present study was to examine β FSH, β LH and GH gene expression during the sexual behavior (nest-building) of male blue gourami.

MATERIALS AND METHODS

Fish

Male blue gouramis, bred and maintained at MIGAL Laboratories in northern Israel, were used in this study. The fish were grown in containers measuring $2 \times 2 \times 0.5$ m, at a temperature of 27°C , and under a light regime of 12L 12D. The fish were fed an artificial diet (45% protein, 7% fat) supplemented by live food (*Artemia salina*). Males and females were maintained separately in groups of about 20 each in containers of 1 m^3 . After two weeks, fish were paired by transferring a male into the presence of a mature female in a small aquarium ($40 \times 20 \times 20$ cm) containing *Elodea*. In most instances, the male started to build the nest on the plant within a few hr.

After weighing and measuring each specimen, testes were taken for histological examination in order to determine the stage of gonadal development, and freshly excised pituitaries were processed immediately for RNA extraction. Donors of pituitary and testes were: 1) mature males that had started to build a nest after coupling (considered as reproductive or sexually active); 2) mature males that were kept in a group without females, hence did not exhibit nest-building behavior (considered as non-reproductive or non-sexually active); and 3) immature males (juveniles).

Histological examination

The pieces of testes were fixed in Bouin and processed until they were embedded in paraffin. Histological sections of $2\text{--}5 \mu\text{m}$ were obtained by using a Reichert-Jung (Austria) microtome. The trichrome of Mallory was used to stain the sections (Humason, 1979).

Measurement of β FSH-I, β LH-II and GH mRNA levels by RT-PCR

The β FSH, β LH and GH mRNA levels were measured by reverse transcriptase (RT) PCR using 18S rRNA as an internal standard (Jackson *et al.*, 1999). This method was employed because the very small size of the blue gourami pituitary gland precluded the use of Northern blotting, which requires relatively large amounts of RNA (Kawasaki, 1990). 18S rRNA was used as an internal standard for the measurements since it is an abundant RNA and its expression is considered stable (Amoureux *et al.*, 1997). The 18S rRNA levels were observed to have a greater uniformity than those of other commonly used internal standards, such as β actin (Ambion Technical Bulletin No. 151, 1997). The 18S level also proved to be stable for the blue gourami and did not vary during different reproductive stages (Jackson *et al.*, 1999).

Total RNA was extracted individually from each pituitary by means of the Rneasy[®] total RNA kit (QIAGEN, Hilden, GR). In most cases, $40\text{--}70 \text{ ng}/\mu\text{l}$ of total RNA were obtained from each pituitary.

In order to measure β FSH, β LH and GH mRNA levels, the cDNA pool for the 3' end was synthesized using an AMV RT (Promega), as described by Frohman (1990). The primer for the synthesis of the 3' cDNA end consisted of an oligo dT (17 bases) linked to a unique 17 mer adapter and a mixture of random decamers. At least five specimens from each of the following stages of the reproductive cycle were used in this study: spermatogenesis (immature males), males maintained together to inhibit nest-building (non-reproductive males) and mature nest-building males (mature reproductive males).

Amplification of cDNA of β FSH, β LH and 18S rRNA

Each amplification of β FSH, β LH and GH cDNA was coupled to an amplification of the 18S rRNA internal standard. All amplifications were conducted in separate tubes since many attempts to use multiplex PCR resulted in considerable interference between the different reactions. The concentration of each primer pair and the number of cycles were calibrated carefully to allow for amplifications that were linearly dependent on the initial concentration of the target

cDNA. β FSH, β LH and GH cDNAs were amplified with gene-specific primers (see Table 1) at a concentration of 6.25 pmol/reaction. 18S rRNA cDNA was amplified with specific primers obtained from Ambion as part of its Quantum RNA kit at a concentration of 5 pmol/reaction. The reagents, except for the cDNA and the primers, were

prepared as a batch for each set of reactions to minimize possible variations that might be independent of the initial cDNA concentrations. The PCR amplifications were carried out in 50 μ l reactions using 2.5 units of Taq polymerase (Promega), a suitable reaction

Table 1. Sequence of primers for cDNA synthesis and PCR

Primer Name	cDNA Target	Use	Nucleotide Sequence
P1	β FSH	Complete cDNA cloning	5'GTCTGTACAGATGTTTAGAGAG
P2	β FSH	Complete cDNA cloning	5'AACGTGGGATGAATGATGAGTG
P3	β LH	Complete cDNA cloning	5'CTGGCTAACCTGCCGCTGACAC
P4	β LH	Complete cDNA cloning	5'TTGCTTTTGGTTTGCTGTGCAG
P5	GH	Complete cDNA cloning	5'AGAAGTGAACCTGAACCTGTATC
P6	GH	Complete cDNA cloning	5'CATTGTGCTGGAACCTGG
dT ₁₇ -adapter	Universal	cDNA synthesis	5' GACTCGAGTCGACATCGA(T) ₁₇
Adapter	Universal	RACE PCR	5'GACTCGAGTCGACATCG




Fig. 1. Determination of the exponential range of amplification for β LH, β FSH, GH and 18S. The amount of PCR products is expressed in arbitrary units of intensity as defined by the densitometry software (PCBAS-2).

buffer (Promega), 1.5 mM MgCl₂, 0.2 mM of each nucleotide, specific primers, 5 µl of cDNA and 20 µl mineral oil.

Determination of cycling parameters

It is important to select the appropriate number of cycles so that the amplification products are clearly visible on an agarose gel, but also that the amplification is in the exponential range and has yet not reached a plateau. The optimal number of cycles was checked for each primer pair using the most concentrated cDNA. Ten reaction tubes for each primer pair were prepared as described above. Following 26 complete cycles, two reaction tubes of each primer pair were removed from the thermocycler (every two cycles, from 26 to 34). The PCR products were electrophoresed in agarose gel and quantified as described below (Fig. 1). For βFSH, βLH, GH and 18S, only 30 cycles produced enough PCR products for the semi-quantitative analysis while maintaining linear dependence on the cDNA concentration. Thus, cycling parameters were: 3 min denaturation at 94°C followed by 30 cycles, each consisting of 1 min denaturation at 94°C, 1 min annealing at 54°C and 1 min extension at 72°C.

Quantification of PCR products

The PCR products (half of the PCR reaction) were electrophoresed on 2% agarose gel at 120 mA for 45 min. The DNA was then stained by ethidium bromide (0.03 mg in 100ml distilled water), and imaged with the Eagle Eye II Image analysis system (STRATAGENE). The gels were imaged at three different exposure times to prevent a possible saturation of the images. The resulting images

were analyzed with the aid of the PCBAS-2 densitometry software (Raytest, GR).

The variations in the expression of the βFSH, βLH and GH genes at the mRNA level were evaluated by first measuring the amount of specific PCR products and then dividing it by the amount of the amplified internal standard. This normalization step was intended to eliminate the effect of random changes in the experimental setup.

Statistical analysis

The differences between the mRNA levels of the various hormones were analyzed by ANOVA, and the difference of means between mRNA levels by the Student's t-test, with the level of significance in different groups set at $p < 0.05$ (Parker, 1986).

RESULTS

Fig. 2 illustrates the gonadal morphology of juvenile males, mature non-reproductive males and mature reproductive males (after spawning). In all three groups, the same cellular stages were found but in different proportions (Table 2). No morphological differences could be observed between the testis of males from the two latter groups. However, there was a decrease in the number of spermatozoa in the center of the lobule of mature reproductive males pro-




Fig. 2. Histological section showing different stages of gonadal development in the blue gourami male *T. trichopterus*: (A) Mature non-reproductive fish (note the concentration of spermatozoa in the middle of the lobule; see arrow); (B). Mature reproductive fish during sexual behavior (note the decrease in the quantity of the spermatozoa in the center of the lobule of reproductive fish when compared to non-reproductive fish; see arrow); (C). Young males at the beginning of spermiogenesis (note the lower concentration of spermatozoa when compared to fully developed fish); and (D) Early cell stages in *T. Trichopterus*: 1 = spermatogonia, 2 = primary spermatocyte, 3 = secondary spermatocyte, 4 = spermatid H.E.

Table 2. Germ cell constitution of gourami testis at different developmental stages




Stage	Juveniles	Mature Reproductive (Nest-building)	Mature Non-reproductive (Non-nest-building)
Spermatogonia	+++	+	+
Spermatocyte I	+	+	+
Spermatocyte II	++	++	++
Spermatid	+	++	++
Spermatozoa	+ / ++	+++	+++

cessed after spawning. The testes of mature reproductive and mature non-reproductive fish were similar; those of juvenile fish contained relatively few spermatozoa (Fig. 2).

Variations in β FSH and β LH and GH mRNA levels during the reproductive cycle

Amplification of β FSH cDNA produced a 306 bp product, that of β LH cDNA a 284 bp product, and that of the internal standard (the cDNA of 18S rRNA) a 430 bp product. The identity of each PCR product was confirmed by DNA sequencing.

The variations of mRNA levels of β FSH are shown in Fig. 3. There was no significant difference (t-test; $p < 0.05$) in the β FSH between mature fish, whether non-nest-builders

**Fig. 3.** Variations in β FSH transcript levels during the gonadal cycle. Each histogram represents an average of five independent measurements (mean \pm SE). The amount of PCR products is expressed in arbitrary units of intensity as defined using the densitometry software (PCBAS-2).**Fig. 4.** Variations in β LH transcript levels during the gonadal cycle. Each histogram represents an average of five independent measurements (mean \pm SE).**Fig. 5.** Variations in GH transcript levels during the gonadal cycle. Each histogram represents an average of five independent measurements (mean \pm SE).

(mature non-reproductive) or nest-builders (mature reproductive). Both levels were significantly higher than in juveniles.

The variations of mRNA levels of β LH are shown in Fig. 4. Amplifications of cDNA from juveniles and non-reproductive mature fish were low, showing no significant difference (t-test; $p > 0.05$) between them. β LH expression rose significantly (ANOVA; $p < 0.05$) at the nest-building stage, as manifested by the increased amount of PCR products.

The product generated by the amplification of GH was 733 bp long. The identity of the PCR product was confirmed by DNA sequencing. There were no significant differences in the GH mRNA level (ANOVA or t-test; $p > 0.05$) between juveniles, mature non-reproductive and mature reproductive specimens (Fig. 5).

DISCUSSION

In the present study, differences in the biosynthesis of β FSH, β LH and GH during gonadal development and the male sexual behavior of blue gourami were determined by quantifying their respective mRNAs. The results support the hypothesis that FSH participates in spermatogenesis. FSH transcripts were significantly lower in immature fish in comparison to mature fish. However, spermatozoa could be found in the center of the lobules of immature fish.

In males sampled immediately after spawning, there was a remarkable decrease in the quantity of spermatozoa in the center of the lobule, as observed in histological preparations. It is noteworthy that the blue gourami is a multi-spawning species, and after successful spawning, the same male is able to fertilize another female 24 hr later. Consequently, spermatogenesis occurs continuously in mature fish during sexual behavior.

In females blue gourami, the gonadal development follows an asynchronous pattern, with oocytes found at different developmental stages (Jackson *et al.*, 1994). In concerning β FSH expression similar results were observed in females. β FSH transcripts were first detected in immature females during low vitellogenesis (i.e., when the percentage of oocytes found at vitellogenesis was low) and an accentuated increase occurred in mature non-reproductive females

found at high vitellogenesis (i.e., when the majority of the oocytes were at vitellogenesis). The β FSH mRNA level also remained high in post-vitellogenic females during the final stage of oocyte maturation and spawning. These mature reproductive females able to spawn again after a 24–48 hr break (Jackson *et al.*, 1999).

The β LH mRNA level was low in young and mature non-reproductive males, increasing rapidly only in reproductive males (nest-builders) (Fig. 4). The β LH level was also significantly higher in the pituitaries and plasma of reproductive males than in non-reproductive males, as was found by Mañanos *et al.* (1997). Thus, a temporal correlation exists between transcription (β LH mRNA in the pituitary), translation (LH levels in the pituitary) and release (LH levels in the plasma) with respect to this hormone. LH is probably related to the final stages of spermatogenesis/spermiation and triggers the final maturation (acquiring of motility) and release of sperm, probably via steroids (Nagahama *et al.*, 1994).

The expression of β LH in the male gourami differs from that found in females (Jackson *et al.*, 1999). The expression of β LH gene in females begins during low vitellogenesis (immature or maturing females), with a significant increase in non-reproductive mature females (high vitellogenic females). It also remains high in mature reproductive females (females found at the final oocyte maturation stage, FOM). Thus, a significant increase of β LH in females is already detected in non-reproductive females (contrary to males, where the increase in β LH level occurs only in reproductive specimens). In females, LH is stored in the pituitary and is released to the plasma only in the presence of the male (Degani *et al.*, 1997). The situation in males is different. The delay between LH production and release is not required; LH is probably produced and released by mature reproductive males during nest-building.

The results of this study show that the GH mRNA level remains high, not only during the male's growth (juvenile fish) but also during spermatogenesis and sexual behavior (Fig. 5). Therefore, the GH mRNA level in blue gourami males (as found in the present study) and females (Degani *et al.*, 2002) remains high during final stages of the reproductive cycle and during sexual behavior. An increase in GH levels during the final stages of gonadal development was also detected in male and female rainbow trout (Gomez *et al.*, 1999). Histological changes in somatotrophs during ovulation and spawning indicate that these cells become markedly more active (Melamed, 1998).

Nest-building males exhibit significantly higher levels of plasma steroids (testosterone, progesterone) than do non-nest-builders (Mañanos *et al.*, 1997). This increase in steroid levels could have a positive effect on GH production. The interaction between GH and steroids has been studied recently in teleosts. The injection of recombinant salmon GH in killifish stimulated the growth of the gonads and raised the level of sex steroids circulating in the blood (Singh *et al.*, 1988). Injected testosterone can stimulate pituitary mRNA (Huggard *et al.*, 1996) or cause an increase in serum con-

centration of GH in rainbow trout (Holloway and Leatherland, 1997). Le Gac *et al.* (1992) studied the action of GH in the testis of trout, and suggested that at the end of reproductive cycle high GH levels are associated with 17α -hydroxy, 20β -dihydroprogesterone, a progestin necessary for efficient spawning in this fish. Together, these results demonstrate the participation of GH in the reproductive cycle, but more information is needed since there is a difference of action according to species. The specific action of steroids and GnRH on GH gene expression is currently being studied in this laboratory.

REFERENCES

- Ambion Technical Bulletin No. 151 (1997) Use of Internal and External Standards or Reference RNAs for Accurate Quantitation of RNA Levels, <http://www.ambion.com/techlib/tb/tb151.html>
- Amoureux MC, Van Gool D, Herrero MT, Dom R, Colpaert F, Pauwels P (1997) Regulation of metallothionein-III (GIF) mRNA in the brain of patients with Alzheimer's disease is not impaired. *Mol Chem Neuropathol* 32: 101–121
- Björnsson BT, Taranger GL, Hansen T, Stefánsson S, Haux C (1995) The interrelationship between photoperiod, growth hormone and sexual maturation of adult Atlantic salmon (*Salmo salar*). *Gen Comp Endocr* 97: 70–81
- Cheal M, Davis RE (1974) Sexual behavior: Social and ecological influences in the anabantoid fish *Trichogaster trichopterus*. *Behav Biol* 20: 435–455
- Copeland KC, Nair KS (1994) Acute growth hormone effects on amino acid and lipid metabolism. *J Clin Endocrin Metab* 78: 1040–1047
- Corin RE, Guller S, Wu KY, Sonenberg M (1990) Growth hormone and adipose differentiation: Growth hormone-induced antimitogenic state in 3T3-F442A preadipose cells. *Proc Natl Acad Sci USA* 87: 7507–7511
- Degani G, Boker R, Jackson K (1996) Growth hormone, gonad development and steroid levels in female carp. *Comp Biochem Physiol* 115: 133–140
- Degani G, Mananos EL, Jackson K, Abraham M, Zohar Y (1997) Changes in plasma and pituitary in female blue gourami during the end of vitellogenesis and final maturation. *J Exp Zool* 279: 377–385
- Degani G, Boker R, Jackson K (1998) Growth hormone, sexual maturation and steroids in male carp (*Cyprinus carpio*). *Fish Physiol Biochem* 120C: 433–440
- Degani G, Goldberg D, Jackson K (2002) The β FSH, β LH and growth hormone in female and male *Trichogaster trichopterus*. Cloning and sequencing of its cDNA and analysis of its mRNA expression during oogenesis and subsequent treatment with GnRH analog. *Biol Reprod* 66: 22
- Frohman, MA (1990) RACE: Rapid Amplification of cDNA Ends. In "PCR Protocols: A Guide to Methods and Applications" Ed by MA Innis, DH Gelfand, JJ Sninsky, TJ White, Academic Press Inc, San Diego, pp 23–28
- Gomez JM, Weil C, Ollitrault M, Le Bail PY, Breton B, Le Gac F (1999) Growth hormone (GH) and gonadotropin subunit gene expression and pituitary and plasma changes during spermatogenesis and oogenesis in rainbow trout (*Oncorhynchus mykiss*). *Gen Comp Endocr* 113: 413–428
- Hodges WR, Behre EH (1953) Breeding behavior, early embryology and melanophore development in the anabantoid fish (*T. trichopterus*). *Copeia* 1953: 100–107
- Holloway AC, Leatherland JF (1997) Effect of gonadal steroid hormones on plasma growth hormone concentrations in sexually

- immature rainbow trout (*Oncorhynchus mykiss*). Gen Comp Endocr 105: 246–254
- Huggard D, Khakoo Z, Kassam G, Mahmoud SS, Habibi HR (1996) Effect of testosterone maturational gonadotropin subunit messenger ribonucleic acid levels in the goldfish pituitary. Biol Reprod 54: 1184–1191
- Humason GL (1979) Animal Tissue Technique, 4th ed., WH Freeman & Co, San Francisco
- Jackson K, Goldberg G, Ofir M, Abraham M, Degani G (1999) Blue gourami (*Trichogaster trichopterus*) gonadotropic β subunits (I & II) cDNA sequences and expression during oogenesis. J Mol Endocrinol 23: 177–187
- Kawasaki ES (1990) Amplification of RNA. In "PCR Protocols: A Guide to Methods and Applications" Ed by MA Innis, DH Gelfand, JJ Sninsky, TJ White, Academic Press Inc, San Diego, pp 21–27
- Kraak G van der, Rosenblum PM, Peter RE (1990) Growth hormone dependent potentiation of gonadotropin-stimulated steroid production by ovarian follicles of the goldfish. Gen Comp Endocr 79: 233–239
- Lee STL, Kime DE, Lam TJ, Tan CH (1998) Synthesis of 17,20 α/β -dihydroxy-4- pregnen-3-one and 5 β -pregnanes in spermatozoa of primary and 17 α -methyl testosterone- induced secondary male grouper (*Epinephelus coioides*). Gen Comp Endocr 112: 1–9
- Le Gac F, Ollitraut M, Loir M, Bail PY (1992) Evidence for binding and action of growth hormone in trout. Biol Reprod 46: 949–957
- Mañanos E, Zohar Y, Degani G (1997) The relationship between gonadotropin and sexual behavior of male *Trichogaster trichopterus* (Pallas). Ind J Fisher 44: 239–246
- McKinnon, JS, Liley NR (1986) Asymmetric species in response to female sexual pheromones by males of two species of *Trichogaster* (Pisces: Belontiidae). Can J Zool 65: 1129–1134
- Meiri I, Gothliff Y, Knibb WR, Zohar Y, Elizur A (1995) Preovulatory changes in gonadotropin gene expression and secretion in the gilthead sea bream, *Spaurus aurata*. Proceedings of the Fifth International Symposium on Reproductive Physiology of Fish, p 37
- Melamed P, Rosenfeld H, Elizur A., Yaron Z (1998) Endocrine regulation of gonadotropin and growth hormone gene transcription in fish. Comp Biochem Physiol 119: 325–338
- Nagahama Y, Miura T, Kobayashi T (1994) The onset of spermatogenesis in fish, Ciba Foundation Symposium
- Parker RE (1986) Introduction to Statistics for Biology. Ed by E Arnold, Cambridge university press, London, pp 26–28
- Planas JV, Swanson P (1995) Maturation-associated changes in the response of the salmon testis to the steroidogenic actions of gonadotropins (GtH I and GtH II) *in vitro*. Biol Reprod 52: 697–704
- Prat F, Sumpter JP, Tyler CR (1996) Validation of radioimmunoassay for two salmon gonadotropins (GtH I and GtH II) and their plasma concentrations throughout the reproductive cycle in male and female rainbow trout (*Oncorhynchus mykiss*). Biol Reprod 54: 1375–1382
- Sakata S, Nose T, Moriyama S, Hirano T, Kawauchi H (1993) Flounder growth hormone: Isolation, characterization, and effects on growth of juvenile rainbow trout. Gen Comp Endocr 89: 396–404
- Salmon F, Cuneo R, Sonksen PH (1991) Growth hormone and protein metabolism. Horm Res 36 (suppl 1): 41–43
- Sakamoto T, Hirano T (1993) Expression of insulin-like growth factor I gene in osmoregulatory organs during seawater adaptation of the salmonid fish: Possible mode of osmoregulatory action of growth hormone. Proc Natl Acad Sci USA 90: 1912–1916
- Singh H, Griffith RW, Takahashi AP, Kawauchi H, Stegeman JJ (1988) Regulation of gonadal steroidogenesis in (*Fundulus heteroclitus*) by recombinant salmon growth hormone and purified salmon prolactin. Gen Comp Endocr 72: 144–153
- Sohn YC, Yoshiura Y, Kobayashi M, Aida K (1999) Seasonal changes in mRNA levels of gonadotropin and thyrotropin subunits in the goldfish (*Carassius auratus*). Gen Comp Endocr 113: 436–444
- Swanson P (1991) Salmon Gonadotropins: Reconciling Old and New Ideas, Ed by AP Scott, DE Kime, MS Rolfe, Proceedings of the Fourth International Symposium on Reproduction Physiology of Fish, Sheffield, pp 30–34
- Swanson P, Bernard M, Nozaki M, Suzuku K, Kawauchi H, Dickhoff WW (1999) Gonadotropins I and II in juvenile Coho Salmon. Gen Comp Endocr 113: 413–428
- Venkatesh B, Brenner S (1997) Genomic structure and sequence of the pufferfish (*Fugu rubripes*) growth hormone-encoding gene: A comparative analysis of teleost growth hormone genes. Gene 187: 211–215
- Weil C, Bougoussa-Houadec M, Gallais C, Itoh S, Sekine S, Valotaire Y (1995) Preliminary evidence suggesting variations of GtH 1 and GtH 2 mRNA levels at different stages of gonadal development in rainbow trout (*Oncorhynchus mykiss*). Gen Comp Endocr 100: 327–336

(Received November 7, 2002 / Accepted March 3, 2003)