

Latest early-earliest middle Miocene deep-sea molluscs in the Japan Sea borderland – the warm-water Higashibessho fauna in Toyama Prefecture, central Japan

Authors: Amano, Kazutaka, Hamuro, Toshikazu, and Hamuro, Masui

Source: Paleontological Research, 8(1) : 29-42

Published By: The Palaeontological Society of Japan

URL: <https://doi.org/10.2517/prpsj.8.29>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Latest early-earliest middle Miocene deep-sea molluscs in the Japan Sea borderland – the warm-water Higashibessho fauna in Toyama Prefecture, central Japan

KAZUTAKA AMANO¹, TOSHIKAZU HAMURO² AND MASUI HAMURO²

¹Department of Geoscience, Joetsu University of Education, 1 Yamayashiki, Joetsu City, Niigata Prefecture 943-8512, Japan
(e-mail: amano@juen.ac.jp)

²Toyama Paleontological Research Club, 100 Nishitakagi, Kosugi-machi, Toyama Prefecture 939-0303, Japan

Received June 5, 2003; Revised manuscript accepted September 23, 2003

Abstract. Eighty-seven species of molluscs were obtained from the uppermost lower-lower middle Miocene Higashibessho Formation at Shimo-sasahara, Yatsuo Town in Toyama Prefecture, central Japan. Among them, *Pagodula shojii* is new to science. Judging from the autochthonous species, the Higashibessho Formation was deposited at the lower sublittoral to upper bathyal depth. Both the deep-sea and the derived shallow-water species include many warm-water dwellers. During the latest early-earliest middle Miocene, the deep-sea species migrated from the Pacific side of central Honshu to the Japan Sea through deep-sea pathways.

Key words: Latest early to earliest middle Miocene, deep-sea, Mollusca, Higashibessho Formation

Introduction

In the Oligocene, the “Japanese Islands” were located at the eastern margin of the Asian Continent. The Japan Sea began to open as a marginal sea in the early Miocene (Iijima and Tada, 1990). No deep-sea molluscan faunas have been recorded from the uppermost lower-lower middle Miocene deposits in the central Japan Sea borderland, other than the upper part of the Kurosedani and the lower part of the Higashibessho Formation in Toyama Prefecture (Tsuda, 1960).

Tsuda (1960) reported a deep-sea assemblage IV represented by *Solemya tokunagai*, *Portlandia* (*Megayoldia*) aff. *thraciaeformis*, *Propeamussium transnipponica*, and *Dentalium* (*Fissidentalium*) *yokoyamai* from the upper part of the Kurosedani Formation. Saito (1988) also recorded ill-preserved deep-water molluscs including *Propeamussium* cf. *tateiwai* Kanehara from the Nanamagari Formation in Ishikawa Prefecture and correlated them with Tsuda’s assemblage IV. However, the exact age of the formation and most of the species were uncertain. When they recorded 19 species including warm-water planktonic species such as *Aturia* sp. and *Clio itoigawai* Shibata from the Higashibessho Formation, Shimizu *et al.*

(2000) noted that the formation was deposited in the lower sublittoral to bathyal zones. They also inferred the existence of a cold deep-water mass below a warm-water current. Amano *et al.* (2000) found a deep-water dweller, *Neilo* (*Multidentata*) *multidentata* (Khomenko), in the Higashibessho Formation. This species is widely distributed in the northwestern Pacific in Oligocene to middle Miocene rocks. From the Higashibessho Formation, Amano *et al.* (2001) described the oldest vesicomys in the Japan Sea borderland including *Calypptogena* sp. and *Vesicomya kawadai* (Aoki).

Thus, the Higashibessho fauna (Kaseno, 1964) consists of deep-water species, including some characteristics of the chemosynthetic community. The Kurosedani and Higashibessho Formations were deposited during the Mid-Neogene Climatic Optimum (Tsuchi, 1987; Amano *et al.*, 2001). During this warm age, the shallow-water tropical Kurosedani fauna (Itoigawa, 1988) distributed to Yamagata Prefecture while the subtropical shallow-sea Kadonosawa fauna (originally Otuka, 1939, redefined by Itoigawa, 1988) prevailed to the southwestern part of Hokkaido. Therefore, it is reasonable that shallow warm-water species have been described from the Kurosedani and Higashibessho formations.




Figure 1. Locality of fossils (from Amano *et al.*, 2001, fig. 1; using the topographical map of “Yatsuo”, scale 1:25,000, published by Geographical Survey Institute of Japan).

Chinzei (1978, 1981, 1986) proposed the subsurface cold-current “Oyashio senryu” at the Mid-Neogene Climatic Optimum. Probably taking account of this proposal, Shimizu *et al.* (2000) considered that the deep water was cold during the time of deposition of the Higashibessho Formation as mentioned above. On the other hand, when they proposed a new Pliocene taxodont species of *Acilana*, Noda *et al.* (1989) denied the existence of the “Oyashio senryu” in the Japan Sea borderland at the Mid-Neogene Climatic Optimum. Karasawa *et al.* (1992, 1995) also suggested that the lower sublittoral to bathyal environment in the Japan Sea borderland was warm, based on the distribution of the crustacean *Bathynomus*, which occurred also from the Higashibessho Formation. To resolve this confusion, it is necessary to examine the temperature implied by the deep-water fauna during the latest early-earliest middle Miocene in the Japan Sea borderland in detail.

Moreover, Ozawa *et al.* (1986) pointed out that the muddy bottom associations of the upper part of the Kurosedani Formation are closely related to that of the Oidawara Member of the Mizunami Group in Gifu Prefecture. However, no other comparisons have been made between the Higashibessho and the deep-sea faunas in the central Pacific side of Japan.

Fortunately, we were able to collect many well-preserved specimens from the horizon of the Higashibessho Formation slightly higher than the level from which Amano *et al.* (2001) described the vesicomyids at Shimo-sasahara. In this paper, we list all these species including one new gastropod and discuss the climatic condition and the biogeographic significance of the fauna.

Geological settings

In and around Yatsuo Town, the Neogene deposits

consist of the Nirehara, Iwaine, Iozen, Kurosedani, Higashibessho, Tenguyama, Otokawa and Mita Formations in ascending order (Sakamoto and Nozawa, 1960; Hayakawa and Takemura, 1987; Ogasawara *et al.*, 1989). The Higashibessho Formation (Fujita and Nakagawa, 1948) consists mainly of muddy sediments overlying the Kurosedani Formation (Tsuda and Chiji, 1950). On the other hand, the upper part of the Kurosedani Formation is predominantly siltstone or siltstone-dominated alternation. Sometimes, it is difficult to separate lithologically the Kurosedani from the Higashibessho Formation. The upper limit of the Kurosedani Formation is usually defined as the upper surface of the Yamadanaka Tuff (Tsuda, 1953).

We collected many molluscan specimens from the Higashibessho Formation at about 250 m west of Shimo-sasahara, Yatsuo Town, Toyama Prefecture (Figure 1). This locality is the same as that of Amano *et al.* (2001). The large outcrop consists of a 2.3 m – thick alternation (L) of fine-grained sandstone and mudstone in the lower part yielding vesicomyid fossils, and a 21.2 m-thick black mudstone (UM) bearing calcareous concretions in the upper part (Figure 2).




Figure 2. Columnar section of the fossil locality. A. Black mudstone, B. Sandstone, C. Alternation of fine-grained sandstone and mudstone, D. Calcareous concretions. L, UM, US show horizons of fossils (*see the text*).

Many well-preserved fossils occur in the upper mudstone of the cliff. In the middle part of the mudstone, a 60 cm-thick sandstone lens (US) including many packed shells can be traced for 20 m. In this area, the Yamadanaka Tuff is exceptionally intercalated in the lower part of the Higashibessho Formation (Sakamoto and Nozawa, 1960). Stratigraphically, the upper mudstone of the fossil locality is situated about 75 m lower than the Yamadanaka Tuff. Therefore, the Higashibessho Formation here is a contemporaneous heterotopic facies of the upper part of the Kurosedani Formation to the west of Yatsuo Town.

As already reported by Hasegawa and Takahashi (1992), Yanagisawa (1999) and Amano *et al.* (2001), the planktonic microfossils from the lowest part of the Higashibessho Formation indicate the diatom NPD 3A and 3B of Akiba (1986), the calcareous nannofossils CN3 of Okada and Bukry (1980) and the planktonic foraminiferal zone N8 of Blow (1969) and the radiolarian *Calocyletta costata* zone of Riedel and Sanfilippo (1978). Thus, the Higashibessho Formation at Shimo-sasahara can be assigned to the latest early-earliest middle Miocene. However, based on the magnetostratigraphic data, Itoh *et al.* (1999) assigned the Kurosedani and the lower part of the Higashibessho Formations to the C5Br Chron which is slightly younger than the age based on the microfossils. Although this problem for now remains unsolved, we use the age based on the microfossils in this paper.

Occurrences of molluscs

Molluscan fossils were obtained from three horizons: the lower alternation (L), upper mudstone (UM), and the sandstone lens intercalated in the mudstone (US) (Figure 2; Table 1). In the lower alternation (L), six bivalve species including two vesicomyids sporadically occur as articulated shells. As described by Amano *et al.* (2001), the commissures of vesicomyids are parallel to the bedding plane despite the shells being articulated.

In the upper mudstone (UM), 51 species of bivalves, gastropods and scaphopods (Figures 3, 4) occurred sporadically, including one new species. Most specimens are well preserved, including many articulated bivalves. In contrast, 35 species of bivalves and gastropods (Figure 5) were obtained from the sandstone lens (US) intercalated in the upper mudstone. All bivalve specimens here are disarticulated.

Judging from these occurrences, the assemblages from the lower alternation and upper mudstone are autochthonous while that of the upper sandstone lens is allochthonous.

Table 1. Molluscan fossils from the Higashibessho Formation at Shimo-sasahara. Number shows total number of specimens. Number in parentheses shows number of articulated specimens. * Warm-water genera or species, ** after Higo *et al.* (1999).

Species	L	UM	US	Depth (m)**
* <i>Lamellinucula hokoensis</i> (Kanehara)	8 (5)			50–3000
* <i>Tindaria</i> sp.	1	1		100–3610
<i>Lucinoma acutilienatum</i> (Conrad)	3 (2)	2 (1)	1	100–200
<i>Thyasira tokunagai</i> Kuroda and Habe	7 (1)			5–300
<i>Vesicomya kawadai</i> (Aoki)	10 (1)			100–9050
<i>Calyptogena</i> sp.	7 (4)			500–5960
<i>Ennucula osawanoensis</i> (Tsuda)		9 (9)		0–2320
<i>Acila</i> (<i>Acila</i>) <i>submirabilis</i> Makiyama		3 (3)		50–800
<i>Bathymalletia inermis</i> (Yokoyama)		3 (1)		50–6200
<i>Neilo</i> (<i>Multidentata</i>) <i>multidentata</i> (Khomeenko)		4		–
* <i>Neilonella tsukigawaensis</i> Kurihara		1 (1)		50–1400
<i>Portlandia</i> (<i>Portlandella</i>) <i>lischkei</i> (Smith)		11 (7)		100–1400
<i>P.</i> (<i>Megayoldia</i>) sp.		1		–
<i>Bathymodiolus</i> ? sp.		4 (3)		–
<i>Solamen fornicatum</i> (Yokoyama)		2 (2)		30–300
* <i>Propeamussium tateiwai</i> Kanehara		5		40–2500
<i>Delectopecten</i> sp.		2 (1)		20–3080
<i>Gloripallium izurense</i> Masuda		2		0–20
<i>Acesta goliath</i> (Sowerby)		1		100–1417
<i>Pycnodonte</i> ? sp.		2		–
<i>Ostrea</i> sp.		1 (1)	1	0–600
<i>Macoma</i> (<i>Macoma</i>) sp.		1 (1)		0–1300
<i>Solen</i> sp.		1		0–115
<i>Pandora</i> sp.		1		10–300
<i>Periploma yokoyamai</i> Makiyama		7 (6)		100–1250
<i>P. mitsuganoense</i> Araki		2 (2)		100–1250
<i>Thracia kamayasikiensis</i> Hatai		1		4–300
* <i>Lyonsiella mitsuganoensis</i> Shibata		3 (1)		500–1200
<i>Poromya osawanoensis</i> Tsuda		5 (1)		30–350
<i>Cardiomya mitsuganoensis</i> Shibata		4 (2)		10–800
<i>Ginebis osawanoensis</i> (Tsuda)		1		50–800
<i>Calliostoma</i> (<i>Calotropis</i>) <i>simane</i> Nomura and Hatai		1	2	0–400
<i>Cryptonatica ichishiana</i> (Shibata)		7		0–3000
* <i>Sinum ineptum</i> (Yokoyama)		1		10–50
<i>Semicassis</i> ? sp.		1		–
<i>Liracassis japonica</i> (Yokoyama)		4		–
* <i>Echinophoria etchuensis</i> (Hatai and Nisiyama)		7		150–567
<i>Cymatiidae</i> gen. et sp. indet.		1		–
<i>Boreotrophon osawanoensis</i> (Tsuda)		3		3–1000
<i>Pagodula shojii</i> sp. nov.		5		3–1000
<i>Babylonia kokozurana</i> Nomura		1		0–50
<i>Buccinidae</i> gen. et sp. indet.		1		–
<i>Zeuxis kometubus</i> (Otuka)		11		0–200
* <i>Neadmete nakayamai</i> Habe		7		100–200
<i>Megasurcula yokoyamai</i> (Otuka)		5		–
<i>M.</i> sp.		5		–
<i>Cochlespira osawanoensis</i> (Tsuda)		4		150–300
<i>Tomopleura osawanoensis</i> Tsuda		1		5–130
* <i>Comitas</i> sp.		2		10–600
<i>Eoscapander corpulenta</i> (Yokoyama)		4		100–300
* <i>Bowdenathea</i> sp.		30		–
* <i>Clio</i> sp.		2		0
<i>Dentalium</i> sp.		1		0–150
* <i>Fissidentalium yokoyamai</i> (Makiyama)		13		100–400
<i>F.</i> sp.		1		100–3000
<i>Laevidentalium</i> sp.		6		20–1400
<i>Saccula kongiensis</i> (Otuka)			2	10–450
<i>Barbatia</i> (<i>Savignyarca</i>) <i>osawanoensis</i> Tsuda			1	0–20
<i>Anadara</i> (<i>Anadara</i>) <i>watanabei</i> (Kanehara)			4	0–50

Table 1. Continued

Species	L	UM	US	Depth (m)**
<i>Anadara (Scapharca) makiyamai</i> Hatai and Nisiyama			14	0–60
* <i>Bellucina civica</i> (Yokoyama)			1	50–400
* <i>Notomyrtea</i> sp.			1	30–200
* <i>Nipponocrassatella osawanoensis</i> (Tsuda)			26	0–200
<i>Veremolpa minoensis</i> Itoigawa			7	0–40
<i>Minolia tukiyesiensis</i> (Oyama and Saka)			1	0–100
<i>Protorotella</i> sp.			1	–
<i>Sigaretornus?</i> sp.			2	–
* <i>Vicaryella atukoe</i> (Otuka)			1	–
* <i>Calyptrea tubura</i> Otuka			1	20–300
<i>Euspira meisensis</i> (Makiyama)			1	5–2433
* <i>Polinices mizunamiensis</i> Itoigawa			3	0–100
<i>Cryptonatica</i> sp.			1	0–3000
* <i>Gyrineum osawanoense</i> (Tsuda)			3	0–200
* <i>Chicoreus</i> sp.			1	0–300
<i>Siphonalia osawanoensis</i> Tsuda			2	10–300
* <i>Mitrella (Indomitrella) mizunamiensis</i> Itoigawa			8	0–100
<i>Reticunassa</i> sp.			1	0–50
* <i>Conus (Asprella) toyamaensis</i> Tsuda			6	0–550
<i>Inquisitor kurodai</i> (Tsuda)			3	10–1100
* <i>Gemmula osawanoensis</i> Tsuda			1	20–1000
* <i>Bathytoma osawanoense</i> Tsuda			1	50–1140
* <i>Strioterebrum osawanoense</i> Tsuda			1	0–100
* <i>Subula osawanoensis</i> Tsuda			2	0–100
<i>Pyrgiscus?</i> sp.			3	–
<i>Turbonilla</i> sp.			1	10–800
<i>Rhizorus tokiensis</i> (Itoigawa)			3	5–410
<i>Eocylichna tokiensis</i> Itoigawa			1	10–450
<i>Ringicula minoensis</i> Itoigawa			4	5–1020

Systematic description of a new species

Family Muricidae Rafinesque, 1815
 Subfamily Trophoninae Cossmann, 1903
 Genus ***Pagodula*** Monterosato, 1884

Type species.—*Murex vaginata* Cristofori and Jan, 1832. This fossil species was subsequently designated as the type species by Radwin and D'Attilio (1976).

Remarks.—According to Houart (2001), this genus is characterized by a fusiform shell with shouldered whorls, distinct spines, no or few spiral cords, and long, open siphonal canal. He distinguished *Boreotrophon* Fischer, 1884 from this genus based on the former's lacking spiral cords and having adapically pointed axial ribs. We have chosen to assign our new species to *Pagodula* on the basis of shell characters outlined by Houart (2001).

Most species of *Pagodula* live in the lower sublittoral to bathyal depths of the Mediterranean Sea and the northeastern Atlantic Ocean (Bouchet and Warén, 1985; Houart, 2001). Recently, *P. kosunorum* was found in 200–250 m depth, off Northeast Taiwan (Houart and Lan, 2003). Geologically, the oldest spe-

cies of *Pagodula* has been recorded from the middle Miocene in Europe (La Perna, 1996).

***Pagodula shojii* sp. nov.**

Figures 4.1, 4.2, 4.4, 4.5


Trophon (Boreotrophon) osawanoensis Tsuda, 1959, pl. 4, fig. 12 (*non* pl. 4, figs. 11, 13).

Type specimen.—Holotype, JUE (abbreviation of Joetsu University of Education) no. 15744, 27.6 mm high, 11.0 mm wide; paratype, JUE no. 15745-1, 22.0 mm high, 10.7 mm wide; paratype, JUE no. 15745-2, 23.5 mm high, 9.8 mm wide; paratype, JUE no. 15745-3, 26.0 mm high, 12.2 mm wide; paratype, JUE no. 15745-4, 7.6 mm high, 4.8 mm wide.

Type locality.—Shimo-sasahara, Yatsuo Town, Toyama Prefecture (36°33'54"N, 137°8'45"E).

Diagnosis.—Medium-sized *Pagodula* characterized by rather weak spines, long siphonal canal, few spiral cords (two on last whorl) and numerous axial ribs (11–14 on last whorl).

Description.—Shell medium-sized for genus, attaining 27.6 mm in height, narrow and fusiform; proto-



conch partly preserved, of more than one smooth angulate whorl; teleoconch of six whorls. Spire rather high, occupying about two-fifths of shell height. Suture rather deep. Axial ribs located at shoulder, 12–15 on penultimate whorl and 12–15 on last whorl, ending in rather weak spine. One spiral cord at shoulder of early whorls; one at shoulder and another distinct cord below shoulder of last whorl. Aperture ovate; inner lip covered by thin callus; outer lip thin and smooth on inner side. Siphonal canal open, very long and slightly curved.

Remarks.—*Pagodula shojii* closely resembles the Recent species, *P. kosunorum* Houart and Lan, 2003 from Taiwan in its size (height = 23.47 mm), number of whorls (six and a half) and number of spiral cords (two on the last whorl). However, *P. shojii* has more numerous (10 on the last whorl in *P. kosunorum*) as well as weaker spines than in *P. kosunorum*. *P. shojii* also resembles the Recent species *P. cossmanni* (Locard, 1897) in shell outline, number of whorls (six), weak spines on the shoulder and number of axial ribs on the last whorl. *P. shojii* can be distinguished from *P. cossmanni* in having less numerous spiral cords on the last whorl than in *P. cossmanni* (three-five).

It is possible that this new species might be confused with *Boreotrophon osawanoensis* (Tsuda, 1959) (Figure 4.3, 4.7). *B. osawanoensis* can be distinguished easily from the new species by having no spiral cord, a more flattened area above the shoulder, and more prominent, lamellated axial ribs whose ends are pointed apically. From this point of view, one of the paratypes of *B. osawanoensis* (Tsuda, 1959, pl. 4, fig. 12) can be identified with the new species because of its two spiral cords and 12 axial ribs on the last whorl.

Distribution.—Known from the type locality and Kashio in the Higashibessho Formation.

Etymology.—This new species is named after Dr. Shoji Fujii, who has contributed to the Neogene stratigraphy and paleontology of Toyama Prefecture.


Paleoenvironmental significance

Judging from the autochthonous molluscan fossils from both the lower alternation and the upper mudstone, the Higashibessho Formation at Shimosasahara was deposited at lower sublittoral to upper bathyal depths (Table 1). This is supported by estimations using benthic foraminifers from the upper part of the Kurosedani Formation (upper to middle bathyal depths; Hasegawa and Takahashi, 1992) and crustacean *Bathynomus* from the Higashibessho Formation (lower sublittoral to bathyal depths; Karasawa *et al.*, 1992). In contrast, the upper sandstone lens includes an allochthonous assemblage consisting of shallow marine molluscs (Table 1).

The lower alternation yields two warm-water genera, *Lamellinucula* and *Tindaria*. Many warm-water genera can be found in the upper mudstone assemblage: *Tindaria*, *Neilonella*, *Propeamussium*, *Lyonsiella*, *Sinum*, *Echinophoria*, *Comitas*, *Bowdenatheca* and *Clio*. This is the oldest record of *Neadmete nakayamai* Habe, now living in Tosa Bay and the southern Japan Sea (Higo *et al.*, 1999). It is noteworthy that no cold-water genera can be recognized in the assemblage from the upper mudstone. As discussed by Amano *et al.* (2001), *Neilo* (*Multidentata*) *multidentata* from the northwestern Pacific is a non-boreal species. From the upper sandstone lens, 14 warm-water genera and subgenera can be recognized among 32 genera: *Bellucina*, *Notomyrtea*, *Nipponocrassatella*, *Vicaryella*, *Calyptraea*, *Polinices*, *Gyrineum*, *Chicoreus*, *Mitrella* (*Indomitrella*), *Conus*, *Gemma*, *Bathytoma*, *Strioterebrum* and *Subula*.

From this composition, we infer that the deep water in the latest early-earliest middle Miocene was not so cold as in the Recent Japan Sea and a warm-water current affected the sea surface. Such climatic conditions deny the existence of the cold undercurrent “Oyashio senryu” here and may support the claim of

- ◆ **Figure 3.** Bivalves and scaphopods from the lower alternation (L) and the upper mudstone (UM) of the Higashibessho Formation. **1.** *Lamellinucula hokoensis* (Kanehara), ×2, JUE no. 15746, L. **2.** *Portlandia* (*Portlandella*) *lischkei* (Smith), ×1, JUE no. 15747, UM. **3.** *Neilonella tsukigawaensis* Kurihara, ×2.3, JUE no. 15748, UM. **4.** *Bathymalletia inermis* (Yokoyama), ×1.5, JUE no. 15749, UM. **5.** *Delectopecten* sp., ×3, JUE no. 15750, UM. **6.** *Propeamussium tateiwai* Kanehara, ×2, JUE no. 15751, UM. **7.** *Ennucula osawanoensis* (Tsuda), ×1.5, JUE no. 15752, UM. **8.** *Acila* (*Acila*) *submirabilis* Makiyama, ×1.2, JUE no. 15753, UM. **9.** *Bathymodiolus*? sp., ×1, JUE no. 15754, UM. **10.** *Solamen fornicatum* (Yokoyama), ×1.6, JUE no. 15755, UM. **11.** *Thyasira tokunagai* Kuroda and Habe, ×2, JUE no. 15756, L. **12.** *Lucinoma acutilienatum* (Conrad), ×1, JUE no. 15757, UM. **13.** *Lyonsiella mitsuganoensis* Shibata, ×3.25, JUE no. 15758, UM. **14.** *Periploma mitsuganoense* Araki, ×1, JUE no. 15759, UM. **15.** *Periploma yokoyamai* Makiyama, ×1, JUE no. 15760, UM. **16.** *Poromya osawanoensis* Tsuda, ×2, JUE no. 15761, UM. **17.** *Cardiomya mitsuganoensis* Shibata, ×3.1, JUE no. 15762, UM. **18.** *Neilo* (*Multidentata*) *multidentata* (Khomenko), ×1, JUE no. 15763, UM. **19.** *Acesta goliath* (Sowerby), ×0.75, JUE no. 15764, UM. **20.** *Gloripallium izurense* Masuda, ×1, JUE no. 15765, UM. **21.** *Dentalium* sp., ×1.2, JUE no. 15766, UM. **22.** *Fissidentalium yokoyamai* (Makiyama), ×0.75, JUE no. 15767, UM.






Figure 5. Molluscan fossils from the sandstone lens (US) of the Higashibessho Formation. **1.** *Barbatia (Savignyarca) osawanoensis* Tsuda, $\times 2$, JUE no. 15784. **2.** *Veremolpa minoensis* Itoigawa, $\times 2.4$, JUE no. 15785. **3.** *Notomyrtea* sp., $\times 3$, JUE no. 15786. **4.** *Saccella kongiensis* (Otuka), $\times 2.1$, JUE no. 15787. **5.** *Polinices mizunamiensis* Itoigawa, $\times 3$, JUE no. 15788. **6.** *Anadara (Anadara) watanabei* (Kanehara), $\times 1.2$, JUE no. 15789. **7.** *Anadara (Scapharca) makiyamai* Hatai and Nisiyama, $\times 1$, JUE no. 15790. **8.** *Nipponocrassatella osawanoensis* (Tsuda), $\times 1.5$, JUE no. 15791. **9, 10.** *Gyrineum osawanoense* (Tsuda); 9, $\times 3$, JUE no. 15792-1; 10, $\times 3$, JUE no. 15792-2. **11.** *Conus (Asprella) toyamaensis* Tsuda, $\times 1.5$, JUE no. 15793. **12.** *Siphonalia osawanoensis* Tsuda, $\times 1.65$, JUE no. 15794. **13.** *Inquisitor kurodai* (Tsuda), $\times 1.5$, JUE no. 15795. **14.** *Vicaryella atukoe* (Otuka), $\times 1$, JUE no. 15796. **15.** *Gemmula osawanoensis* Tsuda, $\times 1$, JUE no. 15797. **16.** *Mitrella (Indomitrella) mizunamiensis* Itoigawa, $\times 3$, JUE no. 15798. **17.** *Subula osawanoensis* Tsuda, $\times 1$, JUE no. 15799.

Ogasawara (1994) who denied the existence of a cold current such as the Recent Oyashio during the Miocene.

Biogeographical significance

Similar deep-sea molluscan fossils to the lower alternation and upper mudstone have been found from the upper lower to lower middle Miocene deposits in

the Setouchi, Sanin-Hokuriku, and Saikai Provinces of Southwest Japan and in the Pohang Basin of South Korea (Figure 6).

In the Sanin-Hokuriku Province, Japan Sea side of central to western part of Honshu, the deep-water fauna is known from the Nanamagari Formation in Ishikawa Prefecture (Saito, 1988), the Kounoura Shale Member of Uchiura Group in Fukui Prefecture (Nakagawa and Takemura, 1985) and the Fuganji

➤ **Figure 4.** Gastropods from the upper mudstone (UM) of the Higashibessho Formation. **1, 2, 4, 5.** *Pagodula shojii* sp.nov.; 1a, $\times 1.6$, 1b, $\times 1.5$, JUE no. 15745-1, paratype; 2a, $\times 1.5$, 2b, $\times 1.55$, JUE no. 15745-3, paratype; 4a,b, $\times 1.55$, JUE no. 15744, holotype; 5a, $\times 1.6$, 5b, $\times 1.5$, JUE no. 15745-2. **3, 7.** *Boreotrophon osawanoensis* (Tsuda); 3, $\times 2.1$, JUE no. 15768-1; 7, $\times 1$, JUE no. 15768-2. **6.** *Tomopleura osawanoensis* Tsuda, $\times 1$, JUE no. 15769. **8.** *Cochlespira osawanoensis* (Tsuda), $\times 1.3$, JUE no. 15770. **9.** *Megasurcula yokoyamai* (Otuka), $\times 1$, JUE no. 15771. **10.** *Comitas* sp., $\times 2$, JUE no. 15772. **11, 13.** *Neadmete nakayamai* Habe; 11, $\times 2$, JUE no. 15773-1; 13, $\times 2$, JUE no. 15773-2. **12.** *Eoscapander corpulenta* (Yokoyama), $\times 1.5$, JUE no. 15774. **14.** *Clio* sp., $\times 1.7$, JUE no. 15775. **15, 16.** *Bowdenathea* sp.; 15, $\times 1.3$, JUE no. 15776-1; 16, $\times 1.3$, JUE no. 15776-2. **17.** *Sinum ineptum* (Yokoyama), $\times 1$, JUE no. 15777. **18.** *Ginebis osawanoensis* (Tsuda), $\times 2$, JUE no. 15778. **19.** *Calliostoma (Calotropis) simane* Nomura and Hatai, $\times 2$, JUE no. 15779. **20.** *Babylonia kokozurana* Nomura, $\times 1$, JUE no. 15780. **21.** *Cryptonatica ichishiana* (Shibata), $\times 1.5$, JUE no. 15781. **22.** *Echinophoria etchuensis* (Hatai and Nisiyama), $\times 2$, JUE no. 15782. **23.** *Liracassis japonica* (Yokoyama), $\times 1$, JUE no. 15783.



Figure 6. Sites where late early to early middle Miocene deep-water faunas have been collected. Paleogeographic map based on Ogasawara and Nagasawa (1992). 1. Higashibessho Formation (present study). 2. Nanamagari Formation (Saito, 1988). 3. Uchiura Group (Nakagawa and Takemura, 1985). 4. Tottori Group (Akagi *et al.*, 1992). 5. Katsuta Group (Taguchi, 2002). 6. Bihoku Group (Itoigawa and Nishikawa, 1976; Okamoto *et al.*, 1989, 1990; Okamoto, 1992). 7. Susa Group (Okamoto *et al.*, 1983). 8. Hagjeon, Heunghae, Duho formations in Pohang Basin, Japan Sea side of South Korea (Kanehara, 1936; Yoon, 1979; Lee, 1992). 9. Ichishi Group (Shibata, 1970). 10. Morozaki Group (Shikama and Kase, 1976; Yamaoka, 1993). 11. Oidawara Formation (Itoigawa *et al.*, 1981).

Mudstone Member of Tottori Group in Tottori Prefecture (Akagi *et al.*, 1992). Among them, *Acila-Saccella* and *Limatula-Propeamussium* associations of the Kounoura Member include many common species and genera (five species and nine genera) with the Higashibessho fauna (Table 2).

In the eastern part of Setouchi Province, Pacific side of central Honshu, such a deep-water fauna has been recorded from the Oidawara Formation of the Mizunami Group in Gifu Prefecture (Itoigawa *et al.*, 1981), the Morozaki Group in Aichi Prefecture (Shikama and Kase, 1976; Yamaoka, 1993) and the Ichishi Group in Mie Prefecture (Shibata, 1970). The fauna

from the Oidawara Formation contains 10 species and 10 genera in common with the Higashibessho fauna (Table 2). The Morozaki Group yields 10 species and nine genera in common with the Higashibessho fauna. Nine species and seven genera of the Ichishi Group are in common.

In the western part of Setouchi Province, inland of western Honshu, the upper part of the Bihoku Group in Hiroshima Prefecture (Itoigawa and Nishikawa, 1976; Okamoto *et al.*, 1989, 1990; Okamoto, 1992) and the Takakura Formation of the Katsuta Group in Okayama Prefecture (Taguchi, 2002) yield a similar fauna to the Higashibessho. Seven species and 13 genera of the upper part of the Bihoku Group are common to it and the Higashibessho fauna, while the Takakura Formation shares six species and seven genera with it.

From the Maeji Sandstone Member of the Susa Group (Okamoto *et al.*, 1983) in Saikai Province, only two species and three genera are common with the Higashibessho fauna herein treated. There are four species in common between the Higashibessho and the middle Miocene faunas from the Hagjeon, Heunghae, and Duho Formations in Pohang Basin, Japan Sea side of South Korea (Kanehara, 1936; Yoon, 1979; Lee, 1992): *Lamellinucula hokoensis*, *Acila* (*Acila*) *submirabilis*, *Propeamussium tateiwai* and *Lucinoma acutilineatum* (Table 2). *L. hokoensis* has been recorded only from the type locality in Korea and the Higashibessho Formation and is an endemic species of the Japan Sea side.

The lower part of the Higashibessho Formation is a contemporaneous heterotopic facies of the upper part of the Kurosedani Formation. Therefore, it is reasonable that derived shallow-water species in the upper sandstone lens share many species with the shallow-water assemblage of the Kurosedani Formation (Table 3). Of the 25 species in the coarse deposits intercalated in the Yamadanaka Tuff at Tsuzara (Kaneko and Goto, 1992), to the east of Shimo-sasahara, 16 species are shared with the Higashibessho Formation. In the shallow-water assemblages I–III of the Kurosedani Formation (Tsuda, 1960), nine species are shared with the Higashibessho Formation. It is noteworthy that the molluscs from the Shukunohora Facies of the Mizunami Group in Gifu Prefecture, on the central Pacific side of Honshu (Itoigawa *et al.*, 1981) contain 11 species and seven genera in common with the Higashibessho Formation.

Up to this time, few detailed comparisons have been carried out especially on the relationship between the Higashibessho and the upper lower to lower middle Miocene deep-water faunas in the Setouchi Province

Table 2. Comparison between the deep-sea assemblages of the Higashibessho Formation with those of other upper lower to lower middle Miocene formations. A closed circle in the list indicates the same species while an open one means same genus. *Uc = Uchiura Group, Oi = Oidawara Formation, Ic = Ichishi Group, Mo = Morozaki Group, Bh = Bihoku Group, Kt = Katsuta Group, SK = Formations in Pohang Basin, South Korea.

Species	Formation*	Uc	Oi	Ic	Mo	Bh	Kt	SK
<i>Lamellinucula hokoensis</i>			○			○	○	●
<i>Ennucula osawanoensis</i>			●	○	○	●		
<i>Acila (Acila) submirabilis</i>		●	●	○	○	●	○	●
<i>Bathymalletia inermis</i>		○	●	●	●	○	○	
<i>Neilo (Multidentata) multidentata</i>					●			
<i>Neilonella tsukigawaensis</i>		○	○	○	○	○		
<i>Portlandia (Portlandella) lischkei</i>		○	○		○	○	○	○
<i>Solamen fornicatum</i>				●	●			
<i>Propeamussium tateiwai</i>		●		●	●	●	●	●
<i>Delectopecten</i> sp.		○	○		○	○	○	
<i>Acesta goliath</i>		○	○	○	●	○	○	○
<i>Lucinoma acutilienatum</i>		●	●	●	●	●	●	●
<i>Thyasira tokunagai</i>			○	○				○
<i>Macoma (Macoma)</i> sp.		○			○	○		
<i>Calyptogena</i> sp.								○
<i>Periploma yokoyamai</i>		○			●			
<i>P. mitsuganoense</i>		○		●	●	●	●	
<i>Lyonsiella mitsuganoensis</i>				●				
<i>Poromya osawanoensis</i>			○	○		○		
<i>Cardiomya mitsuganoensis</i>			●	●		○	●	
<i>Ginebis osawanoensis</i>			●		○	○		
<i>Cryptonatica ichishiana</i>			○	●	●			○
<i>Sinum ineptum</i>			●					
<i>Liracassis japonica</i>		●	●			●	●	
<i>Boreotrophon osawanoensis</i>			●	○	○	○		○
<i>Babylonia kokozurana</i>			○					
<i>Megasurcula yokoyamai</i>			●			○		
<i>Eoscapander corpulenta</i>				●				
<i>Bowdenathea</i> sp.		○						
<i>Clio</i> sp.					○	○	○	
<i>Fissidentalium yokoyamai</i>		●	○		●	●	●	

because of the scarcity of fossil records from the Higashibessho Formation. In consequence of our huge collection, it has become clear that both deep- and shallow-water assemblages of the Higashibessho Formation resemble the fauna in Setuchi Province. The deep-sea species have in common direct development or a short pelagic larval stage, such as the protobranchs, *Propeamussium*, *Lucinoma*, *Thyasira*, *Calyptogena*, *Lyonsiella*, *Cardiomya*, buccinids and so on (Kasyanov *et al.*, 1990; Bouchet and Warén, 1994; Van Dover, 2000). This leads us to infer that the Japan Sea was connected with the Pacific by deep-sea pathways at least in the latest early-earliest middle Miocene. Moreover, diatom study has revealed that the age of the Morozaki Group on the Pacific side can be assigned to NPD2B of Akiba (1986) (Gladenkov, 1998; Ito *et al.*, 1999). Thus, the deep-water species migrated from the Pacific side into the Japan Sea borderland

just after the formation of a deep-sea basin in the latest early-earliest middle Miocene.

Acknowledgments

We are grateful to Geerat J. Vermeij (University of California at Davis) for his critical reading of the manuscript. We thank Roland Houart (Institut Royal des Sciences naturelles de Belgique) for his information and his help in examining the genus *Pagodula*. We also thank Hiroshi Saito (National Science Museum, Tokyo) and Masanori Shimamoto (Tohoku University) for their help in examining some Recent and fossil specimens. We are grateful to Takashi Matsumura (Museum of Nature and Human Activities, Hyogo) and Alan G. Beu (Institute of Geological and Nuclear Sciences) for reviewing our manuscript and providing many useful comments.

Table 3. Comparison between the shallow-water species of the Higashibessho Formation with those of other upper lower to lower middle Miocene formations. A closed circle in the list indicates the same species while an open one means same genus. *Ky = Kurosedani Formation at Iguridani, Ku = Kurosedani assemblages I–III, Sh = Shukunohora Formation.

Species	Formation*	Ky	Ku	Sh
<i>Sacella kongiensis</i>		●	●	○
<i>Barbatia (Savignyarca) osawanoensis</i>		●	●	○
<i>Anadara (Anadara) watanabei</i>		●		
<i>A. (Scapharca) makiyamai</i>		●	●	●
<i>Lucinoma acutlienatum</i>		●	●	
<i>Bellucina civica</i>				●
<i>Nipponocrassatella osawanoensis</i>		●	●	●
<i>Veremolpa minoensis</i>				●
<i>Calliostoma (Calotropis) simane</i>		●	○	
<i>Protorotella</i> sp.		○	○	○
<i>Calyptrea tubura</i>				●
<i>Euspira meisensis</i>		●	●	●
<i>Polinices mizunamiensis</i>				●
<i>Gyrineum osawanoense</i>		●	●	○
<i>Siphonalia osawanoensis</i>		●	●	○
<i>Mitrella (Indomitrella) mizunamiensis</i>				●
<i>Conus (Asprella) toyamaensis</i>		●	●	
<i>Inquisitor kurodai</i>		●		○
<i>Gemmula osawanoensis</i>		●		●
<i>Bathytoma osawanoense</i>		●		
<i>Strioterebrum osawanoense</i>		●		
<i>Subula osawanoensis</i>		●		
<i>Rhizorus tokiensis</i>				○
<i>Eocylichna tokiensis</i>				●
<i>Ringicula minoensis</i>				●

References

- Akagi, S., Nakashima, H. and Noto, M., 1992: Sedimentary environments of muddy facies, the Middle Miocene Tototori Group – The facies change of the Fuganji Mudstone Member –. *Memoirs of the Geological Society of Japan*, no. 37, p. 189–199. (in Japanese with English abstract).
- Akiba, F., 1986: Middle Miocene to Quaternary diatom biostratigraphy in the Nankai Trough and Japan Trench, and modified lower Miocene through Quaternary diatom zones for middle-to-high latitudes of the North Pacific. In, Kagami, H., Karig, D.E., Coulbourn, W.T. *et al.*, *Initial Reports of the Deep Sea Drilling Project*, vol. 87, p. 393–480, U.S. Government Printing Office, Washington, D.C.
- Amano, K., Khudik, V.D. and Yokoi, H., 2000: Origin and biogeographic history of *Neilo (Multidentata)* (Bivalvia: Malletiidae). *Venus (Japanese Journal of Malacology)*, vol. 59, no. 3, p. 191–199.
- Amano, K., Hamuro, T., Hamuro, M. and Fujii, S., 2001: The oldest vesicomyid bivalves from the Japan Sea Borderland. *Venus (Journal of the Malacological Society of Japan)*, vol. 60, no. 3, p. 189–198.
- Blow, W.H., 1969: Late middle Eocene to Recent planktonic foraminiferal biostratigraphy. In, Bronnimann, P. and Renz, H.H. eds., *Proceedings of the First International Conference on Planktonic Microfossils (Geneva, 1967)*, vol. 1, p. 199–421, E.J. Brill, Leiden.
- Bouchet, P. and Warén, A., 1985: Revision of the Northeast Atlantic bathyal and abyssal Neogastropoda excluding Turridae (Mollusca, Gastropoda). *Bolletino Malacologico, Supplemento* 1, p. 123–296.
- Bouchet, P. and Warén, A., 1994: Ontogenetic migration and dispersal of deep-sea gastropod larvae. In, Young, C.M. and Eckelbarger, K.J. eds., *Reproduction, Larval Biology, and Recruitment of the Deep-Sea Benthos*, p. 98–117, Columbia University Press, New York.
- Chinzei, K., 1978: Neogene molluscan faunas in the Japanese Islands: An ecologic and zoogeographic synthesis. *The Veliger*, vol. 21, no. 2, p. 155–170.
- Chinzei, K., 1981: Marine biogeography of Japan during Miocene: A reconstruction based on benthonic molluscan faunas. *Fossils (Palaeontological Society of Japan)*, no. 30, p. 7–15. (in Japanese)
- Chinzei, K., 1986: Faunal succession and geographic distribution of Neogene molluscan faunas in Japan. *Palaeontological Society of Japan, Special Papers*, no. 29, p. 161–171.
- Cossmann, M., 1903: *Essais de Paleoconchologie Comparée*, vol. 5, 215 p., Paris.
- Cristofori, J. de and Jan, G., 1832: *Catalogus in IV. Sectiones divisus rerum Naturalium in Museo extantium Josephi de Cristofori et Georgii Jan. Sectio II, pars I. Conchologia*, 16 p., Parma.
- Fischer, P., 1880–1884: *Manuel de conchyliologie et de paléontologie conchologique, on histoire naturelle des mollusques vivants et fossiles*. xxiv+1369 p., Librairie F. Savy, Paris.
- Fujii, S., Kaseno, Y. and Nakagawa, T., 1992: Neogene paleogeography in the Hokuriku region, Central Japan, based on the revised stratigraphic correlation. *Memoirs of the Geological Society of Japan*, no. 37, p. 85–95. (in Japanese with English abstract)
- Fujita, K. and Nakagawa, T., 1948: On the Neogene deposits in the Tonami District, Toyama Prefecture. *Journal of the Geological Society of Japan*, vol. 54, no. 637, p. 125. (in Japanese)
- Gladenkov, A. Yu., 1998: Oligocene and lower Miocene diatom zonation in the North Pacific. *Stratigraphy and Geological Correlation*, vol. 6, no. 2, p. 150–163.
- Hayakawa, H. and Takemura, A., 1987: The Neogene System in Yatsuo area, Toyama Prefecture, central Japan. *Journal of the Geological Society of Japan*, vol. 93, no. 10, p. 717–732. (in Japanese with English abstract)
- Hasegawa, S. and Takahashi, T., 1992: Faunal succession of benthic foraminifera in the upper Yatsuo Group of the Hokuriku District, central Japan – A temporal faunal trend during an early-middle Miocene transgression in Japan. In, Ishizaki, K. and Saito, T. eds., *Centenary of Japanese Micropaleontology*: P. 51–66. Terra Scientific Publishing Company, Tokyo.
- Higo, S., Callomon, P. and Goto, Y., 1999: *Catalogue and Bibliography of the Marine Shell-Bearing Mollusca of Japan*. 749 p., Elle Scientific Publication, Yao.
- Houart, R., 2001: *A Review of the Recent Mediterranean and Northeastern Atlantic Species of Muricidae*. 227 p., Edizioni Evolver, Roma.
- Houart, R. and Lan, T.C., 2003: Description of a new species of *Pagodula* (Gastropoda: Muricidae) from northeast Taiwan. *Occasional Papers of the Malacological Society of Taiwan*, no. 4, p. 38–45.
- Iijima, A. and Tada, R., 1990: Evolution of Tertiary sedimentary basins of Japan in reference to opening of the Japan

- Sea. *Journal of the Faculty of Science, the University of Tokyo, Section 2*, vol. 22, no. 2, p. 121–171.
- Ito, C., Irizuki, T. and Iwai, M., 1999: Diatom zonal key species and geologic ages of the Miocene Morozaki, Iwamura and Tomikusa Groups in the First Setouchi Province, central Japan. *Journal of the Geological Society of Japan*, vol. 93, no. 10, p. 717–732. (in Japanese with English abstract)
- Itoh, Y., Yanagisawa, Y. and Watanabe, M., 1999: Magnetostratigraphy and diatom biostratigraphy of Neogene rocks distributed in the Yatsuo area, central Japan. *Bulletin of the Geological Survey of Japan*, vol. 50, no. 3, p. 215–223. (in Japanese with English abstract)
- Itoigawa, J., 1988: The Miocene Kadonosawa Fauna of Japan. *Saito Ho-on Kai, Special Publication*, no. 2, p. 397–403.
- Itoigawa, J. and Nishikawa, I., 1976: A few problems on the Miocene Setouchi Series in the northern part of Okayama-Hiroshima Prefectures, southwest Japan. *Bulletin of the Mizunami Fossil Museum*, no. 3, p. 127–150, pls. 33–35. (in Japanese with English abstract)
- Itoigawa, J., Shibata, H., Nishimoto, H. and Okumura, Y., 1981: Miocene fossils of the Mizunami Group, central Japan. 2. Molluscs. *Monograph of the Mizunami Fossil Museum*, no. 3-A, p. 1–53, pls. 1–52. (in Japanese)
- Kanehara, K. 1936: Neogene shells from South Chôsen (Korea). *Japanese Journal of Geology and Geography*, vol. 13, nos. 1–2, p. 31–37, pl. 10.
- Kaneko, K. and Goto, M., 1992: Miocene fossils from the Ikuridani, Yatsuo Town, Toyama Prefecture. *Special Publications from the Toyama Science Museum*, no. 5, p. 1–86. (in Japanese)
- Karasawa, H., Nobuhara, T. and Matsuoka, K., 1992: Fossil and living species of the giant isopod genus *Palaega* Woodward, 1870 of Japan. *Science Report of the Toyohashi Museum of Natural History*, no. 2, p. 1–12.
- Karasawa, H., Suzuki, A. and Kato, H., 1995: *Bathynomus undecimspinosus* (Karasawa, Nobuhara & Matsuoka, 1992) (Crustacea, Isopoda) from the Miocene Hidarimatagawa Formation of Southwestern Hokkaido, Japan. *Bulletin of the Mizunami Fossil Museum*, no. 22, 121–125.
- Kaseno, Y., 1964: Biostratigraphical problems of the Neogene strata in Hokuriku Region, central Japan. *Fossils (Paleontological Society of Japan)*, no. 7, p. 27–35. (in Japanese)
- Kasyanov, V.L., Kryuchkova, G.A., Kulikova, V.A. and Medvedeva, L.A., 1990: *Larvae of Marine Bivalves and Echinoderms* (Translated by Smithsonian Institution, 1998). 288 p., Science Publishers, New Hampshire.
- La Perna, R., 1996: Phyletic relationships and ecological implications between *Pagodula vaginata* (De Cristofori & Jan, 1832) and *Pagodula echinata* (Kiener, 1840) (Gastropoda, Muricidae). *Bollettino della Società Paleontologica Italiana*, vol. 35, no. 1, p. 81–92.
- Lee, Y. G., 1992: Paleontological study of the Tertiary molluscan fauna in Korea. *Science Reports of the Institute of Geoscience, University of Tsukuba, Section B*, vol. 13, p. 15–125.
- Locard, A., 1897: Mollusques testacés, tome premier. In, Milne-Edwards, A. ed., *Expédition scientifique du Travailleur et du Talisman pendant les années 1880, 1881, 1882, 1883*. vi+516 p., 22 pls., Masson, Paris.
- Monterosato, A., 1884: *Nomenclatura Generica e Specifica di alcune Conchiglie Mediterranee*. 152 p., Virzi, Palermo.
- Noda, H., Kikuchi, Y. and Nikaido, A., 1989: Paleobiogeographical significance of the first occurrence of the extinct Miocene genus *Acilana* (Mollusca; Bivalvia) from the Pliocene Kume Formation in Ibaraki Prefecture, northern Kanto, Japan. *Professor Hidekuni Matsuo Memorial Volume*, p. 53–64. (in Japanese with English abstract and description)
- Ogasawara, K., 1994: Neogene paleogeography and marine climate of the Japanese Islands based on shallow marine molluscs. *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 108, nos. 3–4, p. 335–351.
- Ogasawara, K., Ijima, S. and Kaseno, Y., 1989: Miocene molluscs from the Tenguyama Formation, Toyama Prefecture, Hokuriku District, Japan. *Science Reports of Kanazawa University*, vol. 34, no. 2, p. 67–93.
- Ogasawara, K. and Nagasawa, K., 1992: Tropical molluscan association in the middle Miocene marginal sea of the Japanese Islands: An example of molluscs from the Oyama Formation, Tsuruoka City, Northeast Honshu, Japan. *Transactions and Proceedings of the Palaeontological Society of Japan, New Series*, no. 167, p. 1224–1246.
- Okada, H. and Bukry, D., 1980: Supplementary modification and introduction of code numbers to the low-latitude coccolith biostratigraphic zonation (Bukry, 1973, 1975). *Marine Micropaleontology*, vol. 5, no. 3, p. 321–325.
- Okamoto, K., 1992: Vertical sequence of molluscan assemblages from the Miocene Bihoku Group in Shobara City, Southwest Japan. – Study of the Bihoku Group IV. *Bulletin of the Mizunami Fossil Museum*, no. 19, p. 319–328. (in Japanese with English abstract)
- Okamoto, K., Katsuhara, M., Ueno, Y. and Sumiyoshi, O., 1990: Molluscan assemblages from the Miocene Bihoku Group in the Kaisekidani area, Miyauchi-cho, Shobara City, Southwest Japan. – Study of the Bihoku Group III. *Bulletin of the Mizunami Fossil Museum*, no. 17, p. 35–49. (in Japanese with English abstract)
- Okamoto, K., Miyanomae, H. and Namikawa, T., 1989: Vertical sequence of the fossil molluscan assemblages from the Miocene Bihoku Group at Negita, Shobara City, Hiroshima Prefecture. – Study of the Bihoku Group. *Professor Hidekuni Matsuo Memorial Volume*, p. 43–52. (in Japanese with English abstract and description)
- Okamoto, K., Suyama, Y., Matsuda, I., Nishimoto, Y. and Kakegawa, K., 1983: The Miocene Susa Group in the north-eastern area of Yamaguchi Prefecture, Japan. *Bulletin of the Mizunami Fossil Museum*, no. 10, p. 85–102. (in Japanese with English abstract)
- Otuka, Y., 1939: Tertiary crustal deformations in Japan (with short remarks on Tertiary paleogeography). *Jubilee Publication in the Commemoration of Prof. H. Yabe 60th Birthday*, vol. 1, p. 481–519.
- Ozawa, T., Nakagawa, T. and Takeyama, K., 1986: Middle Miocene molluscan fauna of the Uchiura Group, Wakasa Province, Southwest Japan. *Palaeontological Society of Japan, Special Papers*, no. 29, p. 135–148, pls. 12–15.
- Radwin, G.E. and D'Attilio, A., 1976: *Murex Shells of the World*. 284 p., 32 pls. Stanford University Press, Stanford.
- Rafinesque, C.S., 1815: *Analyse de la nature, ou tableau de L'univers et des corps organisés*. 224 p., Barravechia, Palermo.
- Riedel, W.R. and Sanfilippo, A., 1978: Stratigraphy and Evolution of tropical Cenozoic radiolarians. *Micropaleontology*, vol. 23, p. 61–96.
- Saito, T., 1988: Molluscan fossils from the Nanamagari Formation (Miocene) in Kanazawa City, Ishikawa Prefecture,

- Japan. *Bulletin of the Mizunami Fossil Museum*, no. 14, p. 139–144, pl. 33. (in Japanese)
- Sakamoto, T. and Nozawa, T., 1960: *Explanatory Text of the Geological Map of Japan, Scale 1:50,000*. “Yatsuo”. 69 p., Geological Survey of Japan, Kawasaki. (in Japanese with English abstract)
- Shibata, H., 1970: Molluscan faunas of the First Setouchi Series, Southwest Japan. Part 1. Fauna of the Ichishi Group. *Journal of Earth Sciences, Nagoya University*, vol. 18, no. 1, p. 27–84, pls. 1–4.
- Shikama, T. and Kase, T., 1976: Molluscan fauna of the Miocene Morozaki Group in the southern part of Chita Peninsula, Aichi Prefecture, southwest Japan. *Science Reports of the Yokohama National University, Sec. 2*, no. 23, p. 1–25, pls. 1–2.
- Shimizu, M., Fujii, S. and Hamuro, T., 2000: Newly found *Aturia* and molluscan fossil assemblages from Higashibessho Formation, Hokuriku Group, Toyama Prefecture, Central Japan. *Earth Science (Chikyu Kagaku)*, vol. 54, p. 43–48 (in Japanese).
- Taguchi, E., 2002: Stratigraphy, molluscan fauna and paleo-environment of the Miocene Katsuta Group in Okayama Prefecture, southwest Japan. *Bulletin of the Mizunami Fossil Museum*, no. 29, p. 95–133, pls. 1–8.
- Tsuchi, R., 1987: Mid-Neogene migration of Tethyan tropical Mollusca and larger Foraminifera into northern Japan. In, McKenzie, K. G., ed., *Proceedings of the International Symposium on Shallow Thetys 2, Wagga Wagga, 15–17 September 1986*, p. 455–459.
- Tsuda, K., 1953: Geology of Yatsuo area in Toyama Prefecture. *Journal of the Faculty of Science, Niigata University, Series 2*, vol. 1, no. 2, p. 1–35. (in Japanese with English abstract)
- Tsuda, K., 1959: New Miocene molluscs from the Kurosedani Formation, in Toyama Prefecture, Japan. *Journal of the Faculty of Science, Niigata University, Series II*, vol. 3, no. 2, p. 67–110, pls. 1–7.
- Tsuda, K., 1960: Paleo-ecology of the Kurosedani fauna. *Journal of the Faculty of Science, Niigata University, Series 2*, vol. 3, no. 4, p. 171–203.
- Tsuda, K. and Chiji, M., 1950: Some problems on the geology in the Yatsuo District. *Journal of Geological Society of Japan*, vol. 56, no. 656, p. 303–304. (in Japanese)
- Van Dover, C.L., 2000: *The Ecology of Deep-Sea Hydrothermal Vents*. 424 p., Princeton University Press, Princeton.
- Yamaoka, M., 1993: The molluscan assemblages from the Miocene Morozaki Group. In, Ohe, F., Nonogaki, I., Tanaka, T., Hachiya, K., Mizuno, Y., Momoyama, T. and Yamaoka, T. eds., *Fossils from the Miocene Morozaki Group*, p. 53–76, Tokai Fossil Society, Nagoya. (in Japanese)
- Yanagisawa, Y. 1999. Diatom biostratigraphy of the lower to middle Miocene sequence in the Yatsuo area, Toyama Prefecture, central Japan. *Bulletin of the Geological Survey of Japan*, vol. 50, no. 3, p. 139–165. (in Japanese with English abstract)
- Yoon, S., 1979: Neogene molluscan fauna of Korea. *Memoir of the Geological Society of China*, no. 3, p. 125–130.