

Consumption of plant material by perch (*Perca fluviatilis*)

Authors: Zapletal, Tomáš, Adámek, Zdeněk, Jurajda, Pavel, Roche, Kevin, Všeticková, Lucie, et al.

Source: Folia Zoologica, 65(2) : 95-97

Published By: Institute of Vertebrate Biology, Czech Academy of Sciences

URL: <https://doi.org/10.25225/fozo.v65.i2.a3.2016>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Consumption of plant material by perch (*Perca fluviatilis*)

Tomáš ZAPLETAL^{1,2,4}, Zdeněk ADÁMEK³, Pavel JURAJDA^{3*}, Kevin ROCHE³, Lucie VŠETIČKOVÁ³ and Jan MAREŠ²

¹ Faculty of Science, University of Hradec Králové, Rokitského 62, 500 03 Hradec Králové, Czech Republic

² Faculty of Agronomy, Mendel University, Zemědělská 1/1665, 613 00 Brno, Czech Republic

³ Institute of Vertebrate Biology, Academy of Sciences of the Czech Republic, v.v.i.

Květná 8, 603 65 Brno, Czech Republic; e-mail: jurajda@brno.cas.cz

⁴ River Elbe Board, s.e., Víta Nejedlého 951, 500 03 Hradec Králové, Czech Republic

Received 29 January 2016; Accepted 4 March 2016

Abstract. While removing fish during reservoir biomanipulation, it was noted that the diet of normally piscivorous 5+ to 7+ perch was dominated by macrophyte fragments, with fish eggs sub-dominant. To the best of our knowledge, macrophytes have not previously been reported as a food item in perch. Here, we briefly discuss this finding and its significance for perch diet studies.

Key words: dietary preference, diet breadth, biomanipulation, accidental feeding

Introduction

While 0+ perch (*Perca fluviatilis*) feed almost exclusively on zooplankton (Adámek et al. 2004, Kratochvíl et al. 2008), juvenile fish (1+) usually switch to other feeding strategies (Persson & Greenberg 1990), with a shift to piscivory/benthivory at 100–150 mm standard length (SL) (Jacobsen et al. 2002) and almost obligatory piscivorous feeding habits at >155 mm SL (Horpilla et al. 2000). In addition to predation on other fish species, perch are also known to consume fish eggs and several studies have examined this feeding behaviour in more detail. Zick et al. (2006), for example, reported up to 168 eggs (67 % of digestive tract content) in one perch at Grundlsee (Austria). As far as we are aware, however, there have been no previous reports of macrophytes as a significant dietary item in normally piscivorous perch. During an ongoing biomanipulation project to improve drinking water quality at a reservoir in the Czech Republic (Jurajda et al. 2014), we observed a surprisingly high percentage of macrophyte fragments in the digestive tracts of adult perch in spring. In this short note, we assess the degree to which macrophytes are consumed by perch in the reservoir and discuss its possible significance.

Material and Methods

This study was carried out at the Hamry drinking water reservoir (49°43'52" N, 15°55'1" E; elevation 603 m

a.s.l.) in the Bohemian-Moravian highlands of the Czech Republic. About half of the 42 ha reservoir's shoreline comprises bankside meadows with a low slope and littoral macrophytes that are flooded during higher spring water levels. The rest of the shoreline comprises coniferous forest with steep to vertical gravel banks with limited or absent vegetation. The inlet area is shallow with soft sediment and a thick layer of detritus from decaying flooded vegetation and littoral macrophyte beds (principally reed canary grass, *Phalaris arundinacea*).

During an ongoing biomanipulation experiment (Jurajda et al. 2014), roach, bream and perch were removed using a 100 m beach seine (max. depth 4 m, mesh 20 mm) and Nordic gillnets in spring (May) and late summer (August and September) of 2012. Of the fish caught, 33 were perch aged 5+ to 7+. These were taken for diet analysis as part of the biomanipulation experiment to assess predation rates on small cyprinids. All fish were weighed to the nearest 0.1 g and measured to the nearest 1 mm using digital callipers. Number, mean SL and mean total biomass (WT) of the perch were relatively balanced between spring and summer – spring n = 20, mean SL 213 (162–273), mean WT 189 (65–342); summer n = 13, mean SL 219 (175–300), mean WT 204 (99–525). After measuring, the fish were dissected and the digestive

* Corresponding Author

tract contents preserved in 4 % formaldehyde for later analysis in the laboratory.

A modified gravimetric method was used to determine diet composition. Mucus and mineral particles were removed from the sample and discarded. All macrophytes and detritus were then separated out from other taxa under a 40× magnification binocular microscope and determined under a 40-450× magnification microscope. Data are presented as relative percentage biomass (% W; Hyslop 1980) and frequency of occurrence (% FO; Pivnička 1981).

All aspects of this study were carried out in accordance with Czech regulations regarding animal care and protection.

Table 1. Relative percentage biomass (% W) and frequency of occurrence (% FO) of main dietary components in the diet of perch collected from the Hamry Reservoir in spring and summer of 2012; spring = May, summer = Aug/Sep.

	Spring % W	Spring % FO	Summer % W	Summer % FO
Zooplankton				
Cladocera	0.09	5.00	1.76	28.57
Invertebrates				
<i>Astacus astacus</i>	-	-	1.44	14.29
Ephemeroptera	0.16	10.00	-	-
Trichoptera	0.47	10.00	3.20	14.29
Diptera	1.04	20.00	-	-
Total Invert.	1.76	45.00	6.40	57.14
Fish	15.91	15.00	93.60	57.14
Fish eggs	28.44	75.00	-	-
Macrophyte	53.89	90.00	-	-

Results and Discussion

While all fish sampled in spring (20) had food in their digestive tracts, six of the 13 fish sampled in summer had empty digestive tracts. There was a clear difference in perch diet between spring and summer (Table 1), with summer diet comprised almost entirely of fish and spring diet dominated by macrophytes (*Alopecurus* sp.) and fish eggs. There was no difference in the quantity of detritus, plankton or aquatic invertebrates taken between spring and summer. These results should be treated with some caution, however, as both the relatively high number of fish with empty tracts in summer and the low number of 5+ to 7+ fish caught overall mean that the data could be easily skewed by outliers. Overall, spring perch ate approximately the same number/biomass of fish as summer perch, the only difference being that summer perch diet was restricted almost

solely to fish while spring perch diet included a large quantity of macrophytes and fish eggs. Hence, while the data intuitively suggest diet switching in perch between spring and summer, i.e. switching to a more profitable (in this case stationary) prey source (fish eggs) for a limited time when their absolute density is high (see Stephens & Krebs 1986), it would be more correct to talk of a widening of the diet spectrum in spring to include fish eggs, and possibly macrophytes. In this sense, the inclusion of eggs and vegetation into perch diet is still of interest.

Fish eggs have occasionally been reported as an important perch dietary item (Zick et al. 2006), their consumption being related to availability during the spring spawning season. To the best of the authors' knowledge, however, macrophytes have not been reported as a significant food item in perch diet. Though fragments of vegetation have frequently been noted, they tend to be found at low frequencies and are usually assumed to have been accidentally consumed, either when preying on aquatic insects or as part of the previous meal of fish prey (Adámek & Sukop 2001, Adámek et al. 2006). As many fish eggs were still attached to fragments of littoral vegetation when removed from the gut, much of the macrophyte would appear to have been taken accidentally. Note, however, that macrophytes were found at higher levels than fish eggs, both by % W and % FO (Table 1). Whether this plant material was actively grazed upon as a separate preferred dietary item, however, or taken accidentally during non-specific grazing for eggs, cannot be judged. Similarly, the degree to which plant material is consumed accidentally when preying on aquatic insects or as part of a fish's previous meal is also unknown. The fact that macrophytes were absent from the diet in summer, when perch returned to an almost exclusively piscivorous diet, suggests that plant consumption is strongly associated with egg consumption.

As the perch digestive system is specialised toward animal nutrition, it is not clear whether the fish gains any nutritional benefit from consuming macrophytes. Indeed, it is possible that large-scale macrophyte consumption (accidental or otherwise) may impact on perch body condition by limiting the presumed profitability of egg predation. For example, fish could potentially reach satiation and cease to feed before reaching a critical energy level, or energy used to digest low-benefit macrophyte material may outweigh any energy gained by eating more profitable fish eggs. In conclusion, our data support the concept of a widening of perch diet in spring in the Hamry

Reservoir to take advantage of a seasonal increase in an easily obtained and profitable prey source (fish eggs). As a result, submerged aquatic macrophytes, whether by accident or intentionally, also become a part of the diet. Further studies are needed to assess whether this pattern is general in large perch or the result of individual specialisation, and to fully assess

the implications of such behaviour on perch body condition.

Acknowledgements

This study was supported through research project ECIP P505/12/G112 of the “European Centre of Ichthyoparasitology”. We would like to thank all the members of the Institute of Vertebrate Biology, Czech Academy of Sciences, who helped with field work.

Literature

- Adámek Z., Jurajda P., Musil J. et al. 2006: Perch (*Perca fluviatilis*) diet during the flooding period of the Chabařovice coal mining pit (North-West Bohemia, Czech Republic). *Proceedings of the 5th International Conference on Reservoir Limnology and Water Quality, Brno, Czech Republic*: 67–68.
- Adámek Z., Musil J. & Sukop I. 2004: Diet composition and selectivity in 0+ perch (*Perca fluviatilis* L.) and its competition with adult fish and carp (*Cyprinus carpio* L.) stock in pond culture. *Agric. Conspec. Sci.* 69: 21–27.
- Adámek Z. & Sukop I. 2001: The role of supplementary feeding in food competition between common carp (*Cyprinus carpio*) and perch (*Perca fluviatilis*) in a pond culture. *Krmiva* 43 (4): 175–184.
- Horpilla J., Ruuhijarvi J., Rask M. et al. 2000: Seasonal changes in the diets and relative abundances of perch and roach in the littoral and pelagic zones of a large lake. *J. Fish Biol.* 56: 51–72.
- Hyslop E.J. 1980: Stomach contents analysis – a review of methods and their application. *J. Fish Biol.* 17: 411–429.
- Jacobsen L., Berg S., Broberg M. et al. 2002: Activity and food choice of piscivorous perch (*Perca fluviatilis*) in a eutrophic shallow lake: a radio-telemetry study. *Freshw. Biol.* 47 (12): 2370–2379.
- Jurajda P., Adámek Z., Janáč M. et al. 2014: Evaluation of five-year effort in biomanipulation measures on the Hamry Reservoir. *The conference “Ecology of Fish in Lakes and Reservoirs” (EcoFiL 2014), 8-11 September 2014, České Budějovice, Czech Republic*: 3.
- Kratochvíl M., Peterka J., Kubečka J. et al. 2008: Diet of larvae and juvenile perch, *Perca fluviatilis* performing diel vertical migrations in a deep reservoir. *Folia Zool.* 57: 313–323.
- Persson L. & Greenberg L.A. 1990: Juvenile competitive bottlenecks: the perch (*Perca fluviatilis*) – roach (*Rutilus rutilus*) interaction. *Ecology* 21: 44–56.
- Pivnička K. 1981: Fish ecology: estimates of basic parameters characterising fish populations, 1st ed. SPN, Praha: 251. (in Czech)
- Stephens D.W. & Krebs J.R. 1986: Foraging theory. *Oxford University Press, Oxford, U.K.*
- Zick D., Gassner H., Jagsch A. & Patzner R.A. 2006: Auswirkung und Populationsentwicklung des eingeschleppten Flussbarsches (*Perca fluviatilis*) im Grundlsee (Steiermark). *Österr. Fisch.* 59: 20–27.