

Bryophyte Diversity Along the Eastern Arc

Author: Pócs, T.

Source: Journal of East African Natural History, 87(1) : 75-84

Published By: Nature Kenya/East African Natural History Society

URL: [https://doi.org/10.2982/0012-8317\(1998\)87\[75:BDATEA\]2.0.CO;2](https://doi.org/10.2982/0012-8317(1998)87[75:BDATEA]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

BRYOPHYTE DIVERSITY ALONG THE EASTERN ARC

T. Pócs¹

Eszterházy Teachers' College, Dept of Botany
Eger, Pf. 43, H-3301, Hungary
colura@gemini.ektf.hu

ABSTRACT

The Eastern Arc Mountains possess a bryoflora with high species diversity (about 700 species are known—more than in Uganda). The proportion of endemism (altogether 32 species, 4.57 %) is not high compared to phanerogams, but is high when compared with the bryoflora of similar areas. Monotypic endemic genera are *Cladolejeunea* and *Neorutenbergia*. *Cladolejeunea aberrans* (Steph.) Zwickel occurs in the Usambaras and *Neorutenbergia usagarae* (Dix.) Biz. et Pócs is distributed throughout the Eastern Arc. The latter is a representative of *Rutenbergiaceae* family with its other members living in Madagascar and on the Mascarene Islands. One notable feature of the bryoflora is the high number (45 species, 6.43 %) of Lemurian (Madagascan) species, which is most apparent in the Uluguru Mountains (40 species, 8.16 %). The bryoflora of the Usambara and Uluguru Mountains is well known. We know much less about the Pare, Nguru and Ukaguru Mountains and the bryoflora of Udzungwa Mountains is practically unexplored.

INTRODUCTION

This paper intends to reveal the characteristic features of the bryophyte flora of Eastern Arc Mountains (figure 1) from a biogeographic perspective. The bryoflora of the Eastern Arc has many similarities with the phanerogamic flora, especially its high diversity. But due to the easy dispersal of the diaspores of cryptogam plants, most bryophytes have a wide distribution, hence their rate of endemism is lower than vascular plants (Zanten & Pócs, 1981; Gradstein & Pócs, 1989). On the other hand, due to the older age of more conservative bryophyte groups and their ability to persist in small patches of suitable habitat, they reflect much better the ancient links between the dissected parts of Gondwanaland (Pócs, 1982). For example, in the Eastern Arc Mountains there are higher number of so-called Lemurian (Madagascan) elements, and of palaeotropical species, than in any other part of the African continent!

¹ Correspondence address: FELSŐTÁRKÁNY, Ady E. u. 67, H-3324, Hungary




Figure 1. The position of Eastern Arc Mountains

RESULTS

Two areas in the Eastern Arc are well known bryologically, namely the Usambaras (see Bizot *et al.*, 1978; Jones, 1972, 1974a, 1974b, 1975, 1976, 1977, 1979, 1981, 1985, 1988; Jones & Pócs, 1987; Ochyra & Pócs, 1985; Pócs, 1982, 1985, 1988, 1992; Vanden Berghen, 1976 and many others; for full lists see O'Shea, 1995 and Wigginton & Grolle, 1996), and the Ulugurus (Bizot & Pócs, 1974, 1980, 1983; Orbán, 1977, 1978; Pócs, 1976a, 1980). For some other mountains (Nguru and Ukaguru Mountains) very rich collections are already under investigation by different specialists, but results are not yet available. For other areas (like Pare and Uzungwa Mountains) only preliminary collections are at hand and their bryoflora is still little known. On the basis of existing records, the bryophyte flora of the Eastern Arc Mountains consists of about 700 species, which represents about 58 % of the very rich Tanzanian bryoflora (1,200 species). For comparison, the whole known Ugandan

bryoflora is some 580 species and even the bryoflora of countries such as Cameroon (584), Nigeria (274) and Ethiopia (404) contain fewer taxa! (Kis, 1985; Wigginton & Grolle, 1996; O'Shea, 1995, 1997).

Endemic bryophytes of the Eastern Arc

The endemics can be subdivided into two groups. There are not many of the so called 'narrow endemics' in the sense of Gradstein & Pócs (1989). They are restricted to a small area within each mountain. For example, *Cololejeunea amaniensis* occurs only in two small areas in the Usambara Mountains: near Amani Station and on Shagein Summit. *Diphyscium pocsii* lives in the Uluguru Mountains: on shady granitic cliffs of the west side ridge of Lupunga peak above Morogoro and in the Mlulu valley above Tangeni village.

The endemic taxa of the different parts of Eastern Arc are outlined below.

a) Endemic bryophytes of the Usambara Mountains:

Anthoceros brunnthaleri Steph.
Anthoceros parvifrons Steph.
Cladolejeunea aberrans (Steph.) Zwickel.
Cololejeunea amaniensis Pócs.
Cololejeunea tanneri Pócs.
Isopterygium baurii Broth.
Philonotis usambarica Broth.
Plagiochila austitexta Steph.
Plagiochila rudolfii Pócs.
Pterobryon julaceum Broth.
Radula pseudoflaccida E.W. Jones.

b) Endemic bryophytes of the Nguru Mountains:

Fissidens pocsii Biz. et Dury.
Fissidens inclusus Biz. et Dury ex Pócs.
Plagiochila hiroshiana Pócs, ined.

c) Endemic bryophytes of the Uluguru Mountains:

Cololejeunea borhidiana Pócs.
Cololejeunea grolleana Pócs.
Cololejeunea jonesii Pócs.
Diphyscium pocsii (Biz.) Zander.
Fissidens hirsutus Biz. ex Pócs.
Hookeriopsis pocsii Biz.
Diplasiolejeunea albifolia ssp *orientalis* Pócs, ined.

d) Endemic bryophytes of the Mufindi Escarpment:

Rhynchostegiella tanneri Biz.
Trichosteleum jonesii Biz.

Subendemic bryophyte species of the Eastern Arc

The subendemic species occur on more than one mountain of the Eastern Arc. There is one monotypic endemic genus, *Neorutenbergia*, represented by *N. usagarae* (Dixon) Bizot et Pócs. This species is widespread over all the montane forests in the Eastern Arc (figure 2), where it occurs as a common epiphyte, forming large festoons on the thinner trunks and branches in the mossy forests between 1,500 and 2,200 m altitude. Several other species (e.g. *Pterobryon flagelliferum* Mitt., *Renauldia lycopodioides* Bizot ex Pócs (figures 3, 4) occur in two or three mountains within the Arc. A disjunct distribution pattern between the

Usambaras and Ulugurus is also found in many other species, which might reflect that these are the larger and most persistently wet mountains.




Figure 2. The known distribution of *Neorutenbergia usagarae* (Dix.) Bizot et Pócs.

Such Eastern Arc subendemics are:

- Fissidens jonesii* Biz (Usambara, Nguru, Ukaguru).
- Hyophila holstii* Broth. (Usambara, Uluguru).
- Leucobryum bistratosum* Broth. (Usambara, Uluguru).
- Mittenothamnium stuhlmannii* (Broth.) Card. (Uluguru + Bukoba).
- Neorutenbergia usagarae* (Dix.) Biz. et Pócs (From Pare to Udzungwa Mountains, see figure 2).
- Pterobryon flagelliferum* Mitt. (Usambara, Nguru, Usagara).
- Renauldia lycopodioides* Biz. ex Pócs (Usambara, Ukaguru, Udzungwa, see figures 3, 4).
- Syrrhopodon stuhlmannii* Broth. (Usambara, Uluguru).
- Syrrhopodon usambaricus* Broth. ex Orbán (Usambara, Uluguru).

Other phytogeographically interesting groups

West and Central African species isolated in East Africa

These species are widespread in West Africa, many of them reaching Central Africa, but they occur only sporadically in East Africa after a large gap. This phenomenon is parallel to the eastern occurrence of many phanerogams, mostly lowland rainforest species, like

Palisota schweinfurthii C.B. Cl. They are relics of a once continuous forest belt from West to East Africa broken up several times during drier climatic periods since Tertiary and latest 12–10,000 BC (Moreau, 1966; Hamilton, 1982). Such western lowland forest bryophytes in East Africa are restricted mostly to the rain exposed slopes of the Eastern Arc Mountains. Such species are:

Archilejeunea abbreviata (Mitt.) Vand. Bergh.

Calymperes intralimbatus C. Müll.

Cololejeunea apiculata (E.W. Jones) Schust.

Cololejeunea harrisii Pócs.

Dendroceros africanus Steph.

Distichophyllidium africanum Dem. et P.Varde.

Ectropothecium brevicladulum (C.M.) Broth.

Fissidens parkii Mitt.

Lepidopilidium devexum Mitt.

Lepidopilum dusenii C. Müll.

Macrohymenium megasporum (Duby) Kis.

Rhaphidorrhynchium brevihorridum (Broth.)

Trichosteleum perhamosum Broth.

And others




Figure 3. *Renauldia lycopodioides* Bizot ex Pócs. a: Habit (drawn by G. Kis); b–g: Leaves (drawn by the author), c: shoulder with auricle, d: apex, e: cells from lamina base, f: median lamina cells, g: apex cells. All drawn from the type.




Figure 4. The distribution of *Renauldia lycopodioides* Bizot ex Pócs.

Palaeotropical species

The Eastern Arc Mountains have a stronger affinity to Asia than the rest of continental Africa. There are several Asian species that reach their westernmost occurrence in the Eastern Arc. The disjunct occurrence of species in African and Asian humid forests could be related to the period in the Miocene when the Tethys Sea was closed and before aridification changed Northern Africa and the Middle East (Axelrod & Raven, 1972). This element of the bryoflora has been presented in detail previously (Pócs, 1976b, 1992), so the following species only serve as examples. Their total number in the Eastern Arc Mountains is 114 or 16.29 %, higher than anywhere else in tropical Africa. Example species are:

Calycularia crispula Mitt.
Cuspidatula contracta (Reinw. *et al.*) Steph.
Dendroceros javanicus (Nees.) Nees
Gottschelia schizopleura (Spruce) Grolle
Homali dendron exiguum (Bosch. & Lac.) Fleisch.
Lejeunea alata Gott.
Pelekium velatum Mitt.
Pleurozia gigantea (F. Web.) Lindb.
 and many others.

Lemurian species

The so-called Lemurian elements in the sense of Tixier (1978) and Pócs (1997) are the species of Madagascar and the other Indian Ocean islands (like the Comoros, Seychelles, Réunion and Mauritius), which may occur only sporadically in mainland Africa. In the Eastern Arc Mountains their occurrence is much more typical among bryophytes than among the vascular plants, reflecting the ancient link that existed before the dissection of Gondwanaland, at least before Madagascar split off from the continent (Pócs, 1975). In certain genera (*Leucobryum*, *Leucoloma*, *Macromitrium*, *Syrrhopodon*—Orbán, 1977, 1978, 1985) they are especially frequent. No doubt, air dispersal has also played a role in this affinity, but the phytogeographical contrast between the young volcanic mountains and the old crystalline massifs in East Africa, as it is shown by Pócs (in Bizot & Pócs, 1974; Pócs, 1975), seems to support the above statement.

As there are a large number of Lemurian species in the Eastern Arc Mountains, the list below is only partial and provides examples of the distribution within the Eastern Arc:

Bryum truncorum (Brid.) Brid.—Uluguru.
Daltonia cardotii Biz. et Onraedt—Uluguru.
Diplasiolejeunea villaumei Steph.—Usambara, Uluguru.
Diplasiolejeunea zakiae Tixier—Taita Hills, Uluguru.
Drepanolejeunea cambouena Steph.—Usambara, Uluguru.
Drepanolejeunea madagascariensis (Steph.) Grolle—Taita Hills, Kasigau, Uluguru.
Frullania usambarana Schiffn.—Usambara, Uluguru.
Holomitrium borbonicum Hampe ex Besch.—Uluguru.
Leucobryum isleanum Besch.—Ethiopia, Kilimanjaro, Usambara, Ukaguru, Uluguru, Zambia.
Lepidozia africana Steph.—Nguru, Uluguru.
Leucoloma brotheri Ren.—Uluguru.
Leucoloma grimmoides P. Varde—Uluguru.
Macromitrium mauritanum Schwaegr.—Usambara, Uluguru.
Pilotrichella perrobusta P. Varde—Ukaguru, Uluguru.
Plagiochila cambouena Steph.—Usambara, Nguru, Mufindi.

Plagiochila drepanophylla Sande-Lacoste—Usambara.

Schlotheimia excorugata C. Müll. ex Card.—Ukaguru, Uluguru.

Schlotheimia robillardii Duby—Usambara.

Sematophyllum sinuosulum (Besch.) Broth.—Taita Hills, Usambara, Uluguru.

Syrrhopodon insularum Bizot ex Onraedt—Uluguru.

CONCLUSIONS

As can be seen from the above (see figure 5 and table 1), the bryoflora of Eastern Arc Mountains can be characterised by a high overall species diversity, a relatively high proportion of endemism compared to other montane bryofloras in Africa and by the high number of Lemurian (Madagascan) species. The last is especially marked in the Uluguru Mountains, where 40 such species occur, which forms 8.16 % of the bryoflora (compared to the 23 such species in the Usambaras and only eight on the volcanic Kilimanjaro). This group of Eastern Arc-Madagascar species, together with the palaeotropical (African-Asian disjunct) element are likely to be biogeographical relics.




Figure 5. The percentage of narrow endemic (occurring only in one mountain area), subendemic (in more than one mountain of the Eastern Arc), Lemurian (reaching the Eastern Arc from Madagascar and Indian Ocean islands), West and Central African (mostly lowland) species in the two well known mountains, compared with those of Kilimanjaro.

Further collections in less well known areas and the identification of previously collected specimens (about 30,000) will almost certainly reveal interesting records and probably taxa new to science, some of them endemic to different Eastern Arc Mountains.

Table 1. Percentage of narrow endemic (occurring only in one mountain area), subendemic (in more than one mountains of the Eastern Arc), Lemurian (reaching the Eastern Arc from Madagascar and Indian Ocean islands), West and Central African (mostly lowland) species in the two well known mountains, compared with those of Kilimanjaro.

	Uluguru Mountains		Usambara		Kilimanjaro	
Total species	490	% of total	464	% of total	569	% of total
Narrow endemics	7	1.43	11	2.37	13	2.28
Eastern Arc subendemics	11	2.24	11	2.37	-	-
Lemurian (Indian Ocean)	40	8.16	23	4.96	8	1.40
West and Central African species	15	3.06	12	2.59	6	1.05

The mossy elfin forests and other rainforests of the Eastern Arc Mountains are renowned for their richness in bryophyte species. But we believe that attention should also be directed towards the almost unknown bryoflora of temporarily dry habitats, like the mist-affected miombo and other woodlands, cliffs and Velloziaceae stands of granitic inselbergs.

ACKNOWLEDGEMENTS

The author is indebted to the Sokoine University of Agriculture (Morogoro, Tanzania), to the Hungarian Academy of Sciences, to the Swedish Agency for Research Cooperation with Developing Countries (SAREC), to the Norwegian Agency for International Development (NORAD) and for the Hungarian Research Fund (OTKA) for their financial assistance during his 25 years research on the bryoflora of East Africa.

REFERENCES

- Axelrod, D.I. & P.H. Raven (1972). Evolutionary biology viewed from plate tectonic theory. In J.A. Behnke (ed.), *Challenging Biological Problems. Directions toward their Solution*. Oxford University Press, New York. Pp 218–236.
- Bizot, M., I. Friis, J. Lewinsky & T. Pócs (1978). East African Bryophytes IV. Danish collections. *Lindbergia* 4: 259–284.
- Bizot, M. & T. Pócs (1974). East African bryophytes I. *Acta Academiae Paedagogicae Agriensis*, n.ser. 12: 383–449.
- Bizot, M. & T. Pócs (1980). East African Bryophytes, III. *Acta Botanica Academiae Scientiarum Hungaricae* 25: 223–261.
- Bizot, M. & T. Pócs (1983). East African Bryophytes, V. *Acta Botanica Academiae Scientiarum Hungaricae* 28: 15–64.
- Gradstein, S.R. & T. Pócs (1989). Bryophytes. In H. Lieth & M.J.A. Werger (eds), *Tropical Rain Forest Ecosystems*. Elsevier, Amsterdam. Pp 311–325.
- Hamilton, A.C. (1982). *Environmental History of East Africa. A Study of the Quaternary*. Academic Press, London and New York.
- Jones, E.W. (1972). African Hepatics XXIII. Some species of *Lejeunea*. *Journal of Bryology* 7: 23–45.
- Jones, E.W. (1974a). African Hepatics XXIV. Lejeuneaceae: some little-known species, and

- extensions of range. *Journal of Bryology* 7: 545–561 “1973”.
- Jones, E.W. (1974b). African Hepatics XXVI. The *Lejeunea eckloniana* complex. *Journal of Bryology* 8: 77–91.
- Jones, E.W. (1975). African Hepatics XXVII. *Bazzania*. *Journal of Bryology* 8: 299–316.
- Jones, E.W. (1976). African Hepatics XXVIII. *Schistochia* Dum. *Journal of Bryology* 9: 33–41.
- Jones, E.W. (1977). African Hepatics XXX. The genus *Radula* Dumortier. *Journal of Bryology* 9: 461–504.
- Jones, E.W. (1979). African Hepatics XXXI. Rare or little-known Lejeuneaceae and extensions of range. *Journal of Bryology* 10: 387–400.
- Jones, E.W. (1981). African Hepatics XXXII. Some little-known species and extensions of range. *Journal of Bryology* 11: 311–323 “1980”.
- Jones, E.W. (1985). African Hepatics XXXIV. Little-known or new Lejeuneaceae. *Journal of Bryology* 13: 385–398.
- Jones, E.W. (1988). African Hepatics XXXVIII. *Cheilolejeunea* subgen. *Strepsilejeunea* (Spruce) Schuster with special reference to East Africa. *Journal of Bryology* 15: 149–160.
- Jones, E.W. & T. Pócs (1987). African Hepatics. XXXVI. Three new species of *Colura*. *Journal of Bryology* 14: 495–501.
- Kis, G. (1985). *Mosses of South-East Tropical Africa. An annotated list with distributional data.* Institute of Ecology and Botany of the Hungarian Academy of Sciences, Vácrátót, Hungary.
- Moreau, R.E. (1966). *The Bird Faunas of Africa and its Islands.* Academic Press, New York and London.
- O’Shea, B. (1995). Checklist of the mosses of sub-Saharan Africa. *Tropical Bryology* 10: 91–198.—with an updated electronic “version 2” distributed through Internet: BBS home page main menu (<http://www.rbge.org.uk.bbs/>) document 51.
- O’Shea, B. (1997). The mosses of sub-Saharan Africa 2. Endemism and biodiversity. *Tropical Bryology* 13: 75–85.
- Ochyra, R., & T. Pócs (1985). East African Bryophytes, VIII. The Musci of the Usambara Rain Forest Expedition, 1982. *Acta Botanica Academiae Scientiarum Hungaricae* 31: 135–146.
- Orbán, S. (1977). Studies on African Calymperaceae I. *Acta Botanica Academiae Scientiarum Hungaricae* 23: 167–177.
- Orbán, S. (1978). Studies on African Calymperaceae II. *Acta Botanica Academiae Scientiarum Hungaricae* 24: 113–120.
- Orbán, S. (1985). Studies on African Calymperaceae, IV. Distribution of the African species of *Syrrhopodon* Schwaegr. *Abstracta Botanica* 9, Suppl. 2: 99–107.
- Pócs, T. (1975). Affinities between the bryoflora of East Africa and Madagascar. In J. Miegé & A. Stork (eds), *Origines des flores Africaines et Malgaches. Nature, spéciation. Comptes rendus de la VIIIe réunion de l’AETFAT, 1. Boissiera* 24a: 125–128.
- Pócs, T. (1976a). Correlations between the tropical African and Asian bryofloras, I. *Journal of the Hattori Botanical Laboratory* 41: 95–106.
- Pócs, T. (1976b). New or little known epiphyllous liverworts I. *Cololejeunea* from tropical Africa *Acta Botanica Academiae Scientiarum Hungaricae* 22: 353–375.
- Pócs, T. (1980). New or little known epiphyllous liverworts, II. Three new *Cololejeunea* from East Africa. *Journal of the Hattori Botanical Laboratory* 48: 305–320.
- Pócs, T. (1982). Examples of the significance of historical factors in the composition of bryofloras. *Beihefte zur Nova Hedwigia* 71: 303–309.
- Pócs, T. (1985). East African Bryophytes, VII. The Hepaticae of the Usambara Rain Forest Project Expedition, 1982. *Acta Botanica Academiae Scientiarum Hungaricae* 31: 113–133.
- Pócs, T. (1988). *Plagiochila rudolfii* sp. nov., from Tanzania, East Africa. *Beihefte zur*

Nova Hedwigia 90: 223–233.

- Pócs, T. (1992). Correlation between the tropical African and Asian bryofloras. II. In T. Koponen & J. Hyvönen (eds), *Proceedings of the Congress of East Asiatic Bryology*, Helsinki, August 12–19, 1990. *Bryobrothera* 1: 35–47.
- Pócs, T. (1997). The distribution and origin of the foliicolous bryophyta in the Indian Ocean Islands. In E. Farkas & T. Pócs (eds), *Cryptogams in the Phyllosphaere: Systematics, Distribution, Ecology and Use*. Proceedings of the IAB & IAL Symposium on Foliicolous Cryptogams, 29 August–2 September 1995, Eger, Hungary. *Abstracta Botanica* 21: 123–134.
- Pócs, T. & B. Tóthmérész (1997). Foliicolous bryophyte diversity in tropical rainforests. In E. Farkas, & T. Pócs (eds), *Cryptogams in the Phyllosphaere: Systematics, Distribution, Ecology and Use*. Proceedings of the IAB & IAL Symposium on Foliicolous Cryptogams, 29 August–2 September 1995, Eger, Hungary. *Abstracta Botanica* 21: 135–144.
- Tixier, P. (1978). La spéciation lemuriennne et les Lejeuneacées. Le cas du genre *Diplasiolejeunea*. In C. Suire, ed., *Congres International de Bryologie*, Bordeaux 21–23 Novembre 1977. *Comptes Rendus, Bryophytorum Bibliotheca* 13: 622–645.
- Vanden Berghen, C. (1976). Frullaniaceae (Hépaticacae) Africanacae. *Bulletin du Jardin Botanique de Belgique* 46: 1–220.
- Wigginton, M.J. & R. Grolle (1996). Catalogue of the Hepaticacae and Anthocerotacae of Sub-Saharan Africa. *Bryophytorum Bibliotheca* 50: 1–267.
- Zanten, B.O. & T. Pócs (1981). Distribution and dispersal of bryophytes. In W. Schultze-Motel (ed.), *Advances in Bryology* 1. J. Cramer, Vaduz. Pp 479–562.