

Spiders from steppe habitats of Pláně Nature Monument (Czech Republic) with suggestions for the local conservation management

Authors: Hamřík, Tomáš, and Košulič, Ondřej

Source: Arachnologische Mitteilungen: Arachnology Letters, 58(1) : 85-96

Published By: Arachnologische Gesellschaft e.V.

URL: <https://doi.org/10.30963/aramit5812>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Spiders from steppe habitats of Pláně Nature Monument (Czech Republic) with suggestions for the local conservation management

Tomáš Hamřík & Ondřej Košulič



doi: 10.30963/aramit5812

Abstract. Spiders from the steppe habitats of the Pláně Nature Monument (Czech Republic) were investigated. The spiders were sampled using pitfall traps and through sweeping of the herb vegetation. In total, 11634 individuals from 154 species in 88 genera from 25 families were recorded. Numerous threatened species listed in the Red List of the Czech spiders were found (CR – 1, EN – 1, VU – 11, LC – 23). 36 mainly xerothermic species were recorded, representing a valuable arachnofauna typical for the Pannonian Region. However, the studied area does not belong to this particular region. Several findings represent the northernmost occurrences of rare thermophilic spiders in the Czech Republic. Ten species (*Dysdera moravica*, *Megalephyphantes pseudocollinus*, *Trichoncus affinis*, *Porrhomma errans*, *Alopecosa striatipes*, *Cheiracanthium oncognathum*, *Civizelotes pygmaeus*, *Zelotes aeneus*, *Thanatus arenarius*, *Talavera aperta*) are discussed in details and their records in the Czech Republic are mapped. Suggestions regarding the management of the area are given to improve the conditions for a high and valuable biodiversity.

Keywords: Araneae, biodiversity, inventory, richness estimation, xerothermic species

Zusammenfassung. Spinnen von Steppenhabitaten des Naturdenkmals Pláně (Tschechische Republik) mit Vorschlägen für das lokale Naturschutzmanagement. Die Spinnen von Steppenhabitaten des Naturdenkmals Pláně (Tschechische Republik) wurden untersucht. Die Erfassung erfolgte mit Bodenfallen und Kescherfängen in der krautigen Vegetation. Insgesamt wurden 11634 Individuen aus 154 Arten, 88 Gattungen und 25 Familien gefangen. Zahlreiche gefährdete Arten der Tschechischen Roten Liste wurden gefunden (CR/vom Aussterben bedroht – 1, EN/stark gefährdet – 1, VU/gefährdet – 11, LC/Vorwarnliste – 23). 36 überwiegend xerotherme Arten wurden erfasst, die zur wertgebenden Spinnenfauna der pannonischen Region gehören – allerdings zählt das Untersuchungsgebiet gar nicht zur pannonischen Region. Zehn Arten (*Dysdera moravica*, *Megalephyphantes pseudocollinus*, *Trichoncus affinis*, *Porrhomma errans*, *Alopecosa striatipes*, *Cheiracanthium oncognathum*, *Civizelotes pygmaeus*, *Zelotes aeneus*, *Thanatus arenarius*, *Talavera aperta*) werden detailliert besprochen und Karten ihrer Nachweise in der Tschechischen Republik präsentiert. Einige der Nachweise sind die nördlichsten Vorkommen seltener thermophiler Arten in der Tschechischen Republik. Es werden Vorschläge zum Management des Gebietes gemacht, um die Bedingungen für eine hohe und wertvolle Biodiversität zu verbessern.

Differences in spider communities can be found even in small-scale patches within a given habitat (Samu et al. 1999, 2011, Pearce et al. 2004). At a community level, spiders are able to respond quickly and distinctly to short or sudden changes in the environment and some species reliably indicate the state of that environment (Buchar 1983, Samu et al. 2011, Košulič et al. 2016). Thus, spiders play an important role in practical nature conservation. The existence of species of conservation value should be taken into account during habitat management processing (Marc et al. 1999, Řezáč et al. 2015).

Since the middle of 20th century, intensification of agriculture, together with an abandonment of less productive lands, has been the major threat to invertebrates in Europe (Tscharntke et al. 2005). Pastures are transformed into a large blocks of fields for crop production or abandonment (Stoate et al. 2009). Lower demand for forage and poor-quality hay are the reasons why traditional grazing and mowing are no longer sustainable (Valkó et al. 2014). In the recent cultivated landscape, grasslands have a significant role in maintaining biodiversity by providing important habitats and refuges for many species (Duelli 1997). Thus, faunistic investigations in these habitats can provide important data useful for evaluation of the biodiversity state in the current landscape. For this reason, this research focuses on the Pláně Nature Monument (NM), an abandoned pasture with xeric grasslands in a south and south-eastern orientation.

Several zoological inventories that were carried out in the study area focused on the following taxa: Orthoptera, Man-

todea, Heteroptera, Hymenoptera, Coleoptera, Lepidoptera, Diptera, Reptilia, Aves and Mammalia (Nenádál 1994). Additionally, a botanical inventory was conducted (Tichý 2016). Spiders have never been studied within the area of the Pláně NM; to date, only several findings from this region are known. Křížová (2001) found 146 species of spiders approximately 1.5 km away from this region, in Libochovka Valley, during an arachnological survey in 1999 to 2000. Since then, no further study was conducted in the surrounding areas of the Pláně NM (ČAS 2019).

The main objective of this research is a faunistic contribution on spiders from the steppe habitats of the Pláně NM, which could be beneficial to the area's conservation management. Consequently, relevant conservation management institutions could take into account these new findings and develop plans based on this research, which would eventually strengthen the protection of local biodiversity.

Material and methods

Study area

The Pláně NM is located in the district of Brno-venkov, 500 m north-west from the village of Kuřimská Nová Ves in the Southern Moravia region of the Czech Republic (Fig. 1, 49.35139°N, 16.30361°E) and belongs to the faunistic square 6663 of the faunistic zoological grid mapping system in the Czech Republic. The protected area has 11.1 ha and is mostly composed of forest and shrubs. The south and south-east parts are composed of a slope with rocks and xeric grasslands on shallow soils (approximately 1.2 ha). The natural forest communities of the study area are oak-hornbeam, acidophilic oak forests, and pine forests on the poor sandy soils with *Festuca ovina*, *Agrostis capillaris*, *Genista tinctoria* and *Luzula campestris*. The geological substrate is characterized by ortorula with

Tomáš HAMŘÍK, Ondřej KOŠULIČ, Department of Forest Protection and Wildlife Management, Faculty of Forestry and Wood Technology, Mendel University in Brno, Zemědělská 3, 613 00 Brno, Czech Republic;
E-mail: hamr.tom@seznam.cz, Ondra.Kosulic@seznam.cz

submitted 31.1.2019, accepted 12.8.2019, online 13.9.2019



Fig. 1: Study area (black square in the map) and sites with rectangles where the pitfall traps were placed and sweeping took place (ESRI 2013)

brown forest soils (Matuška 2016). In general, the Pláně NM lies in a distinctly colder region of South Moravia in the phytogeographical region of Mesophyticum along the border of the South Moravian and Highlands region (Skalický 1988, Slavík 1988, Buchar & Růžicka 2002). The average annual temperature is 8 °C and the average annual rainfall is 579 mm (Matuška 2016). The average altitude is 450 m. Its surrounding landscape is heterogeneous with various habitat types like hedgerows with *Prunus spinosa*, mixed forests and cereal fields. The study area was traditionally maintained by sheep grazing, but there has been no management from 2009 to 2016 (Matuška 2016). Unfortunately, the Pláně NM is currently overgrowing with pine forest, invasive *Robinia pseudoacacia*, pioneer shrub species and expansive *Calamagrostis epigejos*.

Sample collection and study sites

The spiders were collected in two seasons: from 7. May to 15. Oct. 2017 and from 19. Apr. to 21. Sep. 2018. Pitfall traps and sweeping were used as the collection methods for epigeic and herb-dwelling spiders respectively. Each pitfall trap consisted of a plastic cup (500 ml, diameter 9 cm, depth 15 cm) buried in the ground where its rim is level with the soil surface. Each trap was filled with a 3–4% solution of formaldehyde and detergent as a killing and fixative fluid. Sweeping sampling was carried out only during favourable weather conditions, i.e. a minimum of 17 °C and sunshine (period between 10:00 h and 17:30 h). After collecting, the samples were preserved in 70% ethanol. The online application BioLib (2018) was used for the mapping of species occurrence, whereas the distribution data were taken from ČAS (2019).

The following design was used to assess the effect of conservation management methods on spider biodiversity in the study location (Hamřík 2019). Three sites in the study area were selected, wherein each had a total of 16 plots (4 m × 5 m) (Fig. 1). Two pitfall traps were placed in the centre of each plot, making a total of 32 pitfall traps for each site. Sweeping of herb vegetation took place inside each plot (20 sweeps per plot). In this publication, we present only the faunistic results.

Site 1 (452–448 m a.s.l.) and **Site 2** (442–439 m a.s.l.): These slightly steep slopes are characterized by southerly (Site

1) and south-east orientations (Site 2). They are often rocky and herbaceous vegetation is sparse with *Festuca ovina* growing on shallow soil. *Calamagrostis epigejos* and *Arrhenatherum elatius* prevail in nutrient-rich parts. Before 2005, sheep grazing and subsequent mowing with removal of the mown material took place at Site 2.

Site 3 (435–421 m a.s.l.): This site is characterized by a very steep slope with a south-easterly orientation. The lower part of the slope is covered with nutrient-rich soil and dense vegetation, represented by *Calamagrostis epigejos* and *Arrhenatherum elatius*. The upper part of the slope is rocky with sparse herbaceous vegetation, mainly *Festuca ovina*. Several trees and shrubs like *Prunus spinosa* and *Crataegus monogyna* are present at the site.

Species identification and classification

Using the identification keys of Heimer & Nentwig (1991) and Nentwig et al. (2019), only adult spiders were used and determined down to species level. Every species was categorized according to hunting strategies (Fig. 2): sensing-web weavers, sheet-web weavers, space-web weavers, orb-web weavers, ambush hunters, other hunters, ground hunters and specialists (Cardoso et al. 2011). The most up-to-date data on nomenclature were taken from the World Spider Catalog (WSC 2019) and the arrangement of families is based on the spider phylogenetic tree of Wheeler et al. (2017). The authors determined almost all species; taxonomically complicated taxa were revised and determined by another specialist (Petr Dolejš, National Museum in Prague). Each species was evaluated according to Buchar & Růžicka (2002) and Řezáč et al. (2015) (Tab. 1): 1) occurrence level, 2) habitat preferences, 3) thermo preferences, 4) conservation status. Occurrence level is categorized according to the abundance based on the estimated number and distribution of grid squares with potential species occurrence (the presence of suitable habitats): very abundant (VA), abundant (A), scarce (S), rare (R), very rare (VR). Another evaluation is based on habitat preferences according to degree of habitat originality: climax preferences (C), seminatural habitats (SN), disturbed (D), artificial (A). Thermo preference is categorized according to occurrence of spider species in phytogeographic districts: Thermophyticum (T), Mesophyticum (M), Oreophyticum (O). The last evaluation is the conservation status which is based on the Red List categories: critically endangered (CR), endangered (EN), vulnerable (VU), least concern (LC). The basic values are written in regular font; the markedly preferred values are in bold; and some marginal but non-negligible values are in parentheses (see Tab. 1).

Species richness estimation

For estimation purposes, if the absolute species number in the 48 samples thoroughly represented alpha diversity of the study area, Jackknife1 resampling with 100 randomizations using the Software EstimateS 9.1 was performed (Colwell 2013).

Results and discussion

Faunistic overview

Overall, 11634 adult spiders belonging to 154 species in 88 genera of 25 families were collected (Tab. 1). Altogether, 133 species from 10532 specimens (91%) were caught by pitfall traps and 49 species from 1102 specimens (9%) were caught




Fig. 2: Guild composition of spiders in steppe habitats of the Pláně NM

by sweeping. Spiders collected by pitfall trapping consisted mainly of actively hunting guilds such as ground and other hunters and several species of specialists from the family Dysderidae (Fig. 2).

The most abundant species were *Alopecosa cuneata* with 2603 individuals and *Pardosa palustris* with 1884 individuals. The next dominant species were *Alopecosa farinosa* with 537 individuals and *Alopecosa pulverulenta* with 487 individuals. The aforementioned spiders of the genus *Alopecosa* are typical for open steppe habitats (Buchar & Růžička 2002). *Alopecosa striatipes*, which is very rare in the Czech Republic and occurs only in the south-eastern part of the country (Kůrka et al. 2015), was caught from a high number of 387 individuals, mostly in Site 1 and can be considered abundant in the Pláně NM. The next result was the significantly high abundance of *Thanatus arenarius*, which is rare and only found in warm areas of central and south-eastern parts of the Czech Republic (Buchar & Růžička 2002, Košulič & Hula 2014). However, in the study area, a high number of 704 individuals were found mostly at Site 2. Both Sites 1 and 2 are typical of sparse herbaceous vegetation and stones/rocks on shallow soil, thus meeting the ecological requirements of these species (Ratschker & Roth 2000).

The family Linyphiidae was the most dominant with 32 species recorded (Tab. 1). However, only a few individuals of each species were caught. *Agyneta rurestris* was found to be quite common due to the non-restriction of ecological requirements and aerial dispersal using silk. The next species-rich families were Gnaphosidae (22 species), Lycosidae (14 species), Salticidae (12 species) and Theridiidae (12 species). The suitability for xerothermic herb-dwelling spiders is corroborated by the high abundance of 149 individuals of the scarce *Evarcha laetabunda*, which inhabits herbs on rock steppes (Buchar & Růžička 2002). The Lycosidae with 7104 and Gnaphosidae with 1367 individuals were the most abundant families. From the family Lycosidae, there were also species that are typical of forest edges (Buchar & Růžička 2002): *Pardosa lugubris* and *Trochosa terricola* were abundantly found at Site 3, which is adjacent to a forest.

At Site 3, the xerothermic species *Alopecosa trabalis*, which inhabits forest steppes and light south-exposed forests (Buchar & Růžička 2002), was also abundant (260 individuals). Site 3 is a good example of a xerothermic slope habi-

tat with a south-eastern exposure that is adjacent to a forest, hence it provides suitable conditions for both typical steppe species and species typical of ecotones like forest edges. In the family Gnaphosidae there were a number of regionally important rare/scarcely species like *Civizelotes pygmaeus*, *Drassyllus pumilus*, *Haplodrassus dalmatensis*, *Micaria formicaria*, *Zelotes aeneus*, *Zelotes aurantiacus*, *Zelotes electus* and *Zelotes longipes*, which all live under stones on rock steppes (Buchar & Růžička 2002). Their presence is due to the large amount of scattered small rocks and stones which serve as shelter for these ground hunters. For the record, these species are mostly found in their northernmost territory of distribution in Southern Moravia. *Zelotes electus* was abundantly found at Site 1 (84 individuals) and Site 2 (86 individuals), whereas *Zelotes aurantiacus* was found at Site 3 (24 individuals). Site 2 had a xeric character while Site 3 had a lower amount of sparse vegetation and was adjacent to the forest. Both species have similar habitat preferences, therefore, it may be an effect of competition like that of other related species (Michalko et al. 2016).

In general, a valuable arachnofauna composition including 36 rare and scarce species (7 %) belonging to the Red List of Czech spiders (Řezáč et al. 2015) were discovered. It includes mainly species whose occurrence is typical of open steppes and grasslands at the early stages of succession: CR (critically endangered): *Alopecosa striatipes*; EN (endangered): *Drassyllus pumilus*; VU (vulnerable): *Neottiura suaveolens*, *Trichoncus affinis*, *Trichopterna cito*, *Cheiracanthium oncognathum*, *Phrurolithus minimus*, *Haplodrassus dalmatensis*, *Micaria formicaria*, *Civizelotes pygmaeus*, *Thanatus arenarius*, *Evarcha laetabunda*, *Talavera petrensis*.

It should be noted that the total spider diversity (154 species) was relatively high and this significantly enriches the surveyed area due to new findings. Altogether, 230 species of spiders are now known from the study area (faunistic square 6663) due to the presented faunistic study and previously published data (Křížová 2001). The occurrence of a wide spectrum of species also confirms the importance of xeric grasslands within the Pláně NM as a refuge for spider communities in the intensified agricultural landscape of Moravia. These stepping stones are very important in the overall biodiversity of arthropods in the modern landscape (Košulič et al. 2014, Šálek et al. 2018). The results also suggest that

Tab. 1: List of recorded species with ecological indicators, in taxonomical order. Occurrence level: VA (very abundant), A (abundant), S (scarce), R (rare), VR (very rare); habitat preference: C (climax), SN (seminatural), D (disturbed), A (artificial); thermo preference: T (thermophyticum), M (mesophyticum), O (oreophyticum); conservation status: CR (critically endangered), EN (endangered), VU (vulnerable), LC (least concern)

Species	Site 1	Site 2	Site 3	Occurrence level	Habitat preferences	Thermo preferences	Conservation status
Segestriidae							
<i>Segestria senoculata</i> (Linnaeus, 1758)	.	.	2	VA	C, SN	(T), M, (O)	–
Dysderidae							
<i>Dysdera moravica</i> Řezáč, 2014	.	1	1	R	C	T	LC
<i>Harpactea hombergi</i> (Scopoli, 1763)	.	.	1	A	C, SN	T, M	–
<i>Harpactea lepida</i> (C. L. Koch, 1838)	2	.	2	VA	C, SN	M , (O)	–
<i>Harpactea rubicunda</i> (C. L. Koch, 1838)	1	.	2	VA	C, SN, A	T, M	–
Theridiidae							
<i>Asagena phalerata</i> (Panzer, 1801)	10	3	8	A	C, SN	(T), M , (O)	–
<i>Crustulina guttata</i> (Wider, 1834)	.	.	1	A	C, SN	M	–
<i>Enoplognatha latimana</i> Hippa & Oksala, 1982	3	1	1	S	SN, D	T, M	–
<i>Enoplognatha thoracica</i> (Hahn, 1833)	2	3	2	A	C, SN, D	T, M	–
<i>Euryopsis flavomaculata</i> (C. L. Koch, 1836)	1	.	.	A	C, SN	T, M	–
<i>Lasaeola tristis</i> (Hahn, 1833)	1	.	.	S	C, SN	M	LC
<i>Neottiura bimaculata</i> (Linnaeus, 1767)	3	.	.	VA	C, SN, D	T, M	–
<i>Neottiura suaveolens</i> (Simon, 1879)	2	.	.	R	C	T	VU
<i>Phylloneta impressa</i> L. Koch, 1881	37	47	5	VA	C, SN, D	T, M, (O)	–
<i>Robertus arundineti</i> (O. P.-Cambridge, 1871)	2	.	1	A	C, SN, D	(T), M	–
<i>Robertus lividus</i> (Blackwall, 1836)	5	.	.	VA	C, SN	T, M, O	–
<i>Robertus neglectus</i> (O. P.-Cambridge, 1871)	1	.	.	S	C, SN	(T), M	–
Linyphiidae							
<i>Agyneta rurestris</i> (C. L. Koch, 1836)	6	3	8	VA	C, SN, D	T, M, O	–
<i>Agyneta saxatilis</i> (Blackwall, 1844)	2	3	2	VA	C, SN, (D)	M	–
<i>Ceratinella brevis</i> (Wider, 1834)	1	2	.	VA	C, SN	M, (O)	–
<i>Ceratinella scabrosa</i> (O. P.-Cambridge, 1871)	3	1	3	S	C, SN	M	–
<i>Diplocephalus cristatus</i> (Blackwall, 1833)	2	1	5	VA	C, SN, (D)	M , (O)	–
<i>Diplocephalus picinus</i> (Blackwall, 1841)	1	1	.	VA	C, SN	(T), M, (O)	–
<i>Diplostyla concolor</i> (Wider, 1834)	2	.	.	VA	C, SN	T, M, O	–
<i>Entelecara acuminata</i> (Wider, 1834)	2	.	.	A	C, SN	M	–
<i>Erigone atra</i> Blackwall, 1833	1	2	.	VA	C, SN, D	T, M, O	–
<i>Erigone dentipalpis</i> (Wider, 1834)	.	2	1	VA	C, SN, D	T, M, O	–
<i>Linyphia hortensis</i> Sundevall, 1830	1	1	.	A	C, SN	(T), M	–
<i>Linyphia triangularis</i> (Clerck, 1757)	1	.	2	VA	C, SN, D	T, M	–
<i>Megalepthyphantes pseudocollinus</i> Saaristo, 1997	.	.	1	VR	C	T	LC
<i>Micrargus herbigradus</i> (Blackwall, 1854)	1	1	1	VA	C, SN	(T), M , O	–
<i>Microlinyphia pusilla</i> (Sundevall, 1830)	1	.	.	VA	C, SN, D	T, M, O	–
<i>Neriere emphana</i> (Walckenaer, 1841)	.	1	.	A	C, SN	M	–
<i>Neriere radiata</i> (Walckenaer, 1841)	.	1	.	A	C, SN	M	–
<i>Palliduphantes pallidus</i> (O. P.-Cambridge, 1871)	5	3	1	VA	C, SN	T, M	–
<i>Pelecopsis elongata</i> (Wider, 1834)	4	1	.	S	C	M	LC
<i>Pelecopsis radicola</i> (L. Koch, 1872)	2	1	1	A	C, SN	M , (O)	–
<i>Porrhomma errans</i> (Blackwall, 1841)	3	.	.	VR	C, D	T	–
<i>Stemonyphantes lineatus</i> (Linnaeus, 1758)	3	.	1	A	C, SN, D	(T), M	–
<i>Tapinocyba insecta</i> (L. Koch, 1869)	1	.	.	A	C, SN	(T), M	–
<i>Tenuiphantes cristatus</i> (Menge, 1866)	.	.	1	VA	C, SN	M , (O)	–
<i>Tenuiphantes flavipes</i> (Blackwall, 1854)	5	6	3	VA	C, SN	T, M	–
<i>Tenuiphantes mendei</i> Kulczyński, 1887	1	.	.	VA	C, SN	T, M, O	–
<i>Tenuiphantes tenuis</i> (Blackwall, 1852)	2	2	.	A	C, SN, D	T, M	–
<i>Tiso vagans</i> (Blackwall, 1834)	1	2	2	A	C, SN, (D)	M , (O)	–
<i>Trichoncus affinis</i> Kulczyński, 1894	.	.	1	R	C, SN	M	VU
<i>Trichopterna cito</i> (O. P.-Cambridge, 1872)	1	.	.	S	C	T , M	VU

Species	Site 1	Site 2	Site 3	Occurrence level	Habitat preferences	Thermo preferences	Conservation status
<i>Walckenaeria antica</i> (Wider, 1834)	1	.	.	VA	C, SN	(T), M, (O)	–
<i>Walckenaeria atrotibialis</i> (O. P.-Cambridge, 1878)	1	1	.	VA	C, SN	T, M, O	–
<i>Walckenaeria dysderoides</i> (Wider, 1834)	1	.	.	VA	C, SN	(T), M	–
Tetragnathidae							–
<i>Metellina segmentata</i> (Clerck, 1757)	.	.	1	VA	C, SN, D	T, M, O	–
<i>Pachygnatha degeerii</i> Sundevall, 1830	.	1	12	VA	C, SN, D	T, M, (O)	–
<i>Pachygnatha listeri</i> Sundevall, 1830	.	.	1	VA	C, SN	(T), M	–
Araneidae							
<i>Aculepeira ceropegia</i> (Walckenaer, 1802)	45	65	26	VA	C, SN, D	(T), M	–
<i>Araneus sturmi</i> (Hahn, 1831)	.	1	.	VA	C, SN	T, M	LC
<i>Argiope bruennichi</i> (Scopoli, 1772)	9	2	2	A	C, SN, D	T, M	–
<i>Cercidia prominens</i> (Westring, 1851)	1	.	.	S	C, SN	T, M	–
<i>Hypsosinga albovittata</i> (Westring, 1851)	18	44	26	S	C, SN	T, M	LC
<i>Hypsosinga pygmaea</i> (Sundevall, 1831)	1	.	.	S	C, SN	M	LC
<i>Hypsosinga sanguinea</i> (C. L. Koch, 1844)	10	11	1	A	C, SN	(T), M	–
<i>Mangora acalypha</i> (Walckenaer, 1802)	164	130	245	VA	C, SN, D	T, M	–
Titanoecidae							
<i>Titanoeca quadriguttata</i> (Hahn, 1833)	.	.	1	A	C, SN	T, M	–
Lycosidae							
<i>Alopecosa cuneata</i> (Clerck, 1757)	1374	663	566	VA	C, SN, D	T, M, (O)	–
<i>Alopecosa farinosa</i> (Latreille, 1817)	183	219	135	A	C, SN	T, M	–
<i>Alopecosa pulverulenta</i> (Clerck, 1757)	150	65	272	VA	C, SN, D	T, M, O	–
<i>Alopecosa striatipes</i> (C. L. Koch, 1837)	191	122	74	VR	C	T	CR
<i>Alopecosa trabalis</i> (Clerck, 1757)	59	15	186	S	C, SN	T, M	–
<i>Aulonia albimana</i> (Walckenaer, 1805)	16	1	52	A	C, SN	T, M	–
<i>Pardosa lugubris</i> (Walckenaer, 1802)	14	1	188	VA	C, SN, D	T, M, O	–
<i>Pardosa palustris</i> (Linnaeus, 1758)	394	1186	304	VA	C, SN, D	T, M, O	–
<i>Pardosa pullata</i> (Clerck, 1757)	22	22	6	VA	C, SN, D	T, M, O	–
<i>Pardosa riparia</i> (C. L. Koch, 1833)	77	42	144	A	C, SN	T, M, O	–
<i>Trochosa ruricola</i> (De Geer, 1778)	20	5	55	VA	C, SN, D	T, M	–
<i>Trochosa terricola</i> Thorell, 1856	42	9	138	VA	C, SN, D	T, M, (O)	–
<i>Xerolycosa miniata</i> (C. L. Koch, 1834)	13	41	26	S	C, SN	T, M	–
<i>Xerolycosa nemoralis</i> (Westring, 1861)	5	5	3	VA	C, SN	T, M, O	–
Pisauridae							–
<i>Pisaura mirabilis</i> (Clerck, 1757)	8	5	19	VA	C, SN, D	T, M	–
Oxyopidae							
<i>Oxyopes ramosus</i> (Martini & Goeze, 1778)	.	.	1	S	C, SN	M	LC
Thomisidae							
<i>Misumena vatia</i> (Clerck, 1757)	1	.	2	VA	C, SN	T, M	–
<i>Ozyptila atomaria</i> (Panzer, 1801)	2	2	3	S	C, SN	T, M	–
<i>Ozyptila clavata</i> (Walckenaer, 1837)	32	18	24	S	C	T, M	LC
<i>Ozyptila praticola</i> (C. L. Koch, 1837)	.	.	1	S	C, SN	T, M	–
<i>Synema globosum</i> (Fabricius, 1775)	2	1	5	R	C, SN	T, M	LC
<i>Xysticus bifasciatus</i> C. L. Koch, 1837	32	18	28	VA	C, SN, D	(T), M, (O)	–
<i>Xysticus cristatus</i> (Clerck, 1757)	75	62	36	VA	C, SN, D	T, M, (O)	–
<i>Xysticus erraticus</i> (Blackwall, 1834)	9	.	2	A	C, SN	(T), M	–
<i>Xysticus kochi</i> Thorell, 1872	38	76	47	A	C, SN, (D)	T, M	–
<i>Xysticus lanio</i> C. L. Koch, 1835	.	.	2	S	C, SN	T, M	–
Agelenidae							
<i>Allagelena gracilis</i> (C. L. Koch, 1841)	1	.	2	A	C, SN, (A)	T, M	–
<i>Coelotes terrestris</i> (Wider, 1834)	3	1	8	VA	C, SN	(T), M, O	–
<i>Histopona torpida</i> (C. L. Koch, 1834)	.	1	2	VA	C, SN	M, (O)	–
<i>Inermocoelotes inermis</i> (L. Koch, 1855)	4	14	73	VA	C, SN	M, O	–

Species	Site 1	Site 2	Site 3	Occurrence level	Habitat preferences	Thermo preferences	Conservation status
<i>Tegenaria campestris</i> C. L. Koch, 1834	2	.	6	S	C, SN	T, M	–
<i>Tegenaria silvestris</i> L. Koch, 1872	1	.	.	A	C, SN	M, (O)	–
Cybaeidae							
<i>Cybaeus angustiarum</i> L. Koch, 1868	.	.	6	A	C, SN	M, O	–
Hahniidae							
<i>Habnia nava</i> (Blackwall, 1841)	.	3	5	S	C, SN	T, M	–
Dictynidae							
<i>Cicurina cicur</i> (Fabricius, 1793)	18	8	8	VA	C, SN, D	(T), M	–
<i>Dictyna arundinacea</i> (Linnaeus, 1758)	8	1	4	VA	C, SN, D	(T), M	–
<i>Nigma flavescens</i> (Walckenaer, 1830)	1	1	.	A	C, SN	T, M	–
Amaurobiidae							
<i>Callobius claustrarius</i> (Hahn, 1833)	.	2	7	A	C, SN	M, O	–
Liocranidae							
<i>Agroeca brunnea</i> (Blackwall, 1833)	6	2	6	VA	C, SN	T, M	–
<i>Agroeca cuprea</i> Menge, 1873	20	18	27	S	C	T, M	LC
<i>Agroeca proxima</i> (O. P.-Cambridge, 1871)	9	3	1	S	C, SN	M, O	–
Clubionidae							
<i>Clubiona neglecta</i> O. P.-Cambridge, 1862	1	5	1	VA	C, SN	(T), M	–
<i>Clubiona terrestris</i> Westring, 1851	.	.	2	VA	C, SN	M	–
Phrurolithidae							
<i>Phrurolithus festivus</i> (C. L. Koch, 1835)	.	.	1	VA	C, SN	T, M	–
<i>Phrurolithus minimus</i> C. L. Koch, 1839	1	.	.	R	C, SN	T, M	VU
Gnaphosidae							
<i>Civizelotes pygmaeus</i> (Miller, 1943)	1	3	6	R	C	T	VU
<i>Drassodes lapidosus</i> (Walckenaer, 1802)	1	2	14	VA	C, SN	T, M	–
<i>Drassodes pubescens</i> (Thorell, 1856)	62	45	31	VA	C, SN	T, M	–
<i>Drassyllus lutetianus</i> (L. Koch, 1866)	1	.	.	A	C, SN, D	(T), M	–
<i>Drassyllus praeficus</i> (L. Koch, 1866)	7	15	14	A	C, SN	T, M	–
<i>Drassyllus pumilus</i> (C. L. Koch, 1839)	3	1	.	R	C	T, M	EN
<i>Drassyllus pusillus</i> (C. L. Koch, 1833)	55	43	30	A	C, SN, (D)	T, M	–
<i>Haplodrassus dalmatensis</i> (L. Koch, 1866)	1	2	1	R	C	T	VU
<i>Haplodrassus signifer</i> (C. L. Koch, 1839)	106	106	111	VA	C, SN, D	T, M, O	–
<i>Haplodrassus silvestris</i> (Blackwall, 1833)	1	2	2	A	C, SN	(T), M	–
<i>Haplodrassus soerenseni</i> (Strand, 1900)	.	.	1	S	C, SN	M	LC
<i>Micaria formicaria</i> (Sundevall, 1831)	21	7	14	R	C, SN	T, (M)	VU
<i>Micaria fulgens</i> (Walckenaer, 1802)	9	4	34	A	C, SN	T, M	LC
<i>Micaria pulicaria</i> (Sundevall, 1831)	.	1	6	VA	C, SN	T, M, O	–
<i>Scotophaeus quadripunctatus</i> (Linnaeus, 1758)	1	.	.	S	C, SN, A	–	–
<i>Trachyzelotes pedestris</i> (C. L. Koch, 1837)	4	2	16	S	C, SN	T, (M)	–
<i>Zelotes aeneus</i> (Simon, 1878)	9	6	2	R	C, SN, D	(T), M	LC
<i>Zelotes aurantiacus</i> Miller, 1967	1	1	24	R	C	T	LC
<i>Zelotes electus</i> (C. L. Koch, 1839)	84	86	9	S	C, SN	T, M	LC
<i>Zelotes latreillei</i> (Simon, 1878)	8	3	4	VA	C, SN, D	(T), M	–
<i>Zelotes longipes</i> (L. Koch, 1866)	6	100	2	R	C	T, (M)	LC
<i>Zelotes petrensis</i> (C. L. Koch, 1839)	71	75	93	A	C, SN	T, M	–
Miturgidae							
<i>Zora nemoralis</i> (Blackwall, 1861)	1	1	.	A	C, SN	(T), M	–
<i>Zora silvestris</i> Kulczyński, 1897	3	.	6	A	C, SN	M	–
<i>Zora spinimana</i> (Sundevall, 1833)	4	2	5	VA	C, SN, D	T, M, (O)	–
Cheiracanthiidae							
<i>Cheiracanthium campestre</i> Lohmander, 1944	.	1	.	VR	SN	T	LC
<i>Cheiracanthium oncognathum</i> Thorell, 1871	1	.	.	R	C, SN	T, M	VU

Species	Site 1	Site 2	Site 3	Occurrence level	Habitat preferences	Thermo preferences	Conservation status
Sparassidae							
<i>Micrommata virescens</i> (Clerck, 1757)	.	.	1	VA	C, SN	M	–
Philodromidae							
<i>Philodromus albidus</i> Kulczyński, 1911	.	.	1	A	C, SN, D	T, M	–
<i>Philodromus cespitum</i> (Walckenaer, 1802)	.	2	.	VA	C, SN, D	T, M	–
<i>Philodromus dispar</i> Walckenaer, 1826	1	.	1	S	C, SN	T, M	–
<i>Thanatus arenarius</i> Thorell, 1872	318	362	24	R	C	T	VU
<i>Thanatus formicinus</i> (Clerck, 1757)	101	39	34	A	C, SN	T, M	LC
<i>Thanatus striatus</i> C. L. Koch, 1845	1	.	.	A	C, SN	(T), M	LC
<i>Tibellus oblongus</i> (Walckenaer, 1802)	1	.	.	S	C, SN	T, M	–
Salticidae							
<i>Euophrys frontalis</i> (Walckenaer, 1802)	.	1	.	A	C, SN	T, M	–
<i>Euophrys petrensis</i> C. L. Koch, 1837	5	4	3	S	C, SN	T, M	VU
<i>Evarcha arcuata</i> (Clerck, 1757)	3	1	7	VA	C, SN	T, M	–
<i>Evarcha laetabunda</i> (C. L. Koch, 1846)	68	42	39	S	C	T, (M)	VU
<i>Heliophanus cupreus</i> (Walckenaer, 1802)	1	1	2	A	C, SN	T, M	–
<i>Heliophanus flavipes</i> (Hahn, 1832)	12	6	17	A	C	(T), M	–
<i>Hypositticus pubescens</i> (Fabricius, 1775)	1	.	.	VA	C, SN, A	M	–
<i>Pellenes tripunctatus</i> (Walckenaer, 1802)	.	1	.	S	C	T	LC
<i>Phlegra fasciata</i> (Hahn, 1826)	34	21	35	A	C, SN	T, M	–
<i>Sibianor aurocinctus</i> (Ohlert, 1865)	1	.	.	A	C, SN	T, M	LC
<i>Talavera aequipes</i> (O. P.-Cambridge, 1871)	3	3	2	A	C, SN	T, M	–
<i>Talavera aperta</i> (Miller, 1971)	1	.	.	R	C, SN	T, M	LC

the study area serves as an important refuge for thermophilic spiders that are normally not present in this otherwise colder mesophytic region. This is supported by the high occurrence of typically Pannonian species, for whom this area represents their northernmost distribution limit (Buchar & Růžička 2002, Kůrka et al. 2015).

Species richness estimation

The species accumulation curve did not reach the asymptote and showed a rising character, which means that the spider diversity was not sampled in their entirety and the diversity is expected to be significantly higher than the 154 species collected. Jackknife 1 estimator calculated a total estimated richness of 194.15 ± 6.26 (standard deviation) species for the study area (Fig. 3).

Faunistically remarkable species

All the species mentioned below belong to rare, xerothermic species and the Pláně NM is their northernmost distribution not only in South Moravia, but even in Moravia as a whole. Aside from the *Thanatus arenarius*, the following findings are the first for the faunistic square 6663.

Dysdera moravica (Fig. 4 a)

This species belongs to a large complex of morphologically similar species which mainly occur in northern Italy and in the northwest of the Balkan Peninsula (Řezáč et al. 2012). *Dysdera moravica* is European species that occurs in Serbia, Croatia, Romania, Hungary, Slovakia, Austria, Germany and the Czech Republic (Nentwig et al. 2019). In the Czech Re-




Fig. 3: Number of species in 48 samples in relation to the number of species expected by a Jackknife1 estimator in the Pláně NM

public, this species occurs only in the Pannonian region and its close surroundings. This is a xerothermic species that live in areas with rugged terrain on south oriented slopes with xerothermic oak forests, oak-hornbeam forests or bushes (Řezáč 2012, Košulič 2017).

Location: Two individuals were captured in pitfall traps. At Site 2, one individual was caught in sparse vegetation and another individual was caught on bare soil at Site 3.

Megalepthyphantes pseudocollinus (Fig. 4 b)

In Europe, the species was found in Hungary, Austria, Czech Republic, Slovakia, Ukraine, Belarus, Russia (Southern European, Central European, Western European, Eastern European) and Finland, outside Europe it occurs in Russia (West Siberia), the Caucasus, Turkey and Iran (Nentwig et al. 2019). Within the Czech Republic, it has been mapped from only a few areas of Pannonian regions (Buchar & Růžicka 2002, Bryja et al. 2005). It was discovered in semixerix habitats in Brtnice-Střížov, the most western area of the species' occurrence in the Czech Republic (Svatoň & Jelinek 1998). It is a very rare species that inhabits xerothermic habitats like stony debris in steppes (Bryja et al. 2005).

Location: One individual of the species was captured in a pitfall trap in mesophilic vegetation at Site 3.

Porrhomma errans (Fig. 4 d)

This species has a north-western and central European distribution and was also recorded in Italy (Růžicka 2018, WSC 2019). It is a rare species with an unclear ecological requirement (Kůrka et al. 2015, BAS 2019). *Porrhomma errans* is known from different habitats like beet fields, steppe habitats or floodplain forests (Miller 1974, Bryja et al. 2005, Hula et al. 2012).

Location: The species was sampled by sweeping at Site 1. Three individuals were sampled both in sparse and dense vegetation.

Alopecosa striatipes (Fig. 4 e)

Alopecosa striatipes is an extra-Mediterranean faunal element with a distribution area that extends from the Atlantic coast (Buchar & Thaler 2004). In Europe, the species occurs in Macedonia, Serbia, Montenegro, Croatia, Slovenia, Bulgaria, Romania, Ukraine, Slovakia, Czech Republic, Poland, Germany, Belgium, Switzerland and France, outside Europe the species was recorded in Turkey and the Caucasus (Nentwig et al. 2019). The species is known very rarely from the south-eastern part of the Czech Republic (Kůrka et al. 2005). The species occurs in xerothermic habitats like rock steppes, xeric grasslands and edges of pine forests (Buchar & Růžicka 2002, Bryja et al. 2005). In general, it is a rare and endangered spider species belonging to most of the Red Lists in Central European countries (Gajdoš & Svatoň 2001, Řezáč et al. 2015, Blick et al. 2016).

Location: The species was very common within the study area and was found at every site. The highest abundance was at Site 1, where 191 individuals were caught. Altogether, there were 386 individuals captured in pitfall traps and one individual was swept. The species has a strong population that needs to be highlighted for nature protection and habitat management of this location.

Cheiracanthium oncognathum (Fig. 4 f)

This species is widely distributed in Europe, but only with sporadic findings all over the known distribution area (Nentwig et al. 2019). In the Czech Republic, the species rarely inhabits deciduous trees and bushes at lower and middle altitudes (Dolanský 2011). The biotopes of this species have a character of forest edges on sandy subsoil, steep slopes with sparse vegetation or rocky outcrops with aluminum benches. This species lives very covertly and sometimes makes webs in detritus or in sand not deep underground (Dolanský 2011). According to Dolanský (2011), *Cheiracanthium oncognathum* probably does not penetrate into steppe habitats, which is contrary to the present study's findings.

Location: One individual was collected in a pitfall trap in sparse vegetation at Site 1. This finding is one of the northernmost occurrences of this rare spider species from the Moravian region.

Civizelotes pygmaeus (Fig. 4 d)

Civizelotes pygmaeus has a distribution area from Europe including Italy, Slovenia, Austria, Hungary, Slovakia, Czech Republic, Poland, Ukraine, Romania, Macedonia, Russia (Eastern European) to Kazakhstan (Nentwig et al. 2019). In the Czech Republic, it is a rare species of epigeic spider occurring under stones on rock steppes, sand quarries and vineyard terraces (Buchar & Růžicka 2002, Košulič & Hula 2014). In the southern part of the Czech Republic, it is quite an abundant species, however it prefers warm locations with presence of early stages of succession (Košulič et al. 2014).

Location: There were 10 individuals collected by pitfall traps in the study area. All individuals were caught in the sparse vegetation or in the spots with a high proportion of stones at all three sites.

Zelotes aeneus (Fig. 4 h)

This species is widely distributed in Europe, except Northern Europe (Nentwig et al. 2019). In the Czech Republic, it is a rare species with occurrence under stones in steppe habitats, post-industrial biotopes in early stages of succession and grazed grasslands (Paschetta et al. 2013). In the southern part of the Czech Republic, this species is very rare with only a few records (ČAS 2019).

Location: Altogether, in the study area, 17 individuals were sampled in the pitfall traps from all sites, but most came from Site 1 and Site 2. The individuals were caught in the sparse and also dense vegetation.

Thanatus arenarius (Fig. 4 i)

This Palearctic species, widely distributed in Europe, rarely occurs in the south-eastern and north-eastern part of the Czech Republic (Szita & Samu 2000, WSC 2019). Habitats of this species are sand dunes, rock steppes, heathlands and it is quite abundant in vineyards, especially on terraces with initial stages of succession (Buchar & Růžicka 2002, Košulič & Hula 2014). Křížová (2001) recorded the species approximately 1.5 km away from the Libochovka Valley in 1999 to 2000.

Location: This xerothermic spider was sampled in pitfall traps with 704 individuals at all three sites. The highest abundance was at Site 2 where 362 individuals were caught. It was found both in sparse and dense vegetation, however, this




Fig. 4: Distribution maps of faunistically remarkable species collected in the study area (red dot). **a.** *Dysdera moravica*; **b.** *Megalephyphantes pseudocollinus*; **c.** *Trichoncus affinis*; **d.** *Porrhomma errans*; **e.** *Alopecosa striatipes*; **f.** *Cheiracanthium oncognathum*; **g.** *Civizelotes pygmaeus*; **h.** *Zelotes aeneus*; **i.** *Thanatus arenarius*; **j.** *Talavera aperta*

study suggests that open and barren surfaces are strong determinants for this species of spider.

Talavera aperta (Fig. 4j)

This species occurs from Europe to Central Asia and is considered a rare European species (Grbić et al. 2015, Nentwig et al. 2019). In Europe, the species was recorded in Serbia, Romania, Ukraine, Russia (Central European), Hungary, Slovakia, Austria, Czech Republic, Poland, Germany, Netherlands, Belgium, Switzerland, France and Italy (Nentwig et al. 2019). This xerothermic epigeic spider lives in various habitats like oak forest, vineyards and dry forest edges from different parts and altitudes of the eastern part of the Czech Republic (Buchar & Růžicka 2002, Bryja et al. 2005).

Location: At Site 1, only one individual was collected by sweeping of the sparse vegetation

Suggestions for conservation managements

The results show that Sites 1 and 2 are the most valuable parts of the study area. These parts are typical for their south or south-easterly orientation and sparse vegetation with scattered small rocks and barren surfaces. A high abundance of the threatened ground-dwelling spiders *Alopecosa striatipes* and *Thanatus arenarius* inhabiting open xeric habitats confirm the high value of these sites. The high abundance of these very rare species at Site 2 is likely due to sheep grazing that took place before 2005 and subsequent mowing with the removal of the mown materials. The recent grazing and mowing led to the reduction of plant biomass, which provided a suitable habitat (initial stages of succession) for xerothermic species such as *Alopecosa striatipes*, *Thanatus arenarius*, *Zelotes aeneus*, *Zelotes electus*, etc. However, part of Sites 1 and 2 is threatened by overgrowth and homogenization caused by the expansion of competitive plant species like *Calamagrostis epigejos* and *Arrhenatherum elatius*. This study suggests performing short-term extensive sheep grazing supplemented by mowing in order to reduce only the ungrazed areas with *Calamagrostis epigejos* (Konvička et al. 2005, Jongepierová et al. 2018). Burning is inappropriate because it supports the spread of expansive *Calamagrostis epigejos* (Házi et al. 2011, Deák et al. 2014). Furthermore, the sites are surrounded by a bush of *Prunus spinosa* and *Rosa canina*, which reduce the size of xeric, steppe patches, hence reduction of these shrubs is strongly recommended.

In contrast, rare and endangered spiders including xerothermic species were discovered in smaller numbers at Site 3. However, there was a higher occurrence of *Alopecosa trabalis* and *Agroeca cuprea*, which are more common in forest-steppes or forest edges than in xerothermic open habitats. A threat to the rocky south-east exposed slope of Site 3 is the overgrowth of expansive shrubs and the accumulation of biomass from leaves, which can strongly change microhabitat conditions (Ausden 2007). In the lower part of the site, there are areas with uniform mesophilic vegetation that are slowly expanding to the upper parts with xerothermic vegetation. In particular, this study proposes the reduction of bushes in the lower part of the slope, thus creating a larger area for xerothermic species. Certainly, regular management in the form of mosaic mowing or grazing is requested. It is necessary to leave some older shrubs such as *Crataegus monogyna* that provide suitable habitats for species like *Synema globosum*, *Trichoncus affi-*

nis, *Cheiracanthium oncognathum* and more species preferring shadier habitats.

Historically, this study acknowledges that grazing was only conducted at small scales, using a limited amount of livestock. This management approach is beneficial to the overall biodiversity of arthropods. Large scale overgrazing, on the other hand, usually has negative impacts on many organisms (Milchunas et al. 1998, Ausden 2007). Therefore, this study suggests the continuation of grazing which should be carried out in a manner that does not allow the entire area of the reserve to be grazed. Small non-grazed fences should be maintained within the grazed areas. These places serve as refuges for invertebrates for whom grazing is not suitable (Konvička et al. 2005). Additionally, prescribed burning in the early spring is suggested in small scales around selected places of the protected area (including all study sites); this can replace mowing. According to Niwa & Peck (2002), this management can reduce the biomass accumulation that is usually caused by homogeneous mowing (e.g., Noordijk et al. 2010). Burning, together with small-scale grazing and mowing, can enhance the diversity of various habitats, which in turn can improve overall biodiversity (Ausden 2007). In this context, it is also appropriate to leave some small-scale unmanaged parts in all of the sites of the protected area to create a mosaic of habitats in different stages of succession as this will provide biotopes for a wide spectrum of arthropods that may not all profit from regular disturbances such as *Agroeca cuprea* living in detritus among vegetation (Batáry et al. 2010).

In conclusion, diversification of management must be carried out in parts of the area throughout the year. Accordingly, each organism has the opportunity to find suitable habitats; if one of the interventions is poorly undertaken, other methods will compensate for its negative effects (Di Giulio et al. 2001, Ausden 2007, Bucher et al. 2016).

Acknowledgements

The authors would like to thank Petr Dolejš (National Museum in Prague) for the determination and revision of spider taxa that were difficult to identify. We likewise extend our warm appreciation to Denise Paulina V. Doble and Nicole H. Cernohorsky for the English proofreading; to Anna Foltýnová, Josef Hamřík, Lukáš Dolíhal, Marie Hamříková, Ondřej Dohnal, Pavla Vymazalová, Šárka Mašová, Viktorie Košuličová, Vojtěch Václavík, Zdeněk Sucharda and the members of ZO ČSOP Eresus and ZO ČSOP Náměšťské rybníky for their assistance during fieldwork. Additionally, we are grateful to the reviewers and editor (Theo Blick) for their comments that significantly improved the manuscript. This research was performed within the Zoological inventory of spiders in the Pláně Nature Monument and was financially supported by the South Moravian Region. The study was financially supported by the Specific University Research Fund of the Faculty of Forestry and Wood Technology, Mendel University in Brno (LDF_PSV_2017004/2017).

References

- Ausden M 2007 Habitat management for conservation: A handbook of techniques. Oxford University Press, New York. 424 pp. – doi: [10.1093/acprof:oso/9780198568728.001.0001](https://doi.org/10.1093/acprof:oso/9780198568728.001.0001)
- BAS (British Arachnological Society) 2019 Spider and harvestman recording scheme website. – Internet: <http://srs.britishtspiders.org.uk> (3. Jun. 2019)
- Batáry P, Baldi A, Sarospataki M, Kohler F, Verhulst J, Knop E, Herzog F & Kleijn D 2010 Effect of conservation management on bees and insect-pollinated grassland plant communities in three

- European countries. – Agriculture, Ecosystems & Environment 136: 35–39 – doi: [10.1016/j.agee.2009.11.004](https://doi.org/10.1016/j.agee.2009.11.004)
- Biolib 2018 Biological library. – Internet: <http://www.biolib.cz> (2. Jan. 2019)
- Blick T, Finch O-D, Harms KH, Kiechle J, Kielhorn K-H, Kreuels M, Malten A, Martin D, Muster C, Nährig D, Platen R, Rödel I, Scheidler M, Staudt A, Stumpf H & Tolke T 2016 Rote Liste und Gesamtartenliste der Spinnen (Arachnida: Araneae) Deutschlands. 3. Fassung, Stand April 2008, einzelne Änderungen und Nachträge bis August 2015. – Naturschutz und Biologische Vielfalt 70 (4): 383–510
- Bryja V, Svatoň J, Chytil J, Majkus Z, Růžicka V, Kasal P, Dolanský J, Buchar J, Chvátalová I, Řezáč M, Kubcová L, Erhart J & Fenclová I 2005 Spiders (Araneae) of the Lower Morava Biosphere Reserve and closely adjacent localities (Czech Republic). – Acta Musei Moraviae, Scientiae biologicae, Brno 90: 13–184
- Buchar J 1983 Klasifikace druhů pavoučí zvířeny Čech, jako pomůcka k bioindikaci kvality životního prostředí [Die Klassifikation der Spinnenarten Böhmens als ein Hilfsmittel für die Bioindikation der Umwelt]. Fauna Bohemiae Septentrionalis 8: 119–135 [in Czech, German summary]
- Buchar J & Růžicka V 2002 Catalogue of spiders of the Czech Republic. Peres, Praha. 351 pp.
- Buchar J & Thaler K 2004 Ein Artproblem bei Wolfspinnen: Zur Differenzierung und vikarianten Verbreitung von *Alopecosa striatipes* (C. L. Koch) und *A. mariae* (Dahl) (Araneae, Lycosidae). – Denisia 12: 271–280
- Bucher R, Andres C, Wedel MF, Entling MH & Nickel H 2016 Biodiversity in low-intensity pastures, straw meadows, and fallows of a fen area – a multitrophic comparison. – Agriculture, Ecosystems & Environment 234: 58–64 – doi: [10.1016/j.agee.2015.12.019](https://doi.org/10.1016/j.agee.2015.12.019)
- Cardoso P, Pekár S, Jocqué R & Coddington JA 2011 Global patterns of guild composition and functional diversity of spiders. – PLoS ONE 6 (6, e21710): 1–10 – doi: [10.1371/journal.pone.0021710](https://doi.org/10.1371/journal.pone.0021710)
- Colwell RK 2013 EstimateS: Statistical estimation of species richness and shared species from samples (Software and user's guide), version 9.1. – Internet: <http://viceroy.eeb.uconn.edu/estimates> (10. Jan. 2019)
- ČAS (Czech Arachnological Society) 2019 Online atlas of the Czech spiders. – Internet: <http://www.arachnology.cz> (14. Jan. 2019)
- Deák B, Valkó O, Török P, Végvári Z, Hartel T, Schmotzer A, Kapocsi I & Tóthmérész B 2014 Grassland fires in Hungary – experiences of nature conservationists on the effects of fire on biodiversity. – Applied Ecology and Environmental Research 12: 267–283
- Di Giulio M, Edwards PJ & Meister E 2001 Enhancing insect diversity in agricultural grasslands: the roles of management and landscape structure. – Journal of Applied Ecology 38: 310–319 – doi: [10.1046/j.1365-2664.2001.00605.x](https://doi.org/10.1046/j.1365-2664.2001.00605.x)
- Dolanský J 2011 Rozšíření a stanovištní nároky západních rodu *Cheiracanthium* (Araneae, Miturgidae) v Česku [Distribution and habitat preferences of spiders of the genus *Cheiracanthium* (Araneae, Miturgidae) in Czechia]. – Východočeský sborník přírodovědný, Práce a studie 18: 125–140 [in Czech, English summary]
- Duelli P 1997 Biodiversity evaluation in agricultural landscapes: An approach at two different scales. – Agriculture, Ecosystems and Environment 62: 81–91 – doi: [10.1016/S0167-8809\(96\)01143-7](https://doi.org/10.1016/S0167-8809(96)01143-7)
- ESRI (Environmental Systems Resource Institute) 2013 ArcMap 10.2 ESRI, Redlands, California. – Internet: <https://www.esri.com/en-us/home> (14. Aug. 2018)
- Gajdoš P & Svatoň J 2001 Red (Ecosozological) List of spiders (Araneae) of Slovakia. In: Baláz D, Marhold K & Urban P (eds) Red List of plants and animals of Slovakia. Nature Conservation, ŠOP SR Banská Bystrica. pp. 80–86
- Grbić G, Hänggi A & Savić D 2015 New faunistic records of spiders (Arachnida, Araneae) from the Fruška Gora Mountain, northern Serbia. – Acta Zoologica Bulgarica 67: 479–486
- Hamřík T 2019 Struktura a distribuce společenstev pavouků (Araneae) v závislosti na ochrannářském managementu přírodní památky Pláně [Structure and distribution of spider communities (Araneae) depending on conservation management of Pláně Nature Monument.]. Diploma thesis, Mendel University, Brno. 82 pp. [in Czech, English abstract]
- Házi J, Bartha S, Szentes S, Wichmann B & Penszka K 2011 Seminal grassland management by mowing of *Calamagrostis epigejos* in Hungary. – Plant Biosystems – An International Journal Dealing with all Aspects of Plant Biology 145: 699–707 – doi: [10.1080/11263504.2011.601339](https://doi.org/10.1080/11263504.2011.601339)
- Heimer S & Nentwig W 1991 Spinnen Mitteleuropas: ein Bestimmungsbuch. Paul Parey, Berlin, Hamburg. 544 pp.
- Hula V, Košulič O & Šťastná P 2012 Spiders (Araneae) of selected sinkholes of Moravský kras Protected Landscape Area (Czech Republic). – Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis 60: 57–70 – doi: [10.11118/actaun201260010057](https://doi.org/10.11118/actaun201260010057)
- Jongepierová I, Prach K & Pešout P 2018 Ekologická obnova v České republice II [Ecological restoration in the Czech Republic II]. Agentura ochrany přírody a krajiny ČR, Praha. 212 pp. [in Czech]
- Konvička M, Beneš J & Čížek L 2005 Ohrožený hmyz nelesních stanovišť: ochrana a management [Threatened insect of non-forest habitats: conservation and management]. Sagittaria, Olomouc. 127 pp. [in Czech]
- Košulič O & Hula V 2014 A faunistic study on spiders (Araneae) from vineyard terraces in the municipalities of Morkůvky and Mutěnice (South Moravia, Czech Republic). – Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis 62: 137–154 – doi: [10.11118/actaun201462010137](https://doi.org/10.11118/actaun201462010137)
- Košulič O, Michalko R & Hula V 2014 Recent artificial vineyard terraces as a refuge for rare and endangered spiders in a modern agricultural landscape. – Ecological Engineering 68: 137–154 – doi: [10.1016/j.ecoleng.2014.03.030](https://doi.org/10.1016/j.ecoleng.2014.03.030)
- Košulič O, Michalko R & Hula V 2016 Impact of canopy openness on spider communities: Implications for conservation management of formerly coppiced oak forests. – PLoS ONE 11 (2, e0148585): 1–18 – doi: [10.1371/journal.pone.0148585](https://doi.org/10.1371/journal.pone.0148585)
- Košulič O 2017 Faunistic study on spiders (Araneae) in the Špraněk National Nature Reserve with suggestion to conservation management of the locality. – Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis 65: 1521–1535 – doi: [10.11118/actaun201765051521](https://doi.org/10.11118/actaun201765051521)
- Křížová V 2001 Pavoučí zvířena okolí obce Říkonín [Spider fauna in the surrounding areas of Říkonín village]. Bachelor thesis, University, Ostrava. 27 pp. [in Czech, English abstract]
- Kůrka A, Řezáč M, Macek R & Dolanský J 2015 Pavouci České republiky [Spiders of the Czech Republic]. Academia, Praha. 623 pp. [in Czech]
- Marc P, Canard A & Ysnel F 1999 Spiders (Araneae) useful for pest limitation and bioindication. – Agriculture Ecosystems and Environment 74: 229–273 – doi: [10.1016/S0167-8809\(99\)00038-9](https://doi.org/10.1016/S0167-8809(99)00038-9)
- Matuška P 2016 Plán péče o přírodní památku Pláně na období 2016–2025 [Conservation plan for Pláně Nature Monument in period 2016–2025]. Deposited in: Krajský úřad Jihomoravského kraje, Brno. 17 pp. [in Czech]
- Michalko R, Košulič O, Hula V & Surovcová K 2016 Niche differentiation of two sibling wolf spider species, *Pardosa lugubris* and *Pardosa alacris*, along a canopy openness gradient. – Journal of Arachnology 44: 46–51 – doi: [10.1636/M15-46.1](https://doi.org/10.1636/M15-46.1)
- Miller F 1974 Pavoučí fauna řepných polí v okolí Chválkovic a Nákla na Hané [Spider fauna of sugar beet fields in the surroundings of Chválkovice and Náklo in Haná]. – Acta Universitatis Palackianae Olomouensis – Biologica 15: 175–181 [in Czech, English summary]
- Milchunas DG, Lauenroth WK & Burke IC 1998 Livestock grazing: animal and plant biodiversity of shortgrass steppe and the relationship to ecosystem function. – Oikos 83: 65–74 – doi: [10.2307/3546547](https://doi.org/10.2307/3546547)
- Nenadál S 1994 Lada u Kuřimské Nové Vsi. Inventarizační výzkum. [Pasture near Kuřimská Nová Ves: Inventory Survey]. Deposited in: Krajský úřad Jihomoravského kraje, Brno. 19 pp. [in Czech]

- Nentwig W, Blick T, Gloor D, Hänggi A & Kropf C 2019 Spiders of Europe. Version 6.2019 – Internet: <http://araneae.nmbe.ch> (5. Jun. 2019) – doi: [10.24436/1](https://doi.org/10.24436/1)
- Niwa CG & Peck RW 2002 Influence of prescribed fire on carabid beetle (Carabidae) and spider (Araneae) assemblages in forest litter in southwestern Oregon. – *Environmental Entomology* 31: 785–796 – doi: [10.1603/0046-225X-31.5.785](https://doi.org/10.1603/0046-225X-31.5.785)
- Noordijk J, Schaffers AP, Heijerman T, Boer P, Gleichman M & Sýkora KV 2010 Effects of vegetation management by mowing on ground-dwelling arthropods. – *Ecological Engineering* 36: 740–750 – doi: [10.1016/j.ecoleng.2010.01.003](https://doi.org/10.1016/j.ecoleng.2010.01.003)
- Paschetta M, La Morgia V, Masante D, Negro M, Rolando A & Isaia M 2013 Grazing history influences biodiversity: a case study on ground-dwelling arachnids (Arachnida: Araneae, Opiliones) in the Natural Park of Alpi Marittime (NW Italy). – *Journal of Insect Conservation* 17: 339–356 – doi: [10.1007/s10841-012-9515-y](https://doi.org/10.1007/s10841-012-9515-y)
- Pearce JL, Venier LA, Eccles G, Pedlar J & McKenney D 2004 Influence of habitat and microhabitat on epigeal spider (Araneae) assemblages in four stand types. – *Biodiversity and Conservation* 13: 1305–1334 – doi: [10.1023/B:BIOC.0000019403.26948.55](https://doi.org/10.1023/B:BIOC.0000019403.26948.55)
- Pech P, Dolanský J, Hrdlička R & Lepš J 2015 Differential response of communities of plants, snails, ants and spiders to long-term mowing in a small-scale experiment. – *Community Ecology* 16: 115–124 – doi: [10.1556/168.2015.16.1.13](https://doi.org/10.1556/168.2015.16.1.13)
- Ratschker UM & Roth M 2000 Studies on ground dwelling spiders (Araneae) of agrarian habitat types in Northeast Germany: ecological and nature conservation aspects. – *Ekologia* 19, Supplement 3: 213–225
- Řezáč M 2012 Rozšíření a stanovištní nároky pavouků šestičok rodu *Dysdera* (Araneae: Dysderidae) v Česku [Distribution of the six-eyed spiders of the genus *Dysdera* (Araneae: Dysderidae) in Czechia]. – *Příroda* 30: 117–162 [in Czech, English abstract]
- Řezáč M, Kůrka A, Růžicka V & Heneberg P 2015 Red List of Czech spiders: 3rd edition, adjusted according to evidence-based national conservation priorities. – *Biologia* 70: 645–666 – doi: [10.1515/biolog-2015-0079](https://doi.org/10.1515/biolog-2015-0079)
- Růžicka V 2018 A review of the spider genus *Porrhomma* (Araneae, Linyphiidae). – *Zootaxa* 4481: 1–75 – doi: [10.11646/zootaxa.4481.1.1](https://doi.org/10.11646/zootaxa.4481.1.1)
- Samu F, Sunderland KD & Szinetár C 1999 Scale-dependent dispersal and distribution patterns of spiders in agricultural systems: a review. – *Journal of Arachnology* 27: 325–332
- Samu F, Szinetár C, Szita É, Fetykó K & Neidert D 2011 Regional variations in agrobiont composition and agrobiont life history of spiders (Araneae) within Hungary. – *Arachnologische Mitteilungen* 40: 105–109 – doi: [10.5431/aramit4012](https://doi.org/10.5431/aramit4012)
- Skalický V 1988 Regionálně fytogeografické členění. [Regional phytogeographical division]. In: Hejný S & Slavík B (eds): *Květena České socialistické republiky 1*. [Flora of the Czech Socialist Republic. Vol. 1]. Praha: Academia. 103–121 [in Czech]
- Slavík B 1988 Regionálně fytogeografické členění [Regional phytogeographical division. A map]. In: *Květena České socialistické republiky 1*. [Flora of the Czech Socialist Republic. Vol. 1]. Praha: Academia, mapová příloha, Praha. 557 pp. [in Czech]
- Stoate C, Báldi A, Beja P, Boatman ND, Herzon I, Doorn A van Snoo GR de, Rakosy L & Ramwell C 2009 Ecological impacts of early 21st century agricultural change in Europe – a review. – *Journal of Environmental Management* 91: 22–46 – doi: [10.1016/j.jenvman.2009.07.005](https://doi.org/10.1016/j.jenvman.2009.07.005)
- Svatoň J & Jelínek A 1998 Příspěvek k poznání pavoučí zvířeny (Araneae) v údolí řeky Brtnice na Českomoravské vysocině [Contribution to the knowledge of spider fauna (Araneae) in the Brtnice river valley in the Bohemian-Moravian Highlands]. *Vlastivědný sborník Vysočiny. Oddíl věd přírodních* 13: 83–109 [in Czech, English summary]
- Szita É & Samu F 2000 Taxonomical review of *Thanatus* species (Phidromidae, Araneae) of Hungary. – *Acta Zoologica Academiae Scientiarum Hungaricae* 46: 155–179
- Šálek M, Hula V, Kipson M, Daňková R, Niedobová J & Gamero A 2018 Bringing diversity back to agriculture: Smaller fields and non-crop elements enhance biodiversity in intensively managed arable farmlands. – *Ecological Indicators* 90: 65–73 – doi: [10.1016/j.ecolind.2018.03.001](https://doi.org/10.1016/j.ecolind.2018.03.001)
- Tichý S 2016 Inventarizační floristický průzkum přírodní památky [Botanical Inventory of Pláně Nature Monument]. Deposited in: Krajský úřad Jihomoravského kraje, Brno. 22 pp. [in Czech]
- Tscharntke T, Klein AM, Kruess A, Steffan-Dewenter I & Thies C 2005 Landscape perspectives on agricultural intensification and biodiversity – ecosystem service management. – *Ecology Letters* 8: 857–874 – doi: [10.1111/j.1461-0248.2005.00782.x](https://doi.org/10.1111/j.1461-0248.2005.00782.x)
- Valkó O, Török P, Deák B & Tóthmérész B 2014 Prospects and limitations of prescribed burning as a management tool in European grasslands. – *Basic and Applied Ecology* 15: 26–33 – doi: [10.1016/j.baae.2013.11.002](https://doi.org/10.1016/j.baae.2013.11.002)
- Wheeler WC, Coddington JA, Crowley LM, Dimitrov D, Goloboff PA, Griswold CE, Hormiga G, Prendini L, Ramírez MJ, Sierwald P, Almeida-Silva LM, Álvarez-Padilla F, Arnedo MA, Benavides LR, Benjamin SP, Bond JE, Griswold CJ, Hasan E, Hedin M, Izquierdo MA, Labarque FM, Ledford J, Lopardo L, Maddison WP, Miller JA, Piacentini LN, Platnick NI, Polotow D, Silva-Dávila D, Scharff N, Szűts T, Ubick D, Vink C, Wood HM & Zhang JX 2017 The spider tree of life: phylogeny of Araneae based on target-gene analyses from an extensive taxon sampling. – *Cladistics* 33: 576–616 – doi: [10.1111/cla.12182](https://doi.org/10.1111/cla.12182)
- WSC 2019 World spider catalog. Version 20.0. Natural History Museum Bern. – Internet: <http://wsc.nmbe.ch> (16. Jan. 2019) – doi: [10.24436/2](https://doi.org/10.24436/2)