

New and noteworthy lichen records from the Ionian island of Kerkira (Corfu), Greece.

Authors: Christensen, Steen N., Pisút, Ivan, and Sipman, Harrie J. M.

Source: Willdenowia, 27(1/2) : 265-272

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.27.2726>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

STEEN N. CHRISTENSEN, IVAN PIŠÚT & HARRIE J. M. SIPMAN

New and noteworthy lichen records from the Ionian island of Kerkira (Corfu), Greece.

Abstract

Christensen, S. N., Pišút, I. & Sipman, H. J. M.: New and noteworthy lichen records from the Ionian island of Kerkira (Corfu), Greece. – Willdenowia 27: 265–272. 1997. – ISSN 0511–9618.

Thirty-six lichen taxa are reported from Kerkira. Of these, nine taxa are new to Greece, namely *Anthraccarpon virescens*, *Bactrospora patellarioides* var. *convexa*, *Gyalecta derivata*, *G. truncigena*, *Lecania koerberiana*, *Leptogium subtile*, *Opegrapha vulgata*, *Parmelia subrudecta*, and *Phaeophyscia hirsuta*, and twelve taxa are new to Kerkira.

Introduction

In a previous contribution to the lichens of Kerkira (Christensen 1995b) also the natural features (topography, climate and vegetation) as well as the existing literature on the lichen flora of the island has been treated. This contribution presents further new and noteworthy records resulting from collections independently made by the three authors. In the light of the present state of knowledge of the Greek lichen flora, also taxa with a single or only few earlier records from Kerkira have been included. The paper should be seen as a small contribution to the knowledge of the Greek lichen flora prior to a future lichen checklist of Greece (Nimis 1996).

Material and methods

The lichen material collected by S. N. Christensen in April 1983 is deposited in the Botanical Museum of Copenhagen (C) and in his private herbarium, the material collected by I. Pišút in September 1995 is deposited in the Institute of Botany of the Slovak Academy of Sciences at Bratislava (SAV) and in his private herbarium, and the specimens collected by H.J.M. Sipman in August 1970 are deposited in the Berlin herbarium (B).

In the list of the taxa, the collectors are indicated by their initials, and the nomenclature follows, with few exceptions, Nimis (1993). Taxa new to Greece are marked with a double asterisk (**) and taxa new to Kerkira with a single asterisk (*). Whenever possible, geographical names are quoted in accordance with The Times Atlas of the World 1990.

Collecting localities

Mt Pantokrator: summit, boulder and stone field, limestone, c. 900 m.

Spartilas: c. 13 km N of Kerkira town, SW slope of Mt Pantokrator, just N of the village Spartilas,

maquis on limestone with *Quercus coccifera*, *Pistacia terebinthus*, *Phlomis fruticosa* and *Cistus*, c. 400 m.

Angelokastro: c. 2 km WNW of Palaeokastritsa, ruined castle on limestone rock, phrygana.

Ropa: c. 10 km W of Kerkira town, plain of Ropa, c. 1 km S of the village of Ropa, on *Prunus domestica* and *Ulmus minor* trees along road in arable land.

Gouvia: old olive grove (cf. Christensen 1995b).

Anemomylos: 2 km S of Kerkira town along the road to Kanoni, near the summer residence of the Greek king, oak forest with mossy trunks, 20 m.

Achillion: palace built by the Empress Elizabeth of Austria, c. 9 km S of Kerkira town, the village of Gastouri, rural landscape in the environs, 100 m.

Benitses: village on the east coast, surrounded by olive covered hills, c. 10 km S of Kerkira town.

Aghii Deka: Mt Aghii Deka, 2 km S of the village Aghii Deka, at the road to the village Stavros, *Cupressus sempervirens* road side trees.

Ano Garuna: c. 9 km SW of Kerkira town, Mt Aghii Deka, c. 100 m N of the village Ano Garuna, maquis, c. 475 m.

Gardiki castle: c. 22 km SSW of Kerkira town, c. 2 km S of Aghios Mattheos.

List of the taxa

***Anthracocarpon virescens* (Zahlbr.) Breuss $\equiv$ *Catapyrenium virescens* (Zahlbr.) Breuss, cf. Breuss (1996).

Benitses: On N-facing rock on hill slope S of the springs, 200 m, *HJMS* 2749.

Hitherto known from Spain and Croatia only (Breuss 1990).

***Bactrospora patellarioides* var. *convexa* (de Lesd.) Egea & Torrente

Benitses: On bark and lignum of *Olea europaea*, *IP*. Ascomata small, up to 0.6 mm in diam., with inconspicuous margins, subhymenium brown.

B. patellarioides var. *patellarioides* is known in Greece from one site ("Corfu: Gastouri, 10 km S of Corfu town, forest on E slope of Achillion, on the stem of *Quercus coccifera*, 100 m, 1970, leg. H. Sipman", cf. Egea & Torrente 1993). The var. *convexa* is up to now known only from numerous sites in Italy (Egea & Torrente 1993).

Caloplaca erythrocarpa (Pers.) Zwackh

Spartilas: On limestone wall in maquis, *SNC* 1438.

Achillion: On calcareous rocks on the E slope of the castle, *IP*.

Reported from Attica (Christensen & al. 1991), Thira (Sipman & Raus 1995), Rodhos (corticolous on *Ficus*!, Cengia-Sambo 1927), and only once from Kerkira (Arnold 1887, as *Blastenia lallavei* Clem.).

**Caloplaca velana* var. *schaereri* (Flörke) Clauzade & Roux

Spartilas: On limestone wall in maquis, *SNC* (cum 1436).

Gouvia: On limestone boulder in olive grove, *SNC* (cum 1386, 1390).

Known from the Ionian island of Levkas (Servít 1935), from Attica (Christensen & al. 1991), the Peloponnisos (Rondon 1970), Evvoia (Krause & Klement 1962), and the Aegean region (Szatala in Rechinger 1943).

Catillaria nigroclavata (Nyl.) Schaer.

Anemomylos: On dead *Pinus* branchlet, *HJMS* 2765D.

There is only one earlier record from Kerkira (Steiner 1919, "Auf Stämmen von *Opuntia*"),

and the species is also known from the Peloponnisos (Servít 1935) and the Aegean island of Delos (Steiner 1919).

***Cladonia pseudopityrea* Vain.**

Benitises: On the wood of *Olea europaea*, *IP*.

Once reported from Kerkira (Ahti & Puntillo 1995).

****Collema auriforme* (With.) Coppins & Laundon**

Anemomylos: On N-facing side of shady limestone wall along road, *HJMS 2781B*.

Benitises: On N-facing side of limestone wall along road towards W through the valley, 50 m, *HJMS 2702*.

Known from the Greek mainland (Christensen 1995a, Degelius 1956, 1974, Szatala in Rechinger 1951, Szatala 1959), the Peloponnisos (Christensen 1994b, Degelius 1974) and the Aegean island of Thira (Sipman & Raus 1995).

****Collema fuscovirens* (With.) Laundon**

Benitises: On flat, upper face of slightly shaded limestone boulder in olive grove, 50 m, *HJMS 2762*.

Known from the Greek mainland (Degelius 1956, 1974).

***Collema polycarpon* subsp. *corcyrense* (Arnold) Pišút**

Achillion: On calcareous rocks in the village of Achillion, *IP*.

From Kerkira (Pass San Pantaleone) reported as *Lethagrium orbiculare* f. *corcyrense* Arn. by Arnold (1887). It is interesting that Arnold already considered a higher taxonomic rank for this taxon ("forsan nova species"). The distribution of this subspecies was mapped by Pišút (1968).

****Collema subflaccidum* Degel.**

Benitises: On bark of *Olea europaea*, *IP*; on trunk of *Robinia pseudacacia*, 50 m, *HJMS 2735*.

This species, resembling *C. flaccidum*, but differing in its granuliform isidia, was not known from Kerkira. Previously Degelius (1954, 1956, 1974) recorded it from the Ionian island of Kefalonia and mainland Greece, and Christensen (1995a) from Mt Olimbos.

***Collema tenax* (Sw.) Ach.**

Achillion: On calcareous soil on the E slope of the palace Achillion, *IP*.

Recorded from Kerkira in Arnold (1887) and Steiner (in Rechinger 1915) as *C. pulposum* (Bernh.) Ach. The species has previously been collected at Achillion (Degelius 1954).

***Diploschistes muscorum* (Scop.) R. Sant.**

Benitises: On bare bark of *Olea europaea*, *IP*. Asci 4-spored, thallus K+ yellow, C+ pink red.

Gardiki castle: On *Cladonia pocillum* (*SNC 1590*) on top of the ring wall.

According to Lumbsch (1989), *D. muscorum* is the only species of the genus occasionally occurring also on bark. Christensen (1995b) reported it from Gouvia, growing on *Cladonia chlorophaea*.

****Evernia prunastri* f. *herinii* (P.A. Duvign.) Hawksw. (≡ *Evernia herinii* P.A. Duvign.)**

Benitises: On the bark of *Olea europaea*, together with *Ramalina fastigiata* (Pers.) Ach. and *R. farinacea* (L.) Ach., *IP*.

This forma, characterized by the grey colour of the upper side of the lobes and the absence of usnic acid, has only been reported from Mt Parnes on the Greek mainland (Degelius 1956).

****Evernia prunastri* (L.) Ach. f. *prunastri***

Achillion: On trunk of *Prunus persica* in orchard, *HJMS 2641*.

E. prunastri is a widespread and common epiphyte recorded from several localities on the Greek mainland (e.g. Christensen 1989, 1994a, Halácsy 1894, Politis 1953, Steiner 1898, 1919,

Szatala 1941, 1959, Wilmanns & Phitos 1960), from Evvoia (Hepp in Unger 1862), Kriti (e.g. Szatala in Rechinger 1943, 1951, Zahlbruckner 1906), the Aegean islands (Steiner 1919, Szatala in Rechinger 1943) and the Ionian island of Kefallinia (Zahlbruckner 1907).

*****Gyalecta derivata* (Nyl.) H. Olivier**

Benitses: On tree trunk near farm, 50 m, *HJMS* 2713.

*****Gyalecta truncigena* (Ach.) Hepp**

Achillion: On trunk of *Quercus coccifera* in forest, *HJMS* 2639B.

*****Lecania koerberiana* J. Lahm**

Anemomylos: On dead branchlet of *Pinus*, *HJMS* 2764B; on branchlet of dead *Ficus*?, *HJMS* 2766C.

****Lecanora meridionalis* H. Magn.**

Ropa: On twigs of *Prunus domestica*, *SNC* (cum 1422a), and of *Ulmus minor*, *SNC* 1433a.

Reported from the Pindos range (Christensen 1989), Evvoia (Krause & Klement 1962) and the Aegean island of Thira (Sipman & Raus 1995).

***Leptogium gelatinosum* (With.) Laundon**

Pantokrator: Summit, among mosses in boulder and stone field, limestone, *SNC* 1507a.

Spartilas: Among mosses on limestone outcrop in maquis, *SNC* 1435a.

Anemomylos: On N-facing side of shady limestone wall along road, *HJMS* 2781A.

As *L. sinuatum* (Huds.) Massal. recorded from Kerkira by Christensen (1995b).

*****Leptogium subtile* (Schrad.) Torss.**

Achillion: On the bark of *Olea europaea* on the E slope below the castle, *IP*; det. A. Guttová.

Benitses: On trunk of *Robinia pseudacacia*, 50 m, *HJMS* 2728B; on trunk of olive in olive grove, *HJMS* 2752B.

This widespread lichen has its closest published locality in N Italy (Jørgensen 1994, Nimis 1993). It does, however, also occur on the Greek mainland (Christensen in prep.).

*****Opegrapha vulgata* Ach.**

Benitses: On trunk of very shaded *Cupressus* tree near waterfall, 100 m, *HJMS* 2737.

***Pannaria olivacea* P.M. Jørg.**

Benitses: On the bark of *Olea europaea*, *IP*. Numerous fertile specimens.

In Kerkira probably not rare. Reported from Achillion by Arnold (1887) as *P. craspedia*. Jørgensen (1978) quoted it from Kariotikon, Kiriaki, the foot of Mt Pantokrator near Spartilas and S. Nivolo. Christensen (1995b) mentioned a sterile specimen from Nimphes.

*****Parmelia subrudecta* Nyl.**

Achillion: On trunk of *Prunus persica* in orchard, *HJMS* 2648.

Benitses: On trunk of *Robinia pseudacacia*, 50 m, *HJMS* 2725.

****Parmelia sulcata* Taylor**

Benitses: On trunk of olive in olive grove, 200 m, *HJMS* 2751.

This widespread species is common on the Greek mainland (Christensen 1989, 1994a, Diamantopoulos & al. 1992, Diapoli 1936, Geissler 1977, Harmand & Maire 1909, Hayek 1928, Papastefanou & al. 1992, Pirintso & al. 1993[a, b], 1995, Politis 1953, Räsänen 1944, Szatala 1940, 1941, 1959, Szatala in Rechinger 1951, Walter 1976, Wilmanns 1959), and is known from Evvoia (Krause & Klement 1962), Kriti (Kleinig 1966, Szatala in Rechinger 1951) and the Aegean region (Servít 1933).

****Parmotrema reticulatum*** (Taylor) M. Choisy

Achillion: On the bark of *Olea europaea* on E slope below the castle, *IP* [medulla and soralia K+ yellow turning red, C-, P+ pale orange-red, containing salazinic acid]; on trunk of *Prunus persica* in orchard, *HJMS* 2649.

Benitses: On trunk of *Robinia pseudacacia*, 50 m, *HJMS* 2716.

A pantemperate species which has a southwestern distribution in Europe. In Italy it occurs along the west coast and in the central northern part of the peninsula (Nimis 1993). Reported as *Rimelia reticulata* (Tayl.) Hale & A. Fletcher from the Aegean island of Thira (Sipman & Raus 1995).

Pertusaria heterochroa (Müll. Arg.) Erichsen

Anemomylos: On branchlet of dead *Ficus*?, *HJMS* 2766A.

Only two earlier records are known from Kerkira (Arnold 1887 as *Pertusaria pustulata* f. *superpallens* Nyl. and Christensen 1995b).

*****Phaeophyscia hirsuta*** (Mereschk.) Essl.

Benitses: On trunk of *Robinia pseudacacia*, 50 m, *HJMS* 2723.

Physcia biziana (Massal.) Zahlbr.

Spartilas: On limestone outcrop in maquis, *SNC* 1437

Ph. biziana has been reported from Ipiros (Christensen 1994b), the Peloponnisos (Rondon 1970), Kriti (Rondon 1969, Szatala in Rechinger 1951), the Aegean island of Thira (Sipman & Raus 1995), and from Kerkira (Christensen 1995b). The present specimen from Kerkira and that from Thira represent var. *leptophylla* Vězda, but this variety has been considered a modification without taxonomic importance (Sipman & Raus 1995).

Physconia distorta (With.) Laundon

Ano Garuna: On *Olea* in olive grove, *SNC* 1415.

The specimen belongs to f. *turgida* (Schaer.) J. Nowak. This sterile form with numerous small lobules on the lobe margins is photographically illustrated by Nowak (1993). It is probably the same as var. *subvenusta* (Clauzade & Roux 1985). A specimen was erroneously published as *P. venusta* (Ach.) Poelt by Christensen (1995b). *P. distorta* has been reported from Kerkira by Körber (1868 as "*Parmelia pulverulenta* (Schreb.)"). The variety *subvenusta* has been reported from Kriti (Zahlbruckner 1906). It is questionable whether this lobulate form is anything but an ecological modification (Moberg 1977).

****Physconia grisea*** (Lam.) Poelt subsp. ***grisea***

Aghii Deka: On the trunk of *Cupressus sempervirens* wayside tree along dirt road, *SNC* 1598.

Ph. grisea has been reported from Thessaloniki (Diamantopoulos & al. 1992), Evvoia (Krause & Klement 1962), the Aegean island of Kithira (Szatala in Rechinger 1943) and Kriti (Kleinig 1966).

****Placolecis opaca*** (Fr.) Hafellner

Achillion: On calcareous rocks of a road-wall on the E slope below Achillion, *IP*.

Reported as *Lecidea opaca* Duf. from the Ionian island of Levkas by Harmand & Maire (1909), from Kriti (Szatala in Rechinger 1951), and Mt Olimbos (Szatala 1959).

****Protoblastenia rupestris*** (Scop.) J. Steiner

Spartilas: On limestone outcrop in maquis, *SNC* 1439.

Known from the Greek mainland (Hayek 1928, Rondon 1970, Steiner 1898, Szatala 1959) and Evvoia (Krause & Klement 1962).

****Roccella phycopsis*** (Ach.) Ach.

Angelokastro: On the limestone walls of the castle, *SNC* 1601, 1602, 1603.

Known from Rodhos (Cengia-Sambo 1924), Kriti (Kleinig 1966, Rondon 1969, Szatala in

Rechinger 1951, Zahlbruckner 1906) and several other places in the Aegean region (Cengia-Sambo 1927, Servít 1933, Steiner 1919, Szatala in Rechinger 1943, Zahlbruckner 1907) as well as from the Peloponnisos (Rondon 1970, Steiner 1919).

***Staurolemma omphalarioides* (Anzi) P.M. Jørg. & Henssen**

Achillion: Together with *Ramalina canariensis* J. Steiner on the bark of *Olea europaea* on the E slope below the castle, *IP*; on trunk of *Prunus persica* in orchard, *HJMS* 2645.

Benitses: On trunk of *Robinia pseudacacia*, 50 m, *HJMS* 2714.

Reported by Steiner (in Rechinger 1915) from Kerkira. The world distribution was mapped by Jørgensen & Henssen (1993).

***Xanthoria calcicola* Oxner**

Aghii Deká: On base of road-facing trunk of a dirt covered *Cupressus sempervirens* wayside tree, *SNC* 1597; a yellowish orange shade form.

This normally saxicolous species of calcareous substrata has previously been recorded from the Greek mainland (Steiner 1898, 1919), Evvoia (Krause & Klement 1962), the Peloponnisos (Rondon 1970), the Aegean region (Servít 1933, Sipman & Raus 1995, Steiner 1919, Szatala in Rechinger 1943, Zahlbruckner 1907), Kriti (Kleinig 1966, Rondon 1969), and the Ionian island of Kefallinia (Hepp in Unger 1862), all as *X. aureola*. Once it was reported from Kerkira (as “*Parmelia parietina* L. *β. aureola* Ach. a. *corticola* Hepp”) by Hepp (in Unger 1862).

Acknowledgments

We thank O. Breuss and A. Guttová for the determination/confirmation of certain specimens.

References

- Ahti, T. & Puntillo, D. 1995: The status of *Cladonia pseudopityrea*, an overlooked Mediterranean lichen. – Pp. 155–160 in: Daniels, F. J. A., Schulz, M. & Peine, J. (ed.), *Flechten Follmann. Contributions to lichenology in honour of Gerhard Follmann*. – Cologne.
- Arnold, F. 1887: Lichenologische Fragmente XXVIII. – *Flora* **70**: 145–164.
- Breuss, O. 1990: Die Flechtengattung *Catapyrenium* (*Verrucariaceae*) in Europa. – *Stapfia* **23**.
- 1996: Ein verfeinertes Gliederungskonzept für *Catapyrenium* (lichenisierte Ascomyceten, *Verrucariaceae*) mit einem Schlüssel für die bisher bekannten Arten. – *Ann. Naturhist. Mus. Wien* **98B, Suppl.**: 35–50.
- Cengia-Sambo, M. 1924: Licheni di Rodi raccolti dal prof. Adriano Fiori nell’agosto 1923. – *Bull. Soc. Bot. Ital.* **1924**: 123–125.
- 1927: Licheni di Rodi. – *Nuovo Giorn. Bot. Ital.*, ser. 2, **34**: 829–840.
- Christensen, S. N. 1989: The lichen flora on *Ulmus* in a north Greek mountain village with an introduction to the literature on Greek lichens. – *Graphis Scripta* **2**: 107–113.
- 1994a: Lichens associated with *Pinus nigra* on Mt. Trapezitsa, Epirus, NW Greece. – *Acta Bot. Fenn.* **150**: 11–20.
- 1994b: Lichens on plane trees in two Greek localities with contrasting environments. – *Bot. Hron.* **11**: 31–39.
- 1995a: *Fulgensia klementii* and other lichens from Mt. Olimbos, Makedhonia, Greece. – *Willdenowia* **25**: 283–288.
- 1995b: Lichens from olive groves on the Ionian island Corfu (Kerkyra), Greece. – *Flora Medit.* **5**: 191–209.
- , Alstrup, V. & Voreadou, C. 1991: Lichens from Attica, Greece. – *Graphis Scripta* **3**: 97–103.
- Clauzade, G. & Roux, C. 1985: *Likenoj de Okcidenta Europo. Ilustrita determinlibro*. – *Bull. Soc. Bot. Centre-Ouest*, ser. 2, Num. Spéc. **7**.
- Degelius, G. 1954: The lichen genus *Collema* in Europe. Morphology, taxonomy, ecology. –

- Symb. Bot. Upsal. **13**(2).
- 1956: Studies in the lichen family *Collemataceae* II. On the *Collema* flora of the mainland of Greece. – Svensk Bot. Tidskr. **50**: 496–512.
- 1974: The lichen genus *Collema* with special reference to the extra-european species. – Symb. Bot. Upsal. **20**(2).
- Diamantopoulos, J., Pirintsos, S., Laundon, J. R. & Vokou, D. 1992: The epiphytic lichens around Thessaloniki (Greece) as indicators of sulphur dioxide pollution. – Lichenologist **24**: 63–71.
- Diapouli, Ch. A. 1936: Peri tinon leichinon tou Olimpou. – “To Vouno”, Bull. mens. Club Alpin Hellén. **3**(34): 225–232.
- Egea, J. M. & Torrente, P. 1993: The lichen genus *Bactrospora*. – Lichenologist **25**: 211–255.
- Geissler, P. 1977: Zur Moos- und Flechtenflora Nordgriechenlands. – Bauhinia **6**: 189–213.
- Halácsy, E. von 1894: Botanische Ergebnisse einer im Auftrage der hohen kaiserl. Akademie der Wissenschaften unternommenen Forschungsreise in Griechenland IV. Beitrag zur Flora von Achaia und Arcadien. – Denkschr. Kaiserl. Akad. Wiss. Wien, Math.-Naturwiss. Kl. **61**: 487–535.
- Harmand, J. & Maire, R. 1909: Contribution à l'étude des lichens de la Grèce. – Bull. Soc. Sci. Nancy, ser. 3, **10**: 143–176.
- Hayek, A. 1928: Ein Beitrag zur Kenntnis der Vegetation und der Flora des thessalischen Olymp. – Beih. Bot. Centralbl., Abt. 2, **45**: 220–328.
- Jørgensen, P. M. 1978: The lichen family *Pannariaceae* in Europe. – Opera Bot. **45**.
- 1994: Further notes on European taxa of the lichen genus *Leptogium*, with emphasis on the small species. – Lichenologist **26**: 1–29.
- & Henssen, A. 1993: *Physma omphalarioides* – its taxonomic position and phytogeography. – Graphis Scripta **5**: 12–17.
- Kleinig, H. 1966: Beitrag zur Kenntnis der Flechtenflora von Kreta. – Nova Hedwigia **11**: 513–526.
- Körber, G. W. 1868: Specimen florae cryptogamae septem insularum editum juxta plantas Mazzarianas herbarii Heufleriani et speciatim quoad filices herbarii Tommasiniani V. Lichenes. – Verh. K. K. Zool.-Bot. Ges. Wien **18**: 425–426.
- Krause, W. & Klement, O. 1962: Zur Kenntnis der Flora und Vegetation auf Serpentinstandorten des Balkans 5. Flechten und Flechtengesellschaften auf Nord-Euböa (Griechenland). – Nova Hedwigia **4**: 182–262.
- Lumbsch, H. Th. 1989: Die holarktischen Vertreter der Flechtengattung *Diploschistes* (*Thelotrema* *mataceae*). – J. Hattori Bot. Lab. **66**: 133–196.
- Moberg, R. 1977: The lichen genus *Physcia* and allied genera in Fennoscandia. – Symb. Bot. Upsal. **22**(1).
- Nimis, P. L. 1993: The lichens of Italy. An annotated catalogue. – Mongr. Mus. Region. Sci. Nat. Torino **12**.
- 1996: Towards a checklist of Mediterranean lichens. – Bocconeia **6**: 5–17.
- Nowak, J. 1993: *Physciaceae* sensu stricto. – In: Kochman, J., Nowak, J., Siemińska, J. & Skirgiełło, A. (ed.), Flora Polska, Porosty (Lichenes) **6**(3).
- Papastefanou, C., Manolopoulou, M. & Sawidis, T. 1992: Residence time and uptake rates of ¹³⁷Cs in lichens and mosses at temperate latitude (40°N). – Environm. Int. **18**: 397–401.
- Pirintsos, S. A., Diamantopoulos, J. & Stamou, G. P. 1993[a]: Analysis of the vertical distribution of epiphytic lichens on *Pinus nigra* (Mount Olympos, Greece) along an altitudinal gradient. – Vegetatio **109**: 63–70.
- , — & — 1995: Analysis of the distribution of epiphytic lichens within homogeneous *Fagus sylvatica* stands along an altitudinal gradient (Mount Olympos, Greece). – Vegetatio **116**: 33–40.
- , Vokou, D., Diamantopoulos, J. & Galloway, D. J. 1993[b]: An assessment of the sampling procedure for estimating air pollution using epiphytic lichens as indicators. – Lichenologist **25**: 165–173.
- Pišút, I. 1968: Lichenologische Bemerkungen 3. – Annot. Zool. Bot. **50**: 1–9.
- Politis, J. 1953: Contribution à l'étude de la flore de la Chalcidique. Lichenes. – Pragmateiai

- Akad. Athinon **19**: 88–97.
- Räsänen, V. 1944: Griechische Flechten. Gesammelt von Dr. phil. C. Regel in den Jahren 1931–1939. – *Hedwigia* **81**: 229–233.
- Rechinger, K. 1915: Beiträge zur Kryptogamenflora der Insel Korfu II. Teil IV. Lichenes von J. Steiner. – *Verh. Zool.-Bot. Ges. Wien* **65**: 184–207.
- Rechinger, K. H. 1943: Flora Aegea. Flora der Inseln und Halbinseln des ägäischen Meeres. – *Denkschr. Akad. Wiss. Wien, Math.-Naturwiss. Kl.* **105(1)**.
- 1951: Neue Beiträge zur Flora von Kreta. – *Denkschr. Akad. Wiss. Wien, Math.-Naturwiss. Kl.* **105(2)**.
- Rondon, Y. 1969: Contribution à l'étude des lichens de l'île de Crète. – *Rev. Fac. Ci. Univ. Lisboa, Sér. 2, C, Ci. Nat.* **16**: 105–117.
- 1970: Contribution à l'étude des lichens du Péloponèse. – *Portugalia Acta Biol., Sér. B, Sist.* **11**: 38–50.
- Servít, M. 1933: Bearbeitung der von K. H. Rechinger (fil.) im Jahre 1927 auf den Ägäischen Inseln gesammelten Flechten. – *Ann. Naturhist. Mus. Wien* **46**: 77–90.
- 1935: Beitrag zur Flechtenflora der Ionischen Inseln und des nordwestlichen Peloponnes. – *Repert. Spec. Nov. Regni Veg.* **38**: 60–72.
- Sipman, H. J. M. & Raus, Th. 1995: Lichen observations from Santorini (Greece). – *Bibl. Lichenol.* **57**: 409–428.
- Steiner, J. 1898: Prodromus einer Flechtenflora des griechischen Festlandes. – *Sitzungsber. Kaiserl. Akad. Wiss., Math.-Naturwiss. Kl., Abt. 1*, **107**: 103–189.
- 1919: Beiträge zur Kenntnis der Flora Griechenlands. Bearbeitung der anlässlich der zweiten Wiener Universitätsreise im April 1911 in Griechenland gesammelten Pflanzen. C. Lichenes. – *Verh. Zool.-Bot. Ges. Wien* **69**: 52–101.
- Szatala, Ö. 1940: Adatok a Balkánfélsziget és Kis-Ázsia zuzmóflórájának ismeretéhez. Contributions à la connaissance de la flore lichénologique de la péninsule des Balkans et de l'Asie Mineure. – *Borbásia* **2**: 33–50.
- 1941: Adatok Görögország zuzmóflórájához I. Athos-félsziget (Hagion Oros). Contributions à la connaissance des lichens de la Grèce I. La presqu'île Athos (Hagion Oros). – *Borbásia* **3**: 113–136.
- 1959: Contributions à la connaissance des lichens de la Grèce. II. Mont Olympe. – *Ann. Hist.-Nat. Mus. Natl. Hung.* **51**: 121–144.
- Unger, F. 1862: Wissenschaftliche Ergebnisse einer Reise in Griechenland und in den jonischen Inseln. – *Wien*.
- Walter, E. 1976: Einige Beobachtungen an Flechten während der 15. IPE in Griechenland. – *Veröff. Geobot. Inst. ETH Stiftung Rübel Zürich* **56**: 87–91.
- Wilmanns, O. 1959: Epiphytengesellschaften Nordgriechenlands im Vergleich mit denen Mitteleuropas. – *Phyton* **8**: 175–182.
- & Phitos, D. 1960: Peri ton lichinon tis Parnithos. Zur Epiphytenflora des Parnes. – *Dassika Hron.* **18–19**: 837–841.
- Zahlbruckner, A. 1906: Beitrag zur Flechtenflora Kretas. – *Sitzungsber. Kaiserl. Akad. Wiss., Math.-Naturwiss. Cl., Abt. 1, Wien* **115(1)**: 503–523.
- 1907: Aufzählung der von Dr. H. Bretzl in Griechenland gesammelten Flechten. – *Hedwigia* **47**: 60–65.

Addresses of the authors:

Steen N. Christensen, Botanisk Museum, Gothersgade 130, DK-1123 København K, Denmark.
 Ivan Pišúit, Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 14, 84223 Bratislava, Slovakia.
 Harrie J. M. Sipman, Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin, Königin-Luise-Strasse 6–8, D-14191 Berlin; e-mail: hsipman@zedat.fu-berlin.de.