

## **Isolation of 18 Microsatellite Loci in the Desert Mistletoe *Phoradendron californicum* (Santalaceae) Via 454 Pyrosequencing**

Authors: Arroyo, Juan M., Munguia-Vega, Adrian, Rodríguez-Estrella, Ricardo, and Bascompte, Jordi

Source: Applications in Plant Sciences, 1(12)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1300048>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# ISOLATION OF 18 MICROSATELLITE LOCI IN THE DESERT MISTLETOE *PHORADENDRON CALIFORNICUM* (SANTALACEAE) VIA 454 PYROSEQUENCING<sup>1</sup>

JUAN M. ARROYO<sup>2</sup>, ADRIAN MUNGUÍA-VEGA<sup>3,4,5</sup>, RICARDO RODRÍGUEZ-ESTRELLA<sup>4</sup>,  
AND JORDI BASCOMPTE<sup>2,6</sup>

<sup>2</sup>Department of Integrative Ecology, Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas (CSIC), Calle Américo Vespucio s/n, E-41092 Sevilla, Spain; <sup>3</sup>Conservation Genetics Laboratory, School of Natural Resources and Environment, University of Arizona, BSE-317, 1311 E. 4th St., Tucson, Arizona 85721 USA; <sup>4</sup>Environmental Planning and Conservation Program, Centro de Investigaciones Biológicas del Noroeste (CIBNOR), Mar Bermejo 195, Col. Playa Palo de Santa Rita, La Paz, Baja California Sur 23090, Mexico; and <sup>5</sup>Comunidad y Biodiversidad A.C., Boulevard Agua Marina 297, Colonia Delicias, Guaymas, Sonora 85420, Mexico

- **Premise of the study:** Microsatellite primers were developed for the parasitic mistletoe *Phoradendron californicum* to investigate to what extent population genetic structure depends on host tree distribution within a highly fragmented landscape.
- **Methods and Results:** Fourteen unlinked polymorphic and four monomorphic nuclear microsatellite markers were developed using a genomic shotgun pyrosequencing method. A total of 187 alleles plus four monomorphic loci alleles were found in 98 individuals sampled in three populations from the Sonoran Desert in the Baja California peninsula (Mexico). Loci averaged 13.3 alleles per locus (range 4–28), and observed and expected heterozygosities within populations varied from 0.167–0.879 and 0.364–0.932, respectively.
- **Conclusions:** Levels of polymorphism of the reported markers are adequate for studies of diversity and fragmentation in natural populations of this parasitic plant. Cross-species amplifications in *P. juniperinum* and *P. diguetianum* only showed four markers that could be useful in *P. diguetianum*.

**Key words:** mistletoe; parasitic plant; *Phoradendron californicum*; shotgun sequencing; SSR; toji.

Mistletoes are parasitic plants dispersed by animals, mainly birds (Watson, 2001). The mistletoe *Phoradendron californicum* Nutt. has a broad distribution in North America and is a native parasitic plant of leguminous plants in the Sonoran Desert, mainly *Prosopis articulata* S. Watson and *Cercidium microphyllum* (Torr.) Rose & I. M. Johnst. The interactions among mistletoes, their host plants, and their animal dispersers are currently under study in both natural and fragmented landscapes in the desert of the Baja California peninsula. Mistletoe–host plants’ antagonistic interactions, along with animal dispersers, may form complex networks whose function and structure can be influenced by fragmentation at different scales, e.g., molecular or populations. Thus, the mistletoe–host interaction is ideal for studies on the effects of landscape fragmentation on plant–animal interactions.

Our aim is to determine to what extent landscape fragmentation affects mistletoe–animal interactions, the dispersal process of seeds, and the resulting networks of spatial genetic variability. For these purposes, we isolated and characterized nuclear microsatellite markers that are being successfully applied to describe spatial patterns of genetic structure. To date, microsatellite primers have not been developed for this mistletoe species.

## METHODS AND RESULTS

We extracted genomic DNA using a DNeasy Plant Mini Kit (QIAGEN, Valencia, California, USA) and used ~5 µg from one *P. californicum* individual collected in Santo Domingo Valley, Baja California Sur, Mexico (see Patch 33 population in Appendix 1 for GPS coordinates and voucher), to construct an enriched genomic library digested with *RsaI* (Sigma-Aldrich Corporation, St. Louis, Missouri, USA) for microsatellite loci using the oligonucleotide probes (GT)<sub>15</sub>, (CT)<sub>15</sub>, (GATA)<sub>10</sub>, (GACA)<sub>8</sub>, and (GATGT)<sub>5</sub> (Glenn and Schable, 2005). We sequenced a total of 288 *Escherichia coli* clones, from which 65 (22.5%) contained microsatellite loci but only 17 (5.9%) had flanking sequences adequate for the design of PCR primers, using Primer3 software (Rozen and Skaletsky, 2000). From these, only one polymorphic locus (Phca63B) was successfully amplified and scored. To overcome the low efficiency of enrichment and cloning methods, we used ~5 µg of genomic DNA from the same individual to construct a shotgun genomic library that was sequenced on 1/8th of a plate using 454 GS FLX Titanium chemistry (Roche Applied Science, Indianapolis, Indiana, USA) at the University of Arizona Genetics Core (Tucson, Arizona, USA). We generated 70.1 Mb of quality-filtered data, distributed over 196,512 unique reads with an average length of 357 bp after quality filtering (quality score [Q] ≥ 20 using a 10-bp sliding window). We located microsatellite

<sup>1</sup>Manuscript received 31 May 2013; revision accepted 8 August 2013.

The authors thank Melanie Culver for laboratory support and P. Jordano for useful comments and suggestions. For primer testing, logistical support was provided by the Molecular Ecology Laboratory, Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas (LEM-EBD-CSIC). This study was supported by grants from the Secretaría de Educación Pública, Consejo Nacional de Ciencia y Tecnología (SEP-CONACYT 155956 to R.R.E.) and the European Research Council (Advanced Grant to J.B.).

<sup>6</sup>Author for correspondence: bascomp@ebd.csic.es

doi:10.3732/apps.1300048

*Applications in Plant Sciences* 2013 1(12): 1300048; <http://www.bioone.org/loi/apps> © 2013 Arroyo et al. Published by the Botanical Society of America.  
This work is licensed under a Creative Commons Attribution License (CC-BY-NC-SA).

TABLE 1. Characteristics of 18 microsatellite markers isolated from populations of *Phoradendron californicum*.

| Locus <sup>a</sup> | Primer sequences (5'–3')                                    | Repeat motif   | Allele size range (bp) | A               | T <sub>a</sub> (°C) | Dye <sup>b</sup> | Mix no. for capillary electrophoresis <sup>c</sup> |
|--------------------|---|--|------------------------|-----------------|---------------------|------------------|--|
| PhcaWDS            | F: TGATGATGTTACCCAATAGGCA<br>R: CATAATTCACACCTTAGTGGCA      | (AT) <sub>10</sub>   | 193–219                | 12              | 53                  | FAM              | 1  |
| PhcaIK7            | F: AATCCAAACCACTTCATTATCTTT<br>R: TGAATGGTGGTGATTAATCATAAAA | (TAT) <sub>8</sub>   | 137–191                | 14              | 53                  | VIC              | 1  |
| PhcaK5R            | F: GGCCACCATAGGGAATGATA<br>R: TTACCCCATCAATGGTTTCC          | (AT) <sub>12</sub>   | 200–235                | 15              | 55                  | VIC              | 1  |
| Phca54T            | F: CATTCAACTGATGGATTCCAAA<br>R: GAAATCATTAATAGCATGTCCA      | (ATT) <sub>8</sub>   | 257–322                | 18              | 53                  | VIC              | 1  |
| Phca0Q7            | F: GCACTACTATGATTACAATGACAACA<br>R: TTTTGTAGTGGCAGTAGTTGTGG | (CAA) <sub>8</sub>   | 178–190                | 4               | 53                  | NED              | 1  |
| PhcaZ7Y            | F: AAAGTGAATGTTGGAGGAATTTTT<br>R: TTTGTGTCGAAGATATGTAAATGG  | (GAT) <sub>13</sub>  | 166–222                | 24 <sup>d</sup> | 55                  | PET              | 1  |
| Phca0GH            | F: TAGAACTCCCTCAGACCCCA<br>R: TGCATCAATAAGGATGGGATG         | (TTA) <sub>9</sub>   | 147–212                | 24 <sup>d</sup> | 55                  | FAM              | 2  |
| PhcaREF            | F: ACATAATCATGGGCCGAAAA<br>R: TGGACATTCCATGTCATCA           | (AT) <sub>11</sub>   | 258–280                | 6               | 56                  | FAM              | 2  |
| PhcaCC3            | F: CCTTACAGGCCCATCATTCT<br>R: TAACAGGTGCGCATTTGGCATA        | (CAA) <sub>8</sub>   | 288–308                | 9 <sup>d</sup>  | 54                  | FAM              | 2  |
| Phca63B            | F: CCAAGGACTTTAAACCTCATTCC<br>R: GAGGAAATTATCATCATCCCGTAG   | (GC) <sub>5</sub> (CA) <sub>4</sub> CG(CA) <sub>14</sub> (TA) <sub>5</sub> | 109–159                | 12              | 56                  | VIC              | 2  |
| PhcaUT7            | F: TCAATGTTTGCAAAACAAACA<br>R: GGTGTTTTTACATCAATAACAATGA    | (AAT) <sub>8</sub>   | 276–300                | 7               | 54                  | VIC              | 2  |
| Phca1HP            | F: TGATTTTTTCGGTTTCATGGTT<br>R: GGGTGCCACACATCACAAAT        | (TAA) <sub>10</sub>  | 179–277                | 28              | 53                  | NED              | 2  |
| PhcaGQL            | F: TATTGCAATGGGACAAGGCT<br>R: GTTTCGAACCTTGAGACCCCA         | (TTG) <sub>8</sub>   | 197–246                | 9               | 56                  | PET              | 2  |
| Phca1G8            | F: TTCAAGCTTATCCCCACAGC<br>R: CCAATACAATAGACCCTAAAAACAA     | (AC) <sub>12</sub>   | 271–279                | 5               | 53                  | PET              | 2  |
| PhcaK4A            | F: TCTTGTGTTTTCATCGGTGC<br>R: TCAGCAATCCCTCCATTAC           | (AT) <sub>10</sub>   | 120                    | 1               | 55                  | NED              | 3  |
| PhcaW15            | F: TGAGGAGTGTACTCACCCATGT<br>R: CAACATACCACCAGAGATTG        | (AT) <sub>12</sub>   | 105                    | 1               | 55                  | VIC              | 3  |
| PhcaBVY            | F: ATCATAAAACCATGCTCGGG<br>R: GAAACTTGATTGTGATGTTCTAAGGA    | (AT) <sub>10</sub>   | 129                    | 1               | 55                  | PET              | 3  |
| PhcaSRK            | F: TTCCAGACATGGAGTCGAGA<br>R: AAGACCGGAGCTATCAACCA          | (AT) <sub>10</sub>   | 285                    | 1               | 55                  | PET              | 3  |

Note: A = number of alleles; T<sub>a</sub> = annealing temperature (provided for nontailed primers).

<sup>a</sup>Sequences of 17 loci from the shotgun genomic library were submitted to the Sequence Read Archive (SRA) of the National Center for Biotechnology Information (NCBI; SRP015980 study accession). Locus Phca63B has a JX961681 GenBank accession number.

<sup>b</sup>PCR products were labeled using FAM, VIC, NED, or PET (Applied Biosystems) dyes on an additional 19 bp M13 primer (5'-CACGA-CGTTGTAAAACGAC-3') according to the methods of Boutin-Ganache et al. (2001). Moreover, a palindromic sequence tail (5'-GTGTCTT-3') was added to the 5' end of the reverse primer to improve adenylation and facilitate genotyping.

<sup>c</sup>Mix number indicates loci that were mixed in the same capillary electrophoresis run.

<sup>d</sup>Loci PhcaZ7Y, Phca0GH, and PhcaCC3 showed several alleles (6, 4, and 2, respectively) with a size difference of 2 or 1 bp relative to contiguous alleles; however, no variation outside of the microsatellite region was showed by these loci, and no scoring errors were detected by MICRO-CHECKER software (van Oosterhout et al., 2004). These polymorphisms could result from nonstepwise mutations in the repeat array.

loci containing at least 10 perfect repeats and designed primers using the software QDD (Megléc et al., 2010). We used unique sequence reads and consensus sequences within contigs that grouped sequences with a similarity ≥95% in regions ≥30 bp flanking the repeats to design primers. This step eliminated duplicated loci that have diverged in the flanking regions (except recent duplicates) and reduced null alleles by assembling sequence data from regions with coverage ≥1× for anchoring primers.

We found 115 di-, 376 tri-, and 13 tetranucleotide loci that met our criteria, and selected all of the tetranucleotides, 24 dinucleotides, and 24 trinucleotides with the largest number of repeats for primer synthesis. For primer testing, DNA was isolated from silica-dried stems using a modified cetyltrimethylammonium bromide (CTAB) extraction method (Milligan, 1998) that included tissue grinding in a Mixer Mill MM301 (Retsch, Haan, Germany) and TLE resuspension (10 mM Tris-HCl [pH 8.0], 0.1 mM EDTA). We sampled a total of 98 *P. californicum* individuals from three natural populations located in the Sonoran Desert in the Baja California peninsula, Mexico (Appendix 1). Sampling was designed to capture at least one sample from each parasitized host tree in each patch. Hosts were more aggregated in patch 32 compared to other patches.

PCR amplifications were performed in a 20-μL final volume containing 1× buffer (67 mM Tris-HCl [pH 8.8], 16 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.01% Tween-20), 2.5 mM MgCl<sub>2</sub> (1.5 mM for locus PhcaGQL), 0.01% bovine serum albumin (BSA; Roche Diagnostics, Barcelona, Spain), 0.25 mM dNTP, 0.40 μM dye-labeled M13 primer, 0.40 μM “pig-tailed” reverse primer, 0.04 μM M13-tailed forward primer (see M13 and “pig-tail” sequences in Table 1), 0.5 U *Taq* DNA polymerase (Bioline, London, United Kingdom), and approximately 50–70 ng genomic DNA. Reactions were undertaken in a “touchdown” PCR protocol in a Bio-Rad DNA Engine Peltier Thermal Cycler (Bio-Rad Laboratories, Hercules, California, USA), with an initial 2 min of denaturation at 94°C; 17 cycles at 92°C for 30 s, annealing at 60–44°C for 30 s (1°C decrease in each cycle), and extension at 72°C for 30 s; 25 cycles at 92°C for 30 s, 44°C for 30 s, and 72°C for 30 s; and a final extension of 5 min at 72°C. Amplified fragments were analyzed on an ABI 3130xl Genetic Analyzer (Applied Biosystems, Foster City, California, USA) and sized using GeneMapper 4.0 (Applied Biosystems) and GeneScan 500 LIZ size standard (Applied Biosystems). No multiplexing was attempted at the PCR stage (see mix for capillary electrophoresis in Table 1).

From a total of 61 loci tested from the next-generation library, four were monomorphic, 27 showed complex or nonspecific amplification, six showed

TABLE 2. Results of screening of 14 polymorphic markers in three populations (Patches 32, 33, and 59) of *Phoradendron californicum*.<sup>a</sup>

| Locus   | Patch 32 |          |                       |                       |                      | Patch 33 |          |                       |                       |                      | Patch 59 |          |                       |                       |                      |
|---------|----------|----------|-----------------------|-----------------------|----------------------|----------|----------|-----------------------|-----------------------|----------------------|----------|----------|-----------------------|-----------------------|----------------------|
|         | <i>n</i> | <i>A</i> | <i>H</i> <sub>o</sub> | <i>H</i> <sub>e</sub> | HWE                  | <i>n</i> | <i>A</i> | <i>H</i> <sub>o</sub> | <i>H</i> <sub>e</sub> | HWE                  | <i>n</i> | <i>A</i> | <i>H</i> <sub>o</sub> | <i>H</i> <sub>e</sub> | HWE                  |
| PhcaWDS | 30       | 8        | 0.600                 | 0.789                 | 0.0255 <sup>†</sup>  | 33       | 9        | 0.636                 | 0.752                 | 0.0180               | 33       | 7        | 0.727                 | 0.804                 | 0.1850               |
| PhcaIK7 | 31       | 9        | 0.710                 | 0.843                 | 0.0340               | 33       | 10       | 0.848                 | 0.836                 | 0.9373               | 34       | 12       | 0.706                 | 0.810                 | 0.3190               |
| PhcaK5R | 31       | 10       | 0.677                 | 0.809                 | 0.0437               | 33       | 11       | 0.636                 | 0.778                 | 0.2530               | 34       | 12       | 0.647                 | 0.864                 | 0.0023* <sup>†</sup> |
| Phca54T | 31       | 15       | 0.710                 | 0.915                 | 0.0004* <sup>†</sup> | 33       | 15       | 0.879                 | 0.918                 | 0.1333               | 34       | 10       | 0.824                 | 0.864                 | 0.1622               |
| PhcaOQ7 | 30       | 2        | 0.267                 | 0.364                 | 0.1578               | 30       | 2        | 0.167                 | 0.440                 | 0.0012* <sup>†</sup> | 33       | 4        | 0.212                 | 0.518                 | 0.0001* <sup>†</sup> |
| PhcaZ7Y | 30       | 14       | 0.300                 | 0.883                 | 0.0000* <sup>†</sup> | 33       | 14       | 0.485                 | 0.886                 | 0.0000* <sup>†</sup> | 34       | 18       | 0.500                 | 0.921                 | 0.0000* <sup>†</sup> |
| Phca0GH | 31       | 17       | 0.774                 | 0.916                 | 0.0706               | 33       | 17       | 0.697                 | 0.902                 | 0.0004* <sup>†</sup> | 34       | 15       | 0.500                 | 0.910                 | 0.0000* <sup>†</sup> |
| PhcaREF | 31       | 4        | 0.548                 | 0.668                 | 0.0570               | 33       | 6        | 0.545                 | 0.726                 | 0.0072 <sup>†</sup>  | 34       | 4        | 0.529                 | 0.597                 | 0.0461               |
| PhcaCC3 | 30       | 7        | 0.533                 | 0.733                 | 0.0334 <sup>†</sup>  | 31       | 7        | 0.516                 | 0.754                 | 0.0004* <sup>†</sup> | 33       | 6        | 0.545                 | 0.661                 | 0.0837               |
| Phca63B | 31       | 7        | 0.323                 | 0.540                 | 0.0025* <sup>†</sup> | 33       | 8        | 0.576                 | 0.700                 | 0.1299               | 34       | 8        | 0.382                 | 0.744                 | 0.0000* <sup>†</sup> |
| PhcaUT7 | 31       | 5        | 0.355                 | 0.583                 | 0.0250 <sup>†</sup>  | 33       | 5        | 0.455                 | 0.577                 | 0.3080               | 34       | 5        | 0.500                 | 0.562                 | 0.3970               |
| Phca1HP | 31       | 17       | 0.806                 | 0.924                 | 0.0872               | 33       | 17       | 0.636                 | 0.932                 | 0.0014* <sup>†</sup> | 34       | 18       | 0.500                 | 0.918                 | 0.0000* <sup>†</sup> |
| PhcaGQL | 31       | 7        | 0.355                 | 0.682                 | 0.0001* <sup>†</sup> | 32       | 7        | 0.531                 | 0.798                 | 0.0001* <sup>†</sup> | 34       | 8        | 0.618                 | 0.847                 | 0.0250 <sup>†</sup>  |
| Phca1G8 | 31       | 5        | 0.548                 | 0.764                 | 0.0013* <sup>†</sup> | 33       | 5        | 0.515                 | 0.745                 | 0.0000* <sup>†</sup> | 34       | 4        | 0.353                 | 0.671                 | 0.0002* <sup>†</sup> |

Note: *A* = number of alleles; *H*<sub>e</sub> = expected heterozygosity; *H*<sub>o</sub> = observed heterozygosity; HWE = nominal *P* values for the test of deviations from Hardy–Weinberg equilibrium; *n* = number of individuals successfully genotyped.

<sup>a</sup> See Appendix 1 for GPS coordinates and voucher information.

\* Locus showed significant deviation from Hardy–Weinberg equilibrium (Bonferroni-corrected, *P* < 0.0036).

<sup>†</sup> Locus showed signs for the presence of null alleles.

high frequencies of null alleles, and 11 failed to amplify. Therefore, we obtained a total of 13 polymorphic loci, adding to a total of 14 polymorphic loci among both libraries (Table 1). We observed a total of 187 alleles for our *P. californicum* sample, averaging overall 13.3 alleles per locus (range 4–28; see Table 2 for heterozygosities). The combined probability of individual identity based on the 14 polymorphic loci was estimated at  $1.42 \times 10^{-17}$  with CER-VUS (Kalinowski et al., 2007).

In addition, we tested the 14 polymorphic and the four additional monomorphic loci on 12 and eight individuals of two additional species of *Phoradendron*: *P. diguetianum* Tiegh. and *P. juniperinum* A. Gray (Appendix 1). These species were selected because (1) they belong to the same genus; (2) they are also distributed in the Sonoran Desert, and *P. diguetianum* occurs in the same sampling area; and (3) both species are common and, therefore, easy to collect. Sampling was designed to capture a wider range of potential variation among *Phoradendron* species, covering a more extensive geographic area. DNA isolation of *P. diguetianum* samples was performed as described for *P. californicum*, whereas *P. juniperinum* samples were extracted with the DNeasy Plant Mini Kit (QIAGEN). PCR conditions were as described above for *P. californicum* except for labeling PCR products of the 14 polymorphic loci (see Table 3). Three positive and one negative control were run with each PCR to ensure accurate scoring. *Phoradendron diguetianum* showed four loci (Phca0GH, PhcaCC3, Phca63B, and PhcaW15) with ambiguous locus-specific amplification, and the loci PhcaGQL, PhcaK4A, and PhcaSRK were monomorphic; *P. juniperinum* showed no PCR amplifications.

Table 2 summarizes the features of the 14 loci after inspecting their number of alleles, observed and expected heterozygosities (CERVUS 3.0.3; Kalinowski et al., 2007), and testing for deviations from Hardy–Weinberg equilibrium (HWE; Arlequin 3.5.1.3; Excoffier and Lischer, 2010), gametic disequilibrium (GENEPOP 4.1.4; Roussel, 2008), and the presence of null alleles according to the van Oosterhout method (MICRO-CHECKER 2.2.3; van Oosterhout et al., 2004). We used Bonferroni-corrected *P* values to assess the significance of the results obtained. Several loci among populations showed a significant deviation from HWE (Bonferroni-corrected *P* < 0.05/14 = 0.0036) due to a deficit of heterozygote individuals (Table 2). Only loci PhcaZ7Y and Phca1G8 showed this deviation in all three populations, and PhcaOQ7, Phca0GH, Phca63B, Phca1HP, and PhcaGQL showed deviations in two populations. Significant gametic disequilibrium (*P* < 0.0036) was detected for two pairs of loci, but patterns were not consistent among populations (only Phca0GH–PhcaCC3 in population 33 and Phca0GH–Phca54T in population 59 were significant). Signs for the presence of null alleles were suggested by the general excess of homozygotes in eight loci for each population (Table 2). Thus PhcaZ7Y, Phca1G8, and PhcaGQL consistently showed null allele frequencies between 0.128–0.320 in all three populations. In contrast, loci PhcaOQ7, Phca0GH, Phca63B, Phca1HP, and PhcaCC3 showed evidence of null alleles at frequencies ≤ 0.2576 in two populations. These results were not unexpected and could stem from the characteristic pattern of aggregation of parasitic plants, imposing limitations to pollen- and seed-mediated gene flow within aggregations and reduced genetic

TABLE 3. Cross-species amplification testing of *Phoradendron californicum* microsatellite markers in *P. juniperinum* and *P. diguetianum*.<sup>a</sup>

| Marker name <sup>b</sup> | Species                            |                       |
|--------------------------|------------------------------------|-----------------------|
|                          | <i>P. diguetianum</i> <sup>c</sup> | <i>P. juniperinum</i> |
| Fam PhcaWDS (174–200)    | NSA                                | NA                    |
| Pet PhcaIK7 (118–172)    | NA                                 | NA                    |
| Vic PhcaK5R (181–216)    | NA                                 | NA                    |
| Ned Phca54T (238–303)    | NSA                                | NA                    |
| Fam PhcaOQ7 (159–171)    | MP (90–200)                        | NA                    |
| Pet PhcaZ7Y (147–203)    | NA                                 | NA                    |
| Ned Phca0GH (128–193)    | Null <sup>d</sup> (116–222)        | NA                    |
| Pet PhcaREF (239–261)    | NSA                                | NA                    |
| Vic PhcaCC3 (269–289)    | LSA <sup>e</sup> (143–219)         | NA                    |
| Vic Phca63B (90–140)     | LSA <sup>f</sup> (173–174)         | NA                    |
| Vic PhcaUT7 (257–281)    | NA                                 | NA                    |
| Ned Phca1HP (160–258)    | MP (92–201)                        | NA                    |
| Vic PhcaGQL (178–227)    | Monomorphic (235)                  | NA                    |
| Pet Phca1G8 (252–260)    | NA                                 | NA                    |
| Ned PhcaK4A (120*)       | Monomorphic (120)                  | NA                    |
| Vic PhcaW15 (105*)       | LSA <sup>g</sup> (97–109)          | NA                    |
| Pet PhcaBVY (129*)       | NA                                 | NA                    |
| Pet PhcaSRK (285*)       | Monomorphic (272)                  | NA <sup>h</sup>       |

Note: LSA = microsatellite locus-specific amplification; MP = multiple peaks (>2) were observed for each individual; NA = no amplification; NSA = nonspecific amplification was evidenced by the presence of several peaks in all samples with an unscorable pattern; Null = high frequency of null alleles.

<sup>a</sup> PCR products were labeled using FAM, VIC, NED, or PET (Applied Biosystems) dyes on the 5' end of the forward primer. A sequence tail (5'-GTGTCTT-3') was added to the 5' end of the reverse primer.

<sup>b</sup> Values in parentheses represent the size range (in base pairs) in *P. californicum* using the labeled forward primer.

<sup>c</sup> Values in parentheses represent the size range (in base pairs) in *P. diguetianum* using the labeled forward primer.

<sup>d</sup> Five individuals failed to amplify and seven showed a total of six alleles that differed by 1 bp (in a 116–122 bp range).

<sup>e</sup> One individual showed 143/219-bp genotype. The remaining 11 individuals were 152-bp homozygotes.

<sup>f</sup> Six individuals showed 173/174-bp genotype, five of them 173/173, and one of them 174/174. All showed unscorable peaks in an 83–104 bp range.

<sup>g</sup> The number of alleles was five.

<sup>h</sup> Three individuals showed a 272-bp weak peak.

\* The size range was obtained using the 19-bp labeled M13 primer.



differentiation among individuals within these clumps. Consistent and significant deviations from HWE and gametic disequilibrium within but not among populations are unlikely to be caused by null alleles or physical proximity between pairs of loci, and could indicate the presence of marked genetic structure within sampled populations (Francois and Durand, 2010).

CONCLUSIONS

Observed levels of polymorphism suggest that the reported markers are adequate for characterizing spatial genetic structure at reduced spatial scales and for studying aggregation patterns in *Phoradendron* host-parasite relationships. Cross-species amplifications were unsuccessful in *P. juniperinum* and ambiguous in *P. diguetianum*, most likely due to high genetic divergence among the studied *Phoradendron* species.

LITERATURE CITED

BOUTIN-GANACHE, I., M. RAPOSO, M. RAYMOND, AND C. F. DESCHEPPER. 2001. M13-tailed primers improve the readability and usability of microsatellite analyses performed with two different allele-sizing methods. *BioTechniques* 31: 24–27.

EXCOFFIER, L., AND H. E. L. LISCHER. 2010. Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources* 10: 564–567.

FRANCOIS, O., AND E. DURAND. 2010. Spatially explicit Bayesian clustering models in population genetics. *Molecular Ecology Resources* 10: 773–784.

GLENN, T. C., AND N. A. SCHABLE. 2005. Isolating microsatellite DNA loci. *Methods in Enzymology* 395: 202–222.

KALINOWSKI, S. T., M. L. TAPER, AND T. C. MARSHALL. 2007. Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* 16: 1099–1106.

MEGLÉCZ, E., C. COSTEDOAT, V. DUBUT, A. GILLES, T. MALAUSA, N. PECH, AND J. F. MARTIN. 2010. QDD: A user-friendly program to select microsatellite markers and design primers from large sequencing projects. *Bioinformatics (Oxford, England)* 26: 403–404.

MILLIGAN, B. G. 1998. Total DNA isolation. In A. R. Hoelzel [ed.], *Molecular genetic analysis of populations: A practical approach*, 2nd ed., 43–44. Oxford University Press, Oxford, United Kingdom.

ROUSSET, F. 2008. GENEPOP’007: A complete re-implementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources* 8: 103–106.

ROZEN, S., AND H. SKALETSKY. 2000. Primer3 on the WWW for general users and for biologist programmers. In S. Misener and S. A. Krawetz [eds.], *Methods in molecular biology*, vol. 132: Bioinformatics methods and protocols, 365–386. Humana Press Inc., Totowa, New Jersey, USA.

VAN OOSTERHOUT, C., W. F. HUTCHINSON, D. P. M. WILLS, AND P. SHIPLEY. 2004. MICRO-CHECKER: Software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* 4: 535–538.

WATSON, D. M. 2001. Mistletoe: A keystone resource in forests and woodlands worldwide. *Annual Review of Ecology and Systematics* 32: 219–249.

APPENDIX 1. Voucher, number of individuals sampled, and location information for *Phoradendron* species in this study. Vouchers are deposited at the University of Arizona Herbarium, Tucson, Arizona (ARIZ), and the herbarium of the Centro de Investigaciones Biológicas del Noroeste, La Paz, Baja California Sur, Mexico (HCIB).

| Species                | Locality            | Population         | Latitude (N) | Longitude (W) | No. of individuals | Herbarium ID | Collection date  |
|------------------------|---------------------|--------------------|--------------|---------------|--------------------|--------------|------------------|
| <i>P. californicum</i> | Baja California Sur | Patch 32           | 25°12'53.81" | 111°42'28.97" | 31                 | HCIB 28108   | 5 February 2008  |
| <i>P. californicum</i> | Baja California Sur | Patch 33           | 25°12'48.34" | 111°41'52.21" | 33                 | HCIB 28109   | 5 February 2008  |
| <i>P. californicum</i> | Baja California Sur | Patch 59           | 25°06'13.67" | 111°41'47.93" | 34                 | HCIB 28110   | 5 February 2008  |
| <i>P. diguetianum</i>  | Baja California Sur | Los Planes         | 23°59'37.38" | 110°03'23.03" | 4                  | HCIB 28112   | 16 July 2012     |
| <i>P. diguetianum</i>  | Baja California Sur | Conquista Agraria  | 24°05'33.72" | 110°45'09.36" | 2                  | HCIB 28113   | 10 July 2012     |
| <i>P. diguetianum</i>  | Baja California Sur | Todos Santos       | 23°27'50.37" | 110°14'42.03" | 2                  | HCIB 28114   | 19 July 2012     |
| <i>P. diguetianum</i>  | Baja California Sur | Cancún             | 24°42'23.44" | 111°33'31.03" | 1                  | HCIB 28115   | 29 July 2012     |
| <i>P. diguetianum</i>  | Baja California Sur | San José Comondú   | 26°03'30.47" | 111°48'25.18" | 2                  | HCIB 28116   | 25 July 2012     |
| <i>P. diguetianum</i>  | Baja California Sur | Rancho Cantarranas | 24°50'56.00" | 111°04'46.00" | 1                  | HCIB 28117   | 9 June 2012      |
| <i>P. juniperinum</i>  | Arizona             | Tucson, AZ         | 32°22'50.61" | 110°57'34.57" | 8                  | ARIZ 398223  | 11 November 2009 |