# Correction to "Whole Genome Shotgun Sequences for Microsatellite Discovery and Application in Cultivated and Wild Macadamia (Proteaceae)"

Authors: Nock, Catherine J., Elphinstone, Martin S., Ablett, Gary, Kawamata, Asuka, Hancock, Wayne, et al.

Source: Applications in Plant Sciences, 5(8)

Published By: Botanical Society of America

URL: https://doi.org/10.3732/apps.1300089_ERR

Applications in Plant Sciences 2017 5(8): 1300089_ERR

ERRATUM

# CORRECTION TO "WHOLE GENOME SHOTGUN SEQUENCES FOR MICROSATELLITE DISCOVERY AND APPLICATION IN CULTIVATED AND WILD *MACADAMIA* (PROTEACEAE)"

CATHERINE J. NOCK, MARTIN S. ELPHINSTONE, GARY ABLETT, ASUKA KAWAMATA, WAYNE HANCOCK, CRAIG M. HARDNER, AND GRAHAM J. KING

Vol. 2 No. 4: 1300089, April 2014, doi:10.3732/apps.1300089

The authors discovered an error in a primer sequence reported in Table 1. The corrected primer sequence (locus Mac0010, reverse primer sequence) is listed below in boldface. The authors apologize for the error. The published article (full-text and PDF) has been corrected.

TABLE 1. Correction to: Characterization of 12 polymorphic microsatellite loci developed in *Macadamia integrifolia*.[a]

| Locus | Primer sequences (5′–3′) | Repeat motif | Fluorescent label | Allele size range (bp) | $T_a$ (°C) | GenBank accession no. |
|---|---|---|---|---|---|---|
| Mac001 | F: GTGACTGGTGGACACCAAAACCCA<br>R: GCACTAGGTGTCACCCCCACTTCT | $(AT)_{11}$ | VIC | 412–420 | 60 | KF130888 |
| Mac002 | F: CCCAACTGGGTTTGCAAGGACCAA<br>R: AGTAGCCGCGAGCTGATCGAAGAT | $(CT)_8$ | NED | 283–297 | 60 | KF130889 |
| Mac003 | F: TGGACCATTGAGGAGTTGGACTGT<br>R: TCCACCGTTTCACTTTCGTCAGCC | $(AT)_9$ | FAM | 258–276 | 60 | KF130890 |
| Mac004 | F: CAAGAGTGTCCAGCGAGGGAATGC<br>R: GGGAGACATCATACTTTTGACACATGCC | $(AT)_{11}$ | NED | 224–240 | 60 | KF130891 |
| Mac005 | F: CATAGCATGAGTTTCAAGGGATAA<br>R: ATTACAAACCCACTCTTCGATTT | $(AAG)_{10}$ | FAM | 331–343 | 60 | KF130892 |
| Mac006 | F: TTTCATCATTGATCATCATAGGTACA<br>R: GAGCTAATACTTAACCAGGTGAACA | $(AG)_{11}$ | PET | 322–360 | 55 | KF130893 |
| Mac007 | F: AGGCCTTGGGATGTTCCAGTGTGA<br>R: GCAATCAACACAAGCACCTGTGGC | $(CT)_{11}$ | NED | 368–390 | 60 | KF130894 |
| Mac008 | F: AACGGTTATGTCAAGTGCAACAGGA<br>R: TGACTTTAGCCCTCACTTCAAAGCCA | $(AT)_{10}$ | FAM | 388–398 | 60 | KF130895 |
| Mac009 | F: CAACTCTCTCTCCCTCAGATTCTC<br>R: TAAATCTATGCCACATCACTAGGC | $(AAG)_{13}$ | VIC | 241–244 | 60 | KF130896 |
| Mac010 | F: GCAACTGGATCAGCACATAAGAAT<br>R: **TCCGATCATAGTCTTAGCATTTCA** | $(AG)_{11}$ | PET | 259–297 | 55 | KF130897 |
| Mac011 | F: AGAGGGCGAGATCCCTGACTCTGA<br>R: TGAAATTTGGCGTGGGGAAAGCGT | $(CT)_9$ | FAM | 175–199 | 60 | KF130898 |
| Mac012 | F: TATCAGGACCATCAACAATGATTT<br>R: GCCTGTTGTAGGTAAAGTGGAGAT | $(AC)_{10}$ | VIC | 309–321 | 60 | KF130899 |

*Note*: $T_a$ = annealing temperature used for all *Macadamia* species and cultivars.
[a]Values based on 22 samples representing *Macadamia* cultivars located at Clunes Varietal Trial M2, New South Wales, Australia.

doi:10.3732/apps.1300089_ERR