



Development of Microsatellite Markers for the Coastal Shrub *Scaevola taccada* (Goodeniaceae)

Authors: Ando, Haruko, Emura, Naoko, Denda, Tetsuro, Nakahama, Naoyuki, Inoue-Murayama, Miho, et al.

Source: Applications in Plant Sciences, 2(5)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1300094>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

DEVELOPMENT OF MICROSATELLITE MARKERS FOR THE COASTAL SHRUB *SCAEVOLA TACCADA* (GOODENIACEAE)¹

HARUKO ANDO^{2,6}, NAOKO EMURA³, TETSURO DENDA⁴, NAOYUKI NAKAHAMA²,
MIHO INOUE-MURAYAMA⁵, AND YUJI ISAGI²

²Laboratory of Forest Biology, Division of Forest and Biomaterials Science, Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan; ³Department of Life Science, Rikkyo University, Tokyo 171-8501, Japan; ⁴Faculty of Science, University of the Ryukyus, Okinawa 903-0213, Japan; and ⁵Wildlife Research Center, Kyoto University, Kyoto 606-8203, Japan

- **Premise of the study:** Microsatellite markers were developed for the coastal shrub species *Scaevola taccada* to estimate the population genetic structure, which may reflect different seed dispersal patterns.
- **Methods and Results:** Thirteen microsatellite primer sets were developed for *S. taccada* using 454 pyrosequencing. The primer sets were tested on 64 individuals sampled from two populations in Japan. Fragments were amplified using the primers, with one to 10 alleles per locus, and the expected heterozygosity ranged from 0.00 to 0.85.
- **Conclusions:** These results indicate the utility of markers in *S. taccada* for broad estimations of the population genetic structure of this species.

Key words: 454 sequencing; genetic structure; Goodeniaceae; *Scaevola taccada*; seed dispersal.

Scaevola taccada (Gaertn.) Roxb. (Goodeniaceae) is a coastal shrub species widely distributed along coastal areas of the Pacific and Indian oceans (Howarth et al., 2003). The white fleshy exocarp and the underlying corky layer of the fruit (Howarth et al., 2003) allow it to be transported in the avian gut (Kawakami et al., 2009; Emura et al., 2012) and by oceanic floating (Nakanishi, 1988). Due to its unique seed dispersal system, *S. taccada* has been found growing in various environments. Seashores (sandy shores and cliffs, consistent with dispersal by oceanic floating and birds) represent the primary growth environment of this species, but some populations exist in island interiors (dispersal by birds) (Satake et al., 1989; Emura et al., unpublished data). Following such a distribution pattern, some isolated populations might be in the process of speciation, which may be indicated by population genetic structure. To estimate genetic differentiation caused by the different seed dispersal patterns, highly variable genetic markers are required. Here, we report 13 nuclear microsatellite loci for *S. taccada* developed using 454 next-generation sequencing, which will be useful for estimating the population genetic structure of this species.

¹Manuscript received 5 December 2013; revision accepted 5 February 2014.

This study was supported by a Grant-in-Aid from the Japan Society for the Promotion of Science Fellows. The Roche GS Junior Sequencer was operated by the Cooperation Research Program of the Wildlife Research Center at Kyoto University.

⁶Author for correspondence: ando.haruko.75a@st.kyoto-u.ac.jp

doi:10.3732/apps.1300094

METHODS AND RESULTS

A fresh leaf sample of *S. taccada* was collected from an individual growing in a coastal area of the Ogasawara Islands, Japan, and genomic DNA was extracted using the DNeasy Plant Mini Kit (QIAGEN, Germantown, Maryland, USA). The GS Junior Titanium Series Kit (Roche, Mannheim, Germany) and the SPRIworks Fragment Library System (Beckman Coulter, Brea, California, USA) were used for construction of a DNA library for *S. taccada*. A 500-ng aliquot of genomic DNA was nebulized at 0.24 MPa for 1 min, purified, end-repaired, and A-tailed using the SPRIworks Fragment Library Kit II (Beckman Coulter) and ligated to the Rapid Library Adapter (Roche) using RL Ligase (Roche). Suitably sized DNA fragments were selected by removing short fragments using SPRIworks Fragment Library Kit II (Beckman Coulter). Emulsion PCR (emPCR) was constructed for the desired fragments mixed with capture beads using the GS Junior Titanium emPCR Kit (Roche). After emPCR, the beads capturing the DNA library were enriched to selectively capture beads with sufficient amounts of template DNA for sequencing. The enriched beads were annealed with sequencing primers, and the amplified fragments were sequenced using the GS Junior Benchtop System (Roche). In the GS Junior sequencing, 30,497 DNA sequences were obtained. The sequences were screened to find potential microsatellite loci using the MSATCOMMANDER program (Faircloth, 2008). After screening, 423 repeat regions containing eight or more dinucleotide repeats were identified using Primer3 software (Rozen and Skaletsky, 2000) embedded in MSATCOMMANDER, and primers were successfully designed for a total of 72 repeats.

Twenty-three primer pairs were selected for amplification trials in eight *S. taccada* individuals, based on the repeat structure and avoiding sequences containing mononucleotide repeats. All of the forward primers designed from the selected loci were synthesized with a tag sequence (Boutin-Ganache et al., 2001) for fluorescent labeling. The sequence of each tag is indicated in the notes for Table 1. A modified protocol for the QIAGEN Multiplex PCR Kit (QIAGEN) was used for PCR amplification. The final volume of the PCR reaction mixture was 5 µL and contained the following: 16 ng extracted DNA, 2.5 µL Multiplex PCR Master Mix, 0.01 µM forward primer, 0.2 µM reverse primer, and 0.1 µM M13 (fluorescently labeled) primer. The PCR program was as follows: an initial denaturation at 95°C for 15 min; 30 cycles at

TABLE 1. Characteristics of 13 microsatellite primers developed for *Scaevola taccada*. All values are based on 64 samples from two populations on the Okinawa and Ogasawara islands in Japan.

Locus	Primer sequences (5′–3′)	Repeat motif	Fluorescent label ^a	T _a (°C)	Allele size range (bp)	GenBank accession no.
Stac05	F: TGGTGGTTAACAAGTGCCAAG R: GTTACTCAAGTCTCTTATTAGCAGTTC	(AG) ₈	FAM	57	175–179	AB872256
Stac06	F: GGCACCAGCTTCTGTAAACC R: GCAGTGGTGGATGCATTCTG	(AG) ₉	FAM	57	270–274	AB872257
Stac09	F: CATGAGCCAAGGAAACGTCC R: GCTTCGGCTCTTCACAAGG	(AG) ₉	VIC	57	325–335	AB872258
Stac10	F: CAGCAGGAATGCACTAAGACC R: GGAGGGATGGAAGGTCCG	(AT) ₁₁	VIC	57	223–253	AB872259
Stac11	F: TTTCGCAAGATCCCGGC R: TGGAAGTTGGTAAATGGGTCAG	(AT) ₈	NED	57	254	AB872260
Stac13	F: TCTTAACGACTCCTGTCACC R: CCGTCTCCATCCCTTCGTG	(AC) ₉	FAM	57	209	AB872261
Stac15	F: ACTCATCAGACAAGGTACGG R: AGCAGTCTACTTCCCACGC	(AC) ₁₀	VIC	57	368–372	AB872262
Stac17	F: CTGCAACGGTCATTTCGTCC R: TGCAACCTATATGCTATGTGCTC	(AT) ₉	NED	57	289–303	AB872263
Stac18	F: GAGGGAAGGTACAGAGGTGC R: TCAGACAAGTGAATAGAAGGTCATC	(AG) ₉	FAM	57	189–193	AB872264
Stac19	F: AAAGTCGGGCAGTAGGTGC R: GGAGTTGATGAATGCTCGGC	(AG) ₈	FAM	57	250	AB872265
Stac21	F: CCCTCAAGGCTACTGTGG R: TTGCGTTCTTCCGCAATCC	(CT) ₁₃	FAM	57	297–304	AB872266
Stac24	F: ATTGATTAGTAGGTGAGAATCGTG R: TCGGTATTGTCTTAATCTTCCG	(AT) ₁₀	NED	57	191–193	AB872267
Stac27	F: ATACATCATCGTACCCAAATCC R: GAGCTCCTGAAATGTCGCC	(AT) ₈	NED	57	199–203	AB872268

Note: T_a = annealing temperature.
^a Sequence of the fluorescent labels: FAM = 5′-CACGACGTTGTAAAACGAC-3′, NED = 5′-CTATAGGGCACGCGTGGT-3′, VIC = 5′-TGTGGAA-TTGTGAGCGG-3′.

94°C for 30 s, 57°C for 1.5 min, and 72°C for 1 min; and a final extension at 60°C for 30 min. The PCR product size was determined using the ABI PRISM 3100 Genetic Analyzer (Applied Biosystems, Foster City, California, USA) and GeneMapper software (Applied Biosystems). Thirteen primer pairs amplified well and could be scored easily and unambiguously (Table 1). These primer pairs were used for amplification in 64 individuals sampled from two populations located on the Ogasawara Islands (26.61184°N, 142.17543°E) and the Okinawa Islands (26.13975°N, 127.79644°E), using the PCR program used for amplification described above. One specimen from each population was collected and deposited at the Kyoto Univer-

sity Museum herbarium (accession numbers: KYO 00037380 and KYO 00037381).
The number of alleles for a given locus ranged from one to 10 (mean: 2.5). The observed and expected heterozygosities were 0.00–0.91 (mean: 0.23) and 0.00–0.85 (mean: 0.27), respectively (Table 2). Deviation from Hardy–Weinberg equilibrium (HWE) and the linkage disequilibrium (LD) between loci were tested using FSTAT version 2.9.3 (Goudet, 1995). Two loci (Stac21 and Stac24) exhibited a significant deviation from HWE (*P* < 0.05) in the Ogasawara population. There was no evidence of LD for any loci pairs. Fixation index was calculated using FSTAT and its values were –0.18–1.00.

TABLE 2. Results for primer screening of all samples for 13 microsatellite loci in two populations of *Scaevola taccada*.

Locus	Okinawa (<i>N</i> = 32)				Ogasawara (<i>N</i> = 32)			
	<i>A</i>	<i>H</i> _o ^a	<i>H</i> _e	<i>F</i> _{IS}	<i>A</i>	<i>H</i> _o	<i>H</i> _e	<i>F</i> _{IS}
Stac05	2	0.22	0.28	0.25	3	0.53	0.60	0.13
Stac06	3	0.44	0.46	0.07	2	0.41	0.36	–0.12
Stac09	4	0.22	0.20	–0.07	1	0.00	0.00	NA
Stac10	10	0.91	0.85	–0.05	4	0.44	0.48	0.11
Stac11	1	0.00	0.00	NA	1	0.00	0.00	NA
Stac13	1	0.00	0.00	NA	1	0.00	0.00	NA
Stac15	3	0.47	0.52	0.11	3	0.47	0.53	0.13
Stac17	2	0.03	0.03	0.00	5	0.53	0.65	0.20
Stac18	3	0.53	0.60	0.13	3	0.50	0.42	–0.18
Stac19	1	0.00	0.00	NA	1	0.00	0.00	NA
Stac21	3	0.08	0.47	0.84	3	0.06	0.15	0.58
Stac24	2	0.00	0.12	1.00	1	0.00	0.00	NA
Stac27	2	0.25	0.38	0.35	2	0.03	0.03	0.00

Note: *A* = number of alleles; *F*_{IS} = fixation index; *H*_e = expected heterozygosity; *H*_o = observed heterozygosity; *N* = sample size; NA = not applicable.
^a Numbers in boldface show deviations from Hardy–Weinberg equilibrium.

CONCLUSIONS

We have characterized 13 microsatellite loci for *S. taccada*. These microsatellite loci will be useful for estimating population genetic structure possibly resulting from the various seed dispersal patterns of *S. taccada*.

LITERATURE CITED

BOUTIN-GANACHE, I., M. RAPOSO, M. RAYMOND, AND C. F. DESCHEPPER. 2001. M13-tailed primers improve the readability and usability of microsatellite analyses performed with two different allele-sizing methods. *BioTechniques* 31: 24–26.
EMURA, N., K. KAWAKAMI, T. DEGUCHI, AND K. SONE. 2012. Potential role of frugivorous birds in the recovery process of forest vegetation after feral goat eradication in Mukojima Island, the Bonin Islands. *Journal of Forest Research* 17: 352–359.
FAIRCLOTH, B. C. 2008. MSATCOMMANDER: Detection of microsatellite repeat arrays and automated, locus-specific primer design. *Molecular Ecology Resources* 8: 92–94.
GOUDET, J. 1995. FSTAT: A computer program to calculate *F*-statistics, version 1.2. *Journal of Heredity* 86: 485–486.

- HOWARTH, D. G., M. H. GUSTAFSSON, D. A. BAUM, AND T. J. MOTLEY. 2003. Phylogenetics of the genus *Scaevola* (Goodeniaceae): Implication for dispersal patterns across the Pacific Basin and colonization of the Hawaiian Islands. *American Journal of Botany* 90: 915–923.
- KAWAKAMI, K., L. MIZUSAWA, AND H. HIGUCHI. 2009. Re-established mutualism in a seed-dispersal system consisting of native and introduced birds and plants on the Bonin Islands, Japan. *Ecological Research* 24: 741–748.
- NAKANISHI, H. 1988. Dispersal ecology of the maritime plants in the Ryukyu Islands, Japan. *Ecological Research* 3: 163–173.
- ROZEN, S., AND H. SKALETISKY. 2000. Primer3 on the WWW for general users and for biologist programmers. In S. Misener and S. A. Krawetz [eds.], *Methods in molecular biology*, vol. 132: Bioinformatics methods and protocols, 365–386. Humana Press, Totowa, New Jersey, USA.
- SATAKE, Y., H. HARA, S. WATARI, AND T. TOMINARI. 1989. Wild flowers of Japan—woody plants. Heibonsha, Tokyo, Japan [In Japanese].