

Primers to Amplify SNP Markers in *Epichloë canadensis*(Clavicipitaceae)

Authors: Sullivan, Terrence J., Bultman, Thomas L., and Schoolcraft, Jennifer

Source: Applications in Plant Sciences, 4(3)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1500078>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

PRIMER NOTE

PRIMERS TO AMPLIFY SNP MARKERS IN *EPICHLÖE CANADENSIS* (CLAVICIPITACEAE)¹

TERRENCE J. SULLIVAN^{2,4}, THOMAS L. BULTMAN³, AND JENNIFER SCHOOLCRAFT²
²School of Sciences, Indiana University Kokomo, Kokomo, Indiana 46904 USA; and ³Department of Biology, Hope College, Holland, Michigan 49423 USA

- **Premise of the study:** Primers were designed to produce short amplicons containing single-nucleotide polymorphisms (SNPs) in β -tubulin (*tubB*) and translation elongation factor 1- α (*tefA*) in *Epichloë canadensis* (Clavicipitaceae), an endophytic fungus of *Elymus canadensis* (Poaceae).
- **Methods and Results:** Primers to amplify regions of *tubB* and *tefA* containing suspected SNPs were designed and tested on individuals from six populations. Two *tubB* alleles were identified that differed by a single SNP, and three *tefA* alleles were identified that differed by a combination of two SNPs. All six populations tested were polymorphic for the *tefA* marker, and three of the populations were also polymorphic for the *tubB* marker. These primers are also predicted to amplify these regions in 11 additional epichloid species.
- **Conclusions:** Primers for short amplicons within *tubB* and *tefA* genes can be used to successfully genotype *E. canadensis*, making them useful markers for population genetic or landscape genomic studies.

Key words: Clavicipitaceae; *Elymus canadensis*; endophyte; *Epichloë canadensis*; high-resolution melt analysis; single-nucleotide polymorphism (SNP).

Cool-season grasses often harbor symbiotic endophytic fungi from the genus *Epichloë* (Fr.) Tul. & C. Tul. (Clavicipitaceae) in their aboveground tissue. Some *Epichloë* species have strong effects on their hosts by deterring herbivores, increasing drought resistance, and increasing their host's competitive ability (Clay, 1990). This can lead to significant impacts on their hosts and on their surrounding community (Clay and Holah, 1999). However, there can be significant variation in the interaction based on specific combinations of host–endophyte genotypes (Faeth, 2002; Shymanovich et al., 2014).

Molecular markers, including microsatellites and sequences for the genes β -tubulin (*tubB*) and translation elongation factor 1- α (*tefA*), have proven useful for both identifying and genotyping epichloid endophytes (Moon et al., 1999; Sullivan and Faeth, 2004; Takach and Young, 2014; Young et al., 2014). In this paper, we describe primers producing small amplicons suitable for high-resolution melt (HRM) analysis that can be used for genotyping *tubB* and *tefA* single-nucleotide polymorphisms (SNPs) in *Epichloë canadensis* N. D. Charlton & C. A. Young, an endophyte of Canada wildrye (*Elymus canadensis* L., Poaceae), a native grass widespread in North America.

¹Manuscript received 3 July 2015; revision accepted 20 October 2015.

The authors thank Mark McKone and Nancy Barker for access to the Carleton College Cowling Arboretum and the Iowa Department of Natural Resources for access to Hayden Prairie State Preserve. This research was supported by the National Science Foundation (NSF-IOS 1119775 to T.J.S. and T.L.B.).

⁴Author for correspondence: tjsull@iuk.edu

doi:10.3732/apps.1500078

METHODS AND RESULTS

Epichloë canadensis is a haploid, hybrid species containing sequences from both *E. amarillans* White (typically found in *Agrostis perennans* (Walter) Tuck.) and *E. elymi* Schardl & Leuchtman (found in several *Elymus* L. species) (Charlton et al., 2012). Primers were designed to amplify SNPs in the *E. amarillans*-derived *tubB* and *tefA* genes based on sequences described in Charlton et al. (2012) (*tefA*: JN886775; *tubB*: JN886778), using Primer3 (Untergasser et al., 2012) as implemented in NCBI/Primer-BLAST (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>). The primers were designed to produce an amplicon smaller than 300 bp, only amplify the *E. amarillans*-derived gene, and include the potential SNPs described by Charlton et al. (2012). Primer sequences and their characteristics are given in Table 1.

Genomic DNA used to test the primers came from Canada wildrye plants grown from seeds collected from six populations in August and September 2011. Population locations are provided in Appendix 1. A voucher specimen (PUL N17966), including seeds, from an individual from CC population was deposited in the Purdue University Kriebel Herbarium (West Lafayette, Indiana, USA). Seeds were germinated in potting soil at the Indiana University Kokomo campus until they developed multiple leaves. DNA was extracted from ~100 mg of leaf tissue using the FastDNA Spin Kit (MP Biomedicals, Santa Ana, California, USA) using their recommended protocol for plant material. Amplification of endophyte microsatellites from these samples found multiple alleles for at least one of the loci (data not shown), indicative of hybrid endophytes (Moon et al., 2004) like *E. canadensis*.

Initial HRM screening was done using a StepOne Real-Time PCR system (Applied Biosystems, Grand Island, New York, USA) and the MeltDoctor HRM Master Mix (Applied Biosystems), following the manufacturer's recommendations. Reactions were run in a total volume of 20 μ L with primer concentrations of 0.3 μ M. After an initial 10 min 95°C hot start, samples were cycled 40 times at 94°C for 15 s and 63°C for 60 s. The melt curve was generated immediately following amplification. After an initial denaturing stage of 95°C for 60 s, samples were annealed at 60°C and heated to 95°C using the continuous ramp mode with a 0.3% ramp rate. In this mode, the block warms at a constant rate throughout the melt curve step and readings are taken as quickly as possible. Melt curves were analyzed using High Resolution Melt software version 3.0.1 (Applied Biosystems). To ensure the consistency of the HRM genotyping, samples were run in duplicate to account for any potential variation

TABLE 1. Characteristics of primers designed for SNP amplification in *Epichloë canadensis*.

Locus	Primer sequences (5′–3′)	T _a (°C)	Allele size (bp)	Based on sequences from	GenBank accession no.
<i>tefA</i> HRM	F: TCAACCCGTCACCTGGTCTT	58.5	205	<i>tefA</i> -1 allele, JN886775	KT214347–KT214349
	R: GATGGTGATACCACGCTCAC	58.4			
<i>tubB</i> HRM	F: GAGCCCTGATTTTCGTACC	57.6	242	<i>tubB</i> -1 allele, JN886778	KT214345–KT214346
	R: TGCCCAATGAATGTGAGTT	55.8			

Note: T_a = annealing temperature.

in amplification and only samples with threshold cycle (C_t) values less than 30 were used. Typical difference plots for *tubB* and *tefA* amplicons are shown in Fig. 1. To verify the genetic variation described by the software, PCR products from 10 individuals including representatives of each identified genotype were cloned into the pCR 4.0 plasmid vector using the TOPO-TA cloning kit (Invitrogen, Carlsbad, California, USA) for both the *tefA* and *tubB* PCR products. Sanger sequencing of the clones was performed by Functional Biosciences (Madison, Wisconsin, USA). Initial sequencing of seven to 10 clones confirmed only a single allele was being amplified for each individual, and subsequent sequencing was performed directly from PCR products by Functional Biosciences. In total, 85 individuals were sequenced to confirm *tefA* variation, and 52 individuals were sequenced to confirm *tubB* variation. Sequence alignments




Fig. 1. Representative difference plots for the *tubB* (A) and *tefA* (B) SNP alleles of the *Epichloë amarillans*-derived gene in *E. canadensis*.

were constructed using the ClustalW algorithm in MEGA 5.2.2 (Tamura et al., 2011), and representatives for each genotype were used in a BLAST search to confirm their identity as *E. amarillans*-derived alleles in *E. canadensis*. No *E. elymi*-derived alleles were amplified by these primers.

An A/G SNP was found in the *tubB* amplicon. Direct sequencing found that both amplicons were most similar to the *E. canadensis* isolate CWR5 *tubB*-1 allele (JN886778), with one allele being identical and the other only differing by the SNP. Both alleles had 99% identity with the *E. amarillans tubB* gene (L78272). Representative sequences for the two alleles have been deposited in GenBank (KT214345–KT214346). Two SNPs were found in the *tefA* amplicon, an A/G SNP and an A/C SNP. From these two SNPs, three *E. amarillans*-derived alleles were found containing the SNP combinations AA (identical to *tefA* in *E. amarillans* strain E4668 [KP689563] in *Agrostis hyemalis* (Walter) Britton, Sterns & Poggenb.), GC (identical to *E. amarillans* E57 [KP689562] and *E. canadensis* CWR 34 *tefA*-1 [KF719188]), and AC (identical to *E. canadensis* isolate CWR 5 *tefA*-1 [JN886775]). Representative sequences for each allele have been deposited in GenBank (KT214347–KT214349). The *tefA* SNP was polymorphic in all six populations, and the *tubB* SNP was polymorphic in three of the populations. As *E. canadensis* is haploid, gene diversity (*H*) for each marker in each population was calculated using Arlequin (Excoffier et al., 2005) instead of observed and expected heterozygosity. *H* is equivalent to expected heterozygosity in diploid organisms, but can be used with any organism regardless of ploidy (Nei, 1987). The results are given in Table 2.

The ability of the primers to amplify other epichloid species was tested in silico using Primer-BLAST (Ye et al., 2012). The primer pairs were tested using sequences with the *Epichloë* taxon ID (taxid:5112) in the National Center for Biotechnology Information (NCBI) nr database. Eleven species, not including *E. amarillans* and *E. canadensis*, were identified where both primer sets are expected to match the target sequence exactly and produce an amplicon (Appendix 2).

CONCLUSIONS

Epichloë endophytes are important plant symbionts, and in this paper, we describe primers that can be used to describe genetic diversity within *E. canadensis* using SNPs in the *tubB* and *tefA* genes. Both markers are polymorphic and we expect them to be valuable for future population genetic and landscape genomics studies of epichloid endophytes of cool-season grasses.

LITERATURE CITED

CHARLTON, N. D., K. D. CRAVEN, S. MITTAL, A. A. HOPKINS, AND C. A. YOUNG. 2012. *Epichloë canadensis*, a new interspecific epichloid hybrid symbiotic with Canada wildrye (*Elymus canadensis*). *Mycologia* 104: 1187–1199.

CLAY, K. 1990. Fungal endophytes of grasses. *Annual Review of Ecology and Systematics* 21: 275–297.

CLAY, K., AND J. HOLAH. 1999. Fungal endophyte symbiosis and plant diversity in successional fields. *Science* 285: 1742–1744.

EXCOFFIER, L., G. LAVAL, AND S. SCHNEIDER. 2005. Arlequin ver. 3.0: An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1: 47–50.

FAETH, S. H. 2002. Are endophytic fungi defensive plant mutualists? *Oikos* 98: 25–36.

LEUCHTMANN, A., C. W. BACON, C. L. SCHARDL, J. F. WHITE, AND M. TADYCH. 2014. Nomenclatural realignment of *Neotyphodium* species with genus *Epichloë*. *Mycologia* 106: 202–215.

TABLE 2. Genetic properties of the newly developed SNP primers for six populations of *Epichloë canadensis*.^a

	BF (<i>tubB</i> SNP <i>n</i> = 6; <i>tefA</i> SNP <i>n</i> = 11)		CC (<i>tubB</i> SNP <i>n</i> = 17; <i>tefA</i> SNP <i>n</i> = 16)		HP (<i>tubB</i> SNP <i>n</i> = 9; <i>tefA</i> SNP <i>n</i> = 20)		OFL (<i>tubB</i> SNP <i>n</i> = 2; <i>tefA</i> SNP <i>n</i> = 2)		TLI (<i>tubB</i> SNP <i>n</i> = 10; <i>tefA</i> SNP <i>n</i> = 8)		VNWR (<i>tubB</i> SNP <i>n</i> = 24; <i>tefA</i> SNP <i>n</i> = 9)	
	<i>A</i>	<i>H</i>	<i>A</i>	<i>H</i>	<i>A</i>	<i>H</i>	<i>A</i>	<i>H</i>	<i>A</i>	<i>H</i>	<i>A</i>	<i>H</i>
<i>tubB</i> SNP	1	0.00	1	0.00	2	0.39	1	0.00	2	0.36	2	0.29
<i>tefA</i> SNP	3	0.58	2	0.50	2	0.39	2	1	2	0.25	3	0.42

Note: *A* = number of haplotypes found (total individuals sampled); *H* = gene diversity (probability of two randomly drawn haplotypes being different).
^a Full population names and locations are given in Appendix 1.

MOON, C. D., B. A. TAPPER, AND B. SCOTT. 1999. Identification of *Epichloë* endophytes in planta by a microsatellite-based PCR fingerprinting assay with automated analysis. *Applied and Environmental Microbiology* 65: 1268–1279.

MOON, C. D., K. D. CRAVEN, A. LEUCHTMANN, S. L. CLEMENT, AND C. L. SCHARDL. 2004. Prevalence of interspecific hybrids amongst asexual fungal endophytes of grasses. *Molecular Ecology* 13: 1455–1467.

NEI, M. 1987. Molecular evolutionary genetics. Columbia University Press, New York, New York, USA.

SHYMANOVICH, T., S. SAARI, M. E. LOVIN, A. K. JARMUSCH, S. A. JARMUSCH, A. M. MUSSO, N. D. CHARLTON, ET AL. 2014. Alkaloid variation among epichloid endophytes of sleepygrass (*Achnatherum robustum*) and consequences for resistance to insect herbivores. *Journal of Chemical Ecology* 41: 93–104.

SULLIVAN, T. J., AND S. H. FAETH. 2004. Gene flow in the endophyte *Neotyphodium* and implications for coevolution with *Festuca arizonica*. *Molecular Ecology* 13: 649–656.

TAKACH, J. E., AND C. A. YOUNG. 2014. Alkaloid genotype diversity of tall fescue endophytes. *Crop Science* 54: 667–678.

TAMURA, K., D. PETERSON, N. PETERSON, G. STECHER, M. NEI, AND S. KUMAR. 2011. MEGA5: Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28: 2731–2739.

UNTERGASSER, A., I. CUTCUTACHE, T. KORESSAAR, J. YE, B. C. FAIRCLOTH, M. REMM, AND S. G. ROZEN. 2012. Primer3—New capabilities and interfaces. *Nucleic Acids Research* 40: e115.

YE, J., G. COULOURIS, I. ZARETSKAYA, I. CUTCUTACHE, S. ROZEN, AND T. L. MADDEN. 2012. Primer-BLAST: A tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics* 13: 134.

YOUNG, C. A., N. D. CHARLTON, J. E. TAKACH, G. A. SWOBODA, M. A. TRAMMELL, D. V. HUHMANN, AND A. A. HOPKINS. 2014. Characterization of *Epichloë coenophiala* within the US: Are all tall fescue endophytes created equal? *Chemistry & Biology* 2: 95.

APPENDIX 1. Locations of sampled *Epichloë canadensis* populations.

Population	Abbreviation	Geographic coordinates
Boone Forks Wildlife Management Area, Iowa	BF	42°20'48.42"N, 93°50'39.78"W
Carleton College Cowling Arboretum, Minnesota	CC	44°28'4.79"N, 93°8'27.59"W
Hayden Prairie State Preserve, Iowa	HP	43°26'36.31"N, 92°23'0.04"W
Ottawa State Fishing Lake, Kansas	OFL	39°6'52.05"N, 97°34'16.85"W
Salina, Kansas	TLI	38°40'54.08"N, 97°35'28.11"W
Valentine National Wildlife Refuge, Nebraska	VNWR	42°31'51.45"N, 100°39'18.34"W

APPENDIX 2. *Epichloë* species^a identified by Primer-BLAST to produce amplicons with the described primers.

Species	<i>tubB</i>	<i>tefA</i>
<i>Epichloë australiense</i> (C. D. Moon & Schardl) Leuchtm.	AF323379.1	AF323400.1
<i>Epichloë baconii</i> White	KF042062.1	KF811547.1
<i>Epichloë cabralii</i> Iannone, M. S. Rossi & Schardl	JX679132.1	KJ934942.1
<i>Epichloë chisosae</i> (J. F. White & Morgan-Jones) Schardl	AF457470.1	AF457510.1
<i>Epichloë coenophiala</i> (Morgan-Jones & W. Gams) C. W. Bacon & Schardl	KF811599.1	KF811568.1
<i>Epichloë festucae</i> Leuchtm., Schardl & M. R. Siegel	KF042045.1	KF042045.1
<i>Epichloë festucae</i> var. <i>lolii</i> (Latch & Samuels) C. W. Bacon & Schardl	AY865628.1	AF457540.1
<i>Epichloë melicicola</i> (C. D. Moon & Schardl) Schardl	AF323387.1	AF323404.1
<i>Epichloë occultans</i> (C. D. Moon, B. Scott & M. J. Chr.) Schardl	AF176270.1	AF457541.1
<i>Epichloë siegelii</i> (K. D. Craven, Leuchtm. & Schardl) Leuchtm.	AF308139.1	AF308133.1
<i>Epichloë tembladerae</i> (Cabral & J. F. White) Iannone & Schardl	AF457496.1	AF457545.1

^aSpecies names follow classifications proposed in Leuchtmann et al. (2014).