

IDENTIFICATION OF VIRUSES INVOLVED IN THE 1971 OUTBREAK OF HEMORRHAGIC DISEASE IN SOUTHEASTERN UNITED STATES WHITE-TAILED DEER

Authors: THOMAS, F. C., WILLIS, N., and RUCKERBAUER, G.

Source: Journal of Wildlife Diseases, 10(3) : 187-189

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-10.3.187>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

IDENTIFICATION OF VIRUSES INVOLVED IN THE 1971 OUTBREAK OF HEMORRHAGIC DISEASE IN SOUTHEASTERN UNITED STATES WHITE-TAILED DEER

F. C. THOMAS, N. WILLIS and G. RUCKERBAUER¹

Abstract: Tissues and serum from white-tailed deer (*Odocoileus virginianus*) in the southeastern United States were taken during an outbreak of hemorrhagic disease. These were subjected to virus isolation attempts in tissue culture and several serological tests (plaque neutralization, complement fixation, agar gel precipitation). Both epizootic hemorrhagic disease of deer virus (EHDV) and bluetongue virus (BTV) were isolated and evidence of their presence demonstrated serologically. All of the evidence supported previous suggestions that these two agents are antigenically distinguishable.

INTRODUCTION

During the summer of 1971, a widespread disease of white-tailed deer was observed in the southeastern United States.² Samples (tissue and serums) were supplied to the Animal Diseases Research Institute¹ by the Southeastern Cooperative Wildlife Disease Study² (SCWDS) and the U.S. Bureau of Sport Fisheries and Wildlife³ (BSFW). These included samples from wild deer (SCWDS) and both native and newly introduced penned deer (BSFW). The latter were sentinel deer which were serologically negative for bluetongue virus (BTV) and epizootic hemorrhagic disease virus (EHDV). The clinical, pathological and epidemiological observations will be reported by other investigators.^{2,4}

MATERIAL AND METHODS

Five sets of blood samples and other tissues (from deer no. 1 to 5) of which

three (No. 1, 2, 3) were accompanied by serum, were supplied by SCWDS. All five deer were dead or affected by the described syndrome.^{2,4} Blood samples (for virus isolation) were preserved with equal volumes of OPG.⁵ Serum and other tissues were shipped frozen.

Serum samples were received from BSFW and included 14 from penned deer (all normal or convalescing) in the Mammoth Cave, Kentucky area.

Blood and pooled tissue suspensions (usually spleen, liver, heart, lung and kidney) submitted for virus isolation were inoculated onto L-cells⁵ and WI2⁵ cells. The isolates were identified by neutralization,⁵ plaque type,⁶ and fluorescent antibody staining.¹ In one case (tissue pool 3) two viruses were separated by the conditions of their replication in WI2 cells, without the use of antiserum. They were identified by a fluorescent focus technique employing a gum tragacanth overlay⁸ and specific conjugates.¹ Serum samples were tested with one or more of

¹ Animal Pathology Division, Health of Animals Branch, Agriculture Canada, Animal Diseases Research Institute, P.O. Box 1400, Hull, Quebec, Canada.

² Southeastern Cooperative Wildlife Disease Study, College of Veterinary Medicine, University of Georgia, Athens, Georgia 30601, U.S.A.

³ U.S. Bureau of Sport Fisheries and Wildlife, Mammoth Cave, Kentucky, and Denver Wildlife Research Center, Denver, Colorado.

⁴ Roughton, R. 1974. An outbreak of a hemorrhagic disease of white-tailed deer in Kentucky. Manuscript in preparation.

⁵ O.P.G. — potassium oxalate, 5 ml; phenol, 5 gm; glycerol, 500 ml; Distilled water, 500 ml.

the following tests for EHDV and BTV antibodies; plaque neutralization^a (PN), complement fixation¹ (CF) and agar gel precipitation¹ (AGP).

Serum 1 was EHDV antibody positive, serum 2 BTV antibody positive and serum 3 negative (Table 1).

RESULTS

SCWDS Samples:

BTV was isolated from blood 2 and EHDV from blood 3, the others being negative. Tissue pool 1 yielded no virus, pools 2 and 4 BTV, pool 3 both BTV and EHDV and pool 5 EHDV. These isolations were repeatable.

BSFW Samples:

Of nine native deer serums from this group, one was positive for BTV and EHDV, two questionable for BTV and positive for EHDV and six negative for both viruses. The five samples from the sentinel (introduced) deer yielded one questionable for EHDV, three questionable for BTV and positive for EHDV and one positive for EHDV (Table 2).

TABLE 1. Virus isolation and serological results from samples supplied by Southeastern Cooperative Wildlife Disease Study.

| Specimen | Virus Isolation (blood) | | Virus Isolation (tissues) | | Serology | |
|----------|-------------------------|------|---------------------------|------|-----------------|--------------|
| | BTV | EHDV | BTV | EHDV | BTV | EHDV |
| 1 | — | — | — | — | — ^a | +(CF,AGP,PN) |
| 2 | + | — | + | — | +(AGP,PN) | — |
| 3 | — | + | + | + | — | — |
| 4 | — | — | + | — | ND ^b | ND |
| 5 | — | — | — | + | ND | ND |

^a negative in all 3 tests (CF, AGP, PN).

^b ND — not done

TABLE 2. Serological results on samples supplied by the U.S. Bureau of Sport Fisheries and Wildlife.

| Resident Penned Deer | Serology | |
|-----------------------|---------------------|----------------|
| | BTV | EHDV |
| MC-109 | +(PN) | +(CF, AGP, PN) |
| MC-125 | Q ^a (PN) | +(PN) |
| MC-141 | Q(PN) | +(CF, AGP, PN) |
| Remainder (6 samples) | — ^b | — |
| Sentinel Deer | | |
| MC-236 | — | Q(PN) |
| MC-237 | Q(PN) | +(PN) |
| MC-238 | — | +(CF, AGP, PN) |
| MC-239 | Q(PN) | +(CF, AGP, PN) |
| MC-240 | Q(PN) | +(CF, AGP, PN) |

^a questionable.

^b negative in all 3 tests (CF, AGP, PN).

These results and an indication of the tests on which our conclusions were based are given in Tables 1 and 2.

DISCUSSION

The data indicate that both BTV and EHDV were active during the 1971 deer die-off. Inasmuch as both viruses have been experimentally shown to be capable of producing hemorrhagic lesions and

mortality in white-tailed deer,^{3,7} the laboratory findings suggest these to be the most likely candidates.

It is interesting that both virus isolations and serology confirmed previous suggestions that the viruses are antigenically distinguishable.^{5,6} One could hypothesize that climatic conditions (i.e. those favouring biting anthropods) allowed BTV and EHDV already present in the area to infect large numbers of animals, some of which died.

Acknowledgements

The authors wish to acknowledge the collaboration of Drs. A. K. Prestwood and F. A. Hayes of the Southeastern Cooperative Wildlife Disease Study and Drs. R. Roughton and M. Friend of the U.S. Bureau of Sport Fisheries and Wildlife.

LITERATURE CITED

1. BOULANGER, P., ANDRE GIRARD, G. L. BANNISTER and G. M. RUCKERBAUER. 1970. Serological studies in bluetongue. Proc. 74th Ann. Meeting U.S. Animal Health Assoc.: 537-548.
2. PRESTWOOD, A. K., T. P. KISTNER, F. E. KELLOGG and F. A. HAYES. 1974. The 1971 outbreak of hemorrhagic disease among white-tailed deer of the southeastern United States. J. Wildl. Dis. 10: 217-224.
3. THOMAS, F. C. and D. O. TRAINER. 1970. Bluetongue virus in white-tailed deer. Am. J. vet. Res. 31: 271-278.
4. THOMAS, F. C. and D. O. TRAINER. 1970. Bluetongue virus: (1) In pregnant white-tailed deer. (2) A plaque reduction neutralization test. J. Wildl. Dis. 6: 384-389.
5. THOMAS, F. C. and JANICE MILLER. 1971. A comparison of bluetongue virus and EHD virus: Electron microscopy and serology. Can. J. comp. Med. 35: 22-28.
6. THOMAS, F. C. and D. O. TRAINER. 1971. Bluetongue virus: Some relationship among North American isolates and further comparisons with EHD virus. Can. J. comp. Med. 35: 187-191.
7. WILHELM, A. J. and D. O. TRAINER. 1969. Hematological and virological studies of epizootic hemorrhagic disease of deer. Bull. Wildl. Dis. Assoc. 5: 77-94.
8. WILLIS, N. G., P. N. COREY, J. B. CAMPBELL and L. W. MACPHERSON. 1970. Development and statistical evaluation of a plaque assay for the virus of epizootic hemorrhagic disease of deer. Can. J. Microbiol. 16: 1293-1301.

Received for publication 27 August 1973