



PARASITISM AMONG WHITE-TAILED DEER AND DOMESTIC SHEEP ON COMMON RANGE 1

Authors: PRESTWOOD, ANNIE K., PURSGLOVE, SAMUEL R., and HAYES, FRANK A.

Source: Journal of Wildlife Diseases, 12(3) : 380-385

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-12.3.380>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

PARASITISM AMONG WHITE-TAILED DEER AND DOMESTIC SHEEP ON COMMON RANGE[†]

ANNIE K. PRESTWOOD, SAMUEL R. PURSGLOVE and FRANK A. HAYES,
Southeastern Cooperative Wildlife Disease Study, Department of Parasitology,
College of Veterinary Medicine, University of Georgia, Athens, Georgia 30602, USA

Abstract: Parasitism was studied in white-tailed deer (*Odocoileus virginianus*) and domestic sheep (*Ovis aries*) which shared a common range in eastern West Virginia. Of 30 species of internal parasites, 11 were found in deer and 22 in sheep. Five parasites, *Sarcocystis* sp., *Cysticercus tenuicollis*, *Oesophagostomum venulosum*, *Cooperia punctata*, and *Gongylonema pulchrum*, occurred in both deer and sheep. An index of similarity of 17.2 suggests that the parasite faunas of these hosts are distinct, and that it is unlikely that white-tailed deer are reservoirs of common parasites of domestic sheep in the southern Appalachian region.

INTRODUCTION

According to a recent review, 79 genera containing 137 species of parasites have been reported from North American deer of the genus *Odocoileus*.¹⁷ Of these, approximately 40% also have been reported from domestic livestock. Recent findings¹⁰ have suggested that the helminth faunas of intermingling populations of white-tailed deer and cattle (*Bos taurus*) are distinct with little exchange occurring. These authors concluded that white-tailed deer are not important reservoirs of helminths of cattle. The parasite faunas of white-tailed deer and domestic sheep utilizing the same range, however, have not been described. This report presents findings on parasitism among intermingling populations of white-tailed deer and domestic sheep.

MATERIALS AND METHODS

Animals from Hardy County, West Virginia, were selected for study since white-tailed deer are abundant and sheep production is of major agricultural importance. This county is located in the

mountainous eastern panhandle of West Virginia. Approximately 70% of the area is forested and lies 300 m or more above sea level.

The specific collection site for deer and sheep met the following criteria: 1. deer and sheep commonly were observed in the same pastures; 2. an abundance of deer was present; 3. access to area was relatively easy facilitating collection of deer; and 4. availability of adult sheep and lambs which had not received anthelmintic medication during the previous 6 months.

The collection site was typical of many small, family farms scattered throughout the Appalachian region of the Southeast. The farm was situated in a shallow, rocky valley having relatively poor soils. Forty-eight hectares of unimproved pasture of native grasses occurred in the bottoms and extended onto the hillsides, where a flock of 47 ewes and 70 lambs grazed. The ridges were heavily forested with an oak-hickory vegetative type. Supplemental grain was provided only for ewes during pregnancy. Hay was given sheep on pastures during the winter. An

[†] Supported by an appropriation from the Congress of the United States. Funds were administered and research coordinated by the Federal Aid in Wildlife Restoration Act (50 Stat. 917) and through Contract No. 14-16-0008-638, Fish and Wildlife Service, U.S. Department of the Interior.

anthelmintic was administered to sheep during the previous spring. Unusual disease problems had not occurred.

During September, 1972, 5 (4 adults and 1 fawn) deer were collected at night by shooting. Deer were wrapped in plastic, surrounded by crushed ice and transported to a field laboratory for necropsy the following day. Concomitantly, 5 (3 adults and 2 lambs) sheep were purchased from the same farm where white-tailed deer were obtained. Sheep were necropsied immediately after euthanasia.

Postmortem procedures were similar for deer and sheep. After searching the skin and pelage for ectoparasites, the animals were skinned before further examination. General necropsy and parasitologic techniques have been presented elsewhere.⁹⁻¹⁵ Fecal flotations using saturated sodium nitrate solution were done to detect coccidian oocysts. Sections of tongue, lung, liver, kidney, spleen, lymph node, cardiac muscle, and abomasum

were preserved in buffered 10% formalin. Tissues were processed according to standard procedures, cut at 6 μ m, and stained with Delafield's hematoxylin and eosin. An index of similarity⁶ was used for comparing parasite faunas, and parasite profiles¹⁰ were prepared for each host.

RESULTS

No ectoparasites were found on deer or sheep, but 30 species of internal parasites (Table 1) representing 18 genera were identified, viz., 2 protozoans, 2 cestodes, and 26 nematodes. White-tailed deer were parasitized by 11 species while 24 species were found in sheep. Five parasites, *Sarcocystis* sp., *Cysticercus tenuicollis*, *Oesophagostomum venulosum*, *Cooperia punctata*, and *Gongylonema pulchrum*, occurred in both deer and sheep. An index of similarity of 17.2 was calculated for the parasite faunas of these animals.

TABLE 1. Parasites of white-tailed deer and domestic sheep of West Virginia.

Parasite	Host
PROTOZOA	
<i>Eimeria</i> sp.	Sheep
<i>Sarcocystis</i> sp.	Deer, sheep
CESTODA	
<i>Cysticercus tenuicollis</i>	Deer, sheep
<i>Moniezia expansa</i>	Sheep
NEMATODA	
Threadworms	
<i>Capillaria bovis</i>	Deer
Whipworms	
<i>Trichuris ovis</i>	Sheep
<i>Trichuris skrjabini</i>	Sheep
Nodular worms	
<i>Oesophagostomum columbianum</i>	Sheep
<i>Oesophagostomum venulosum</i>	Deer, sheep
Bowel worms	
<i>Chabertia ovina</i>	Sheep

TABLE 1 (Continued)

Parasite	Host
NEMATODA (Continued)	
Trichostrongyles	
<i>Cooperia curticei</i>	Sheep
<i>Cooperia mcmasteri</i>	Sheep
<i>Cooperia oncophora</i>	Sheep
<i>Cooperia punctata</i>	Deer, sheep
<i>Haemonchus contortus</i>	Sheep
<i>Ostertagia circumcincta</i>	Sheep
<i>Ostertagia dikmansi</i>	Deer
<i>Ostertagia mossi</i>	Deer
<i>Ostertagia trifurcata</i>	Sheep
<i>Skrjabinagia odocoilei</i>	Deer
<i>Trichostrongylus axei</i>	Sheep
<i>Trichostrongylus colubriformis</i>	Sheep
<i>Trichostrongylus vitrinus</i>	Sheep
<i>Nematodirus fillicollis</i>	Sheep
<i>Nematodirus spathiger</i>	Sheep
Lungworms	
<i>Dictyocaulus filaria</i>	Sheep
<i>Dictyocaulus viviparus</i>	Deer
<i>Muellerius capillaris</i>	Sheep
<i>Parelaphostrongylus tenuis</i>	Deer
Spiruroids	
<i>Gongylonema pulchrum</i>	Deer, sheep

Parasite burdens of deer ranged from 243 to 4963 (av. 1475) helminths. Numbers of worms in sheep ranged from 1604 to 37,767 (av. 13,003). Parasite profiles for each host are presented in Figures 1 and 2.

Microscopic examination of tissues revealed *Sarcocystis* in the tongues of 2 deer and in tongue and cardiac muscle of 4 sheep. One to 4 sarcocysts were seen in histological sections. Pathologic alterations associated with eggs and larvae of *Parelaphostrongylus* were present in the lungs of 4 deer. Most lesions were mild and consisted of a granulomatous reac-

tion with eosinophils, lymphocytes, histiocytes, and occasional foreign body giant cells. A cysticercus was found in the lungs of 1 deer; adjacent alveoli were compressed but inflammatory response was minimal. The lungs of 4 sheep contained lesions attributable to lungworm infection. Histologically, there were multifocal areas of granulomatous inflammation, peribronchial lymphoid hyperplasia, smooth muscle hypertrophy, and in 1 animal, multifocal areas of chronic bronchopneumonia. Reaction was mild to severe, and 1 animal was considered to have verminous pneumonia.




FIGURE 1. Parasite profile for white-tailed deer in eastern West Virginia.




FIGURE 2. Parasite profile for domestic sheep in eastern West Virginia.

DISCUSSION

Much of the sheep production in the southeastern United States traditionally has been located in mountainous terrain where farming operations are primarily confined to valleys and gentle slopes. Steeper slopes and ridges in this region usually are forested, and the edge effect provided by the meeting of two distinct types is conducive to production of an abundance of wildlife including white-tailed deer as evidenced by legal harvest of over 2000 deer in this county during 1972. Due to the distribution of farming operations along the slopes and valleys, frequent contact between deer and sheep is afforded. Of the 11 parasites found in the deer examined, 5 (45.4%) also were found in sheep. These findings are similar to those reported previously;¹⁷ however, the low index of similarity (17.2) strongly suggests that the parasite faunas of deer and sheep are quite distinct. Only on one occasion was a "shared" parasite, viz., *Gongylonema pulchrum*, more prevalent in white-tailed deer. Based on

this study, white-tailed deer cannot be considered important reservoirs for common parasites of the lung and gastrointestinal tract of sheep in the southern Appalachians.

Although experimentally infected sheep² were considered relatively resistant to neurologic disease caused by *Parelaphostrongylus tenuis*, neurologic disorders caused by meningeal worm have been reported in sheep of New York,¹⁸ Connecticut,⁷ and West Virginia.¹ White-tailed deer are usual hosts for this helminth, where only occasionally does it produce neurologic signs.^{3,8} The prevalence of meningeal worm in deer of this study was comparable to that previously reported in eastern West Virginia.¹³ Neurologic disease, however, had not occurred in sheep on this or adjacent farms at the time of this study. In view of this and of results of experimental infections,² it is likely that unusual ecologic conditions

which promote contact between sheep and infected intermediate hosts are necessary to precipitate sporadic outbreaks of neurologic disease among sheep.

The intensity of infection with helminths among white-tailed deer was considered low to moderate when compared with deer from other areas of the Southeast. Lungworms (*Dictyoaulus viviparus* and *P. tenuis*) potentially were the most pathogenic parasites present in deer. Microscopic lung lesions were similar to those described previously for *P. tenuis*⁴ and for *Parelaphostrongylus andersoni*.⁵

Among sheep, only 1 animal was considered heavily parasitized, and lungworm pneumonia caused by *D. filaria* and *Miellerius capillaris* was diagnosed histologically. This animal also harbored the greatest number of gastrointestinal helminths. The sum effect of all helminths undoubtedly contributed to the overall poor physical condition of this adult ewe.

Acknowledgments

The assistance of Mr. Gary D. Strawn and other members of the West Virginia Department of Natural Resources is gratefully acknowledged.

LITERATURE CITED

1. ALDEN, C., F. WOODSON, R. MOHAN and S. MILLER. 1975. Cerebrospinal nematodiasis in sheep. J. Am. vet. med. Ass. 166: 784-786.
2. ANDERSON, R. C. and UTA R. STRELIVE. 1966. Experimental cerebrospinal nematodiasis (*Pneumostrongylus tenuis*) in sheep. Can. J. Zool. 44: 889-894.
3. ECKROADE, R. J., G. M. ZU RHEIN and W. FOREYT. 1970. Meningeal worm invasion of the brain of a naturally infected white-tailed deer. J. Wildl. Dis. 6: 430-436.
4. GOBLE, F. C. 1941. Tissue changes in white-tailed deer (*Odocoileus virginianus borealis*) accompanying natural infections of lungworms (Genera *Protostrongylus* and *Dictyoaulus*). J. Wildl. Manage. 5: 141-158.
5. HOLMES, J. C. and R. PODESTA. 1968. The helminths of wolves and coyotes from the forested regions of Alberta. Can. J. Zool. 46: 1193-1204.
6. NETTLES, V. F. and ANNIE K. PRESTWOOD. 1976. Experimental *Parelaphostrongylus andersoni* infections in white-tailed deer. Vet. Path. In Press.
7. NIELSEN, S. W. and J. AFTOSMIS. 1964. Spinal nematodiasis in two sheep. J. Am. vet. med. Ass. 144: 155-158.
8. PRESTWOOD, ANNIE K. 1970. Neurologic disease in a white-tailed deer massively infected with meningeal worm (*Pneumostrongylus tenuis*). J. Wildl. Dis. 6: 84-86.

9. ———, F. A. HAYES, J. H. EVE and J. F. SMITH. 1973. Abomasal parasites of white-tailed deer in southeastern United States, Texas, and the Virgin Islands. J. Am. vet. med. Ass. 163: 556-561.
10. ———, F. E. KELLOGG, S. R. PURSGLOVE and F. A. HAYES. 1975. Helminth parasitism among intermingling insular populations of white-tailed deer, feral cattle, and feral swine. J. Am. vet. med. Ass. 166: 787-789.
11. ———, V. F. NETTLES and F. E. KELLOGG. 1974. Distribution of muscle-worm, *Parelaphostrongylus andersoni*, among white-tailed deer of the southeastern United States. J. Wildl. Dis. 10: 404-409.
12. ——— and S. R. PURSGLOVE. 1974. *Leptostrongylus alpenae* in white-tailed deer (*Odocoileus virginianus*) of Georgia and Louisiana. J. Parasit. 60: 573.
13. ——— and J. F. SMITH. 1969. Distribution of meningeal worm (*Pneumostrongylus tenuis*) in deer in the southeastern United States. J. Parasit. 55: 720-725.
14. ———, J. F. SMITH and J. BROWN. 1971. Lungworms in white-tailed deer of the southeastern United States. J. Wildl. Dis. 7: 149-154.
15. ———, ——— and W. E. MAHAN. 1970. Geographic distribution of *Gongylonema pulchrum*, *Gongylonema verrucosum* and *Paramphistomum liorchis* in white-tailed deer of the southeastern United States. J. Parasit. 56: 123-127.
16. UHAZY, L. S. and J. C. HOLMES. 1971. Helminths of the Rocky Mountain bighorn sheep in western Canada. Can. J. Zool. 49: 507-512.
17. WALKER, MARTHA L. and W. W. BECKLUND. 1970. Checklist of the internal and external parasites of deer, *Odocoileus hemionus* and *O. virginianus*, in the United States and Canada. Special Publication No. 1, Index-Catalogue of Medical and Veterinary Zoology, U.S. Gov. Printing Off., Washington, D.C. 1-45.
18. WHITLOCK, J. H. 1952. Neurofilariosis, a paralytic disease of sheep. Cornell Vet. 42: 125-132.

Received for publication 19 January 1976