



BLOOD PARASITES OF SOME COLUMBIFORM AND PASSERIFORM BIRDS FROM CHILE 1

Author: FORRESTER, DONALD J.

Source: Journal of Wildlife Diseases, 13(1) : 94-96

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-13.1.94>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

BLOOD PARASITES OF SOME COLUMBIFORM AND PASSERIFORM BIRDS FROM CHILE[□]

DONALD J. FORRESTER, Laboratory of Wildlife Disease Research,
College of Veterinary Medicine, University of Florida, Gainesville, Florida 32610, USA

ELLIS C. GREINER, Department of Biology and International Reference Centre for
Avian Haematozoa, Memorial University of Newfoundland,
St. John's, Newfoundland, Canada A1C 5S7

ROBERT W. McFARLANE, Savannah River Ecology Laboratory, Aiken, South Carolina 29801, USA

Abstract: Ninety-one birds of 23 species from Chile were examined for haematozoa; 13 birds of seven species harbored species of *Haemoproteus*, *Leucocytozoon*, *Plasmodium*, *Trypanosoma* and microfilariae. Haemoproteids (representing four species) were the most common parasites and occurred in 10 of the 13 infected birds.

INTRODUCTION

Virtually nothing is known about avian blood parasites from Chile. Lucena,⁶ in his checklist of neotropical avian haematozoa, cites two reports: *Haemoproteus danilewskyi* from *Sturnella defilippii*⁷ and *Leucocytozoon* sp. from *Sycalis arvensis*.⁸ However, both findings involve captive Chilean birds in the Zoological Gardens in London, England, and may not be valid records for Chile.

During 1973, one of us (R.W.M.) had an opportunity to obtain blood films from a small sample of columbiform and passeriform birds from Chile. The results of analyses of these slides are presented.

STUDY AREAS AND METHODS

The Department of Arica, Province of Tarapacá, is the northernmost political subdivision of Chile, bordering both Peru and Bolivia. Although it lies within the tropics at 18° S latitude, its climate is basically arid and temperate. Rainfall is virtually unknown at low elevations and seldom exceeds 25 cm in the Andean highlands. The narrow east-west valleys

which traverse the Atacama Desert provide meager vegetation and birdlife except where agricultural irrigation contributes to greater habitat diversity. Above the desert lie narrow north-south altitudinal zones of vegetation which ascend from columnar cacti to xerophytic scrub (tola) and finally tussock grassland (puna). The occurrence and dispersal of many avian species are thus restricted to certain elevations.

Birds were collected with mist nets from March to June, 1973, from three agricultural valleys and two higher localities: Molinos (930 m elevation) and Chapisca (1010 m) in Lluta Valley, Azapa (250 m) in Azapa Valley, Taltape (780 m) in Camarones Valley, the agricultural community of Putre (3500 m) in the tolar zone, and Chilcaya (4250 m) in the puna zone. Blood films were air-dried in the field, subsequently fixed in 100% methanol and stained with Giemsa's Stain. A minimum of 20,000 erythrocytes was examined on each slide. All positive smears have been deposited in the collection of the International Reference Centre for Avian Haematozoa, Memorial University of Newfoundland, St. John's, Newfoundland, Canada.

[□] Florida Agricultural Experiment Stations Journal Series No. 6208.

RESULTS AND DISCUSSION

Blood films from 91 birds of 23 species representing eight families were examined (Table 1). Thirteen birds (14%) harbored one or more parasites; five of these had multiple infections. The most commonly occurring parasite genus was *Haemoproteus* which was represented by four species: *H. columbae* in two of two *Zenaida asiatica* and one of five *Columbina cruziana*, *H. sacharovi* in one of two *Z. asiatica*, *H. orizivora* in all in-

fectured *Phrygilus fruticeti* and *Zonotrichia capensis* and *H. fringillae* in one of four *P. fruticeti* and all three infected *Z. capensis*. *Leucocytozoon fringillinarum* and *L. majoris* occurred concurrently in *P. fruticeti*. *Plasmodium relictum* was found in *Passer domesticus*. *Trypanosoma* sp. occurred in *P. fruticeti*, but because only one specimen was found in one bird it was not identified to species. Microfilariae were not further identified.

TABLE 1. Blood parasites of some columbiform and passeriform birds from Chile.

Bird Species	Total number		H*	L*	P*	T*	M*
	Examined	Infected					
COLUMBIDAE							
<i>Columbina cruziana</i>	5	1	1				
<i>Zenaida asiatica</i>	2	2	2				
COEREBIDAE							
<i>Conirostrum cinereum</i>	4	1					1
FRINGILLIDAE							
<i>Phrygilus fruticeti</i>	4	4	4	1		1	2
<i>Zonotrichia capensis</i>	15	3	3				
PLOCEIDAE							
<i>Passer domesticus</i>	5	1			1		
TYRANNIDAE							
<i>Elaenia albiceps</i>	2	1					1
Uninfected species (see below)	54						
Totals	91	13 (14%)	10	1	1	1	4

*H = *Haemoproteus*; L = *Leucocytozoon*; P = *Plasmodium*; T = *Trypanosoma*; M = microfilariae.

Uninfected Species (number examined in parentheses): Columbidae—*Metriopelia ceciliae* (1); *Metriopelia melanoptera* (3); Coerebidae—*Conirostrum tamarugensis* (1); Fringillidae—*Catamenia analis* (2); *Phrygilus plebejus* (4); *Sicalis uropygialis* (6); *Spinus magellanicus* (13); *Sporophila telasco* (6); *Xenospingus concolor* (4); Furnariidae—*Asthenes dorbignyi* (1); *Asthenes modesta* (1); *Cinclodes fuscus* (2); *Upucerthia validirostris* (1); Thraupidae—*Thraupis bonariensis* (3); Turdidae—*Turdus chiguanco* (2); Tyrannidae—*Anairetes flavirostris* (4).

Haemoproteid infections occurred in all collection sites except Chapisca and Chilcaya. This would indicate that elevation may not be a factor in haemoproteid vector distribution in Northern Chile. The single bird infected with *Leucocytozoon* was collected at 3500 m whereas the only occurrence of *Plasmodium* was at 250 m.

The low overall prevalence of haematozoan infection observed in this sample of birds from Chile is in agreement with results obtained recently in Columbia,¹ Venezuela^{2,3} and Brasil.^{4,5} Reasons for the low prevalence are unknown but, as suggested by Bennett and Borrero,¹ may be due to a lack of suitable vector species.

Acknowledgments

The authors are grateful for the advice of Drs. G. F. Bennett and C. M. Herman and for the technical assistance of P. P. Humphrey. The Centro de Investigación y Capacitación Agrícola of the Univesidad del Norte graciously provided transportation and assistance in specimen collection. Part of this study was conducted while the senior author was a Visiting Scientist at the International Reference Centre for Avian Haematozoa, Memorial University of Newfoundland, St. John's, Newfoundland.

LITERATURE CITED

1. BENNETT, G. F. and J. I. BORRERO H. 1976. Blood parasites of some birds from Columbia. *J. Wildl. Dis.* 12: 454-458.
2. GABALDON, A., G. ULLOA and A. G. de MONTCOURT. 1974. Encuesta sobre malaria aviaria en Venezuela: Resultados del primer año. *Bol. Dir. Malar. San. Amb.* 19: 80-103.
3. ———. 1975. Encuesta sobre malaria aviaria en Venezuela: Resultados del segundo año. *Bol. Dir. Malar. San. Amb.* 20: 73-92.
4. LAINSON, R., J. J. SHAW and P. S. HUMPHREY. 1970. Preliminary survey of blood-parasites of birds of the area de pesquisas ecológicas do Guamá, Belém, Pará, Brasil. *J. Parasit.* 56 (Sect. II, Part I): 197-198.
5. LOPES, O. de S. and G. F. BENNETT. Blood parasites of some birds from Sao Paulo State, Brasil. Unpublished manuscript in files of International Reference Centre for Avian Haematozoa, Memorial University of Newfoundland, St. John's, Newfoundland.
6. LUCENA, D. T. 1941. Lista dos protozoários hemoparasitas de aves da região neotrópica. *Rev. Fac. Med. Vet. S. Paulo.* 2: 33-66.
7. PLIMMER, H. G. 1913. Report on the deaths which occurred in the Zoological Gardens during 1912, together with the blood-parasites found during the year. *Proc. Zool. Soc. London:* 141-149.
8. SCOTT, H. H. 1926. Report on the deaths occurring in the Society's gardens during the year 1925. *Proc. Zool. Soc. London:* 231-244.

Received for publication 16 August 1976