

THE PREVALENCE OF ANTI-LEPTOSPIRAL AGGLUTININS IN SERA OF WILDLIFE IN SOUTHEASTERN AUSTRALIA

Authors: MILNER, A.R., and WILKS, C.R.

Source: Journal of Wildlife Diseases, 17(2) : 197-202

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-17.2.197>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

THE PREVALENCE OF ANTI-LEPTOSPIRAL AGGLUTININS IN SERA OF WILDLIFE IN SOUTHEASTERN AUSTRALIA

A.R. MILNER[□] and C.R. WILKS, Veterinary Research Institute, Department of Agriculture, Park Drive, Parkville, Victoria.

D.M. SPRATT, Division of Wildlife Research, CSIRO, P.O. Box 84, Lyneham, A.C.T.

P.J.A. PRESIDENTE,[□] Regional Veterinary Laboratory, Department of Agriculture, Bairnsdale, Victoria, Australia.

Abstract: Anti-leptospiral agglutinins were found in the serum from 18 (7 species) of 419 (25 species) animals sampled from various areas of southeastern Australia. Positive serologic reactions were observed in 5 of 25 (20%) brush-tailed possum (*Trichosurus vulpecula*), 1 of 26 (3.8%) tammar wallaby (*Macropus eugenii*), 2 of 12 (16.7%) swamp wallaby (*Wallabia bicolor*), 1 of 3 (33.3%) koala (*Phascogale cinereus*), 3 of 41 (7.3%) common wombat (*Vombatus ursinus*), 2 of 100 (2%) bush rat (*Rattus fuscipes*) and 4 of 12 (25%) rusa deer (*Cervus timorensis*). The majority (55.5%) of serologic reactions were to serovar *hardjo*.

No serologic reactions were observed in samples from echidna (*Tachyglossus aculeatus*), brown antechinus (*Antechinus stuartii*), swainson's antechinus (*Antechinus swainsonii*), long-nosed bandicoot (*Perameles nasuta*), brown bandicoot (*Isodon obesulus*), common ringtail (*Pseudocheirus peregrinus*), greater glider (*Scolinobates volans*), eastern grey kangaroo (*Macropus giganteus*), red-necked wallaby (*Macropus rufogriseus*), rabbit (*Oryctolagus cuniculus*), water rat (*Hydromys chrysogaster*), black rat (*Rattus rattus*), eastern swamp rat (*Rattus lutreolus*), broad-toothed rat (*Mastacomys fuscus*), fox (*Vulpes vulpes*), sambar deer (*Cervus unicolor*), hog deer (*Axis porcinus*) and fallow deer (*Dama dama*).

INTRODUCTION

The most extensive surveys of the prevalence of infection with *Leptospira interrogans* in Australian wildlife have been conducted in the northern, tropical areas of Australia. In northern Queensland it was concluded from serologic and bacteriologic evidence that numerous wildlife species including *Rattus scordius conatus*, *Rattus rattus*, *Rattus norvegicus*, *Rattus assimilis*, *Perameles nasuta*, *Isodon macrourus* and *Mus musculus* were maintenance hosts for various leptospiral serovars.^{1,5} In a more recent survey in this area, 6.1%

of the rodents sampled had evidence of infection and, although the overall prevalence of rodent leptospirosis was low, highly infected foci were observed in wet environments.⁷

In a more temperate area of Australia, Munday¹⁰ surveyed 26 species of Tasmanian fauna for antibodies to a number of serovars but obtained few positive reactions. He considered that wombats (*Vombatus ursinus*) may be susceptible to infection with *pomona* and that *Rattus norvegicus* may be a maintenance host for *icterohaemorrhagiae*. Durfee and Presidente³ in a serologic survey of Vic-

[□] Present address: John Curtin School of Medical Research, Australian National University, Canberra, Australia.

[□] Present address: "Attwood" Laboratory, Department of Agriculture, Mickleham Road, Melbourne, Australia.

torian wildlife for antibodies to leptospire of the Hebdomadis serogroup, found a high prevalence of reactors in brush-tailed possums (*Trichosurus vulpecula*), wombats and rusa deer (*Cervus timorensis*).

A number of reports indicate that Australian wildlife may be of importance in the transmission of leptospirosis to both humans and domestic animals. Sullivan¹³ noted that Weil's disease in humans is acquired from rats, as are most cases of "canefields leptospirosis" caused by serovars *australis* and *zanoni* in Queensland. Battey *et al.*¹ noted that *Rattus sordidus conatus* was responsible for many human infections with *australis* and *grippotyphosa* in northern Queensland. Both the wombat and the feral pig may be involved in the transmission of *pomona* to domestic animals.^{8,10,11}

Because of the potential importance of wildlife in the transmission of leptospirosis to man and domestic animals a survey was performed using 12 leptospiral serovars to determine the prevalence of antibodies in wildlife sera collected in New South Wales and Victoria.

MATERIALS AND METHODS

Sampling Areas. Samples were taken from a number of areas throughout southeastern Australia (Fig. 1). Most locations were forested with only limited possibilities for contact with domestic animals.

Serology. Serologic testing was performed by the microscopic agglutination test (MAT) as reported by Galton *et al.*¹⁶ and modified by Cole *et al.*² Each serum was tested against members of the following serogroups: Australis (serovar *australis*), Autumnalis (*autumnalis*), Ballum (*ballum*), Bataviae (*bataviae*), Canicola (*canicola*), Grippotyphosa (*grippotyphosa*), Hebdomadis (*hardjo*), Icterohaemorrhagiae (*icterohaemorrhagiae* and *copenhageni*), Pomona (*pomona*), Pyrogenes (*zanoni*) and Tarassovi (*tarassovi*).

Each serum was screened at a final dilution of 1:40 against each antigen. All positive reactors were then titrated to an end point. Titers less than 1:32 were regarded as negative.

Animals. Serum samples were obtained from 419 animals (25 species, 12 families) (Table 1). No attempt was made to isolate leptospire from these samples. All samples from *Macropus eugenii*, *Cervus timorensis* and *Dama dama* were from captive animals. All other animals sampled were free ranging.

RESULTS

Anti-leptospiral agglutinins were detected in the serum of 18 animals from 7 species (Table 1). The prevalence of antibodies was highest in rusa deer, brush-tailed possums and swamp wallabies, although 1 of the 3 koala samples was positive.

The serum-antibody titers were low with the exception of the titers of 1:1024 to *hardjo* in one brush-tailed possum and one rusa deer.

The majority (55.5%) of serologic reactions obtained were to *hardjo*.

DISCUSSION

The results of the present survey indicate that 18 of 419 (1.9%) animals examined had evidence of past or present infection with leptospire. It must be emphasised, however, that although representatives of 11 pathogenic serogroups were used as antigens in the present study, no indication of the prevalence of infection with other, untested, leptospire can be gauged because of the serogroup specific nature of the MAT.

The lack of evidence of widespread infection in the rodents examined in this study was surprising. Although it has

been demonstrated that some individual rodents may excrete leptospires without having detectable circulating antibodies^{7,12} it would appear that in southeastern Australia these animals do not constitute a reservoir for any of the

serogroups tested. This is contrary to the situation described in north Queensland by Battey *et al.*¹ and Emanuel *et al.*² who considered that a number of rodent species were responsible for both maintaining and spreading infection.




FIGURE 1. Principle areas of collection (★) of animals in New South Wales and Victoria.

TABLE 1. Prevalence of anti-leptospiral agglutinins in the sera of wildlife in southeastern Australia.

Family	Species	(common name)	Number Examined		Prevalence (%)
			Total	Sero-positive*	
TACHYGLOSSIDAE	<i>Tachyglossus aculeatus</i>	(echidna)	1	—	—
DASYURIDAE	<i>Antechinus stuartii</i>	(brown antechinus)	19	—	—
	<i>Antechinus swainsonii</i>	(swainson's antechinus)	17	—	—
PERAMELIDAE	<i>Perameles nasuta</i>	(long-nosed bandicoot)	8	—	—
	<i>Isodon obesulus</i>	(brown bandicoot)	3	—	—
PHALANGERIDAE	<i>Trichosurus vulpecula</i>	(brush-tailed possum)	25	5	20
PETAURIDAE	<i>Pseudocheirus peregrinus</i>	(common ringtail)	4	—	—
	<i>Scoinoobates volans</i>	(greater glider)	8	—	—
MACROPODIDAE	<i>Macropus eugenii</i>	(tammar wallaby)	26	1	3.8
	<i>Macropus giganteus</i>	(eastern grey kangaroo)	20	—	—
	<i>Macropus rufogriseus</i>	(red-necked wallaby)	23	—	—
	<i>Wallabia bicolor</i>	(swamp wallaby)	12	2	16.7
PHASCOLARCTIDAE	<i>Phascolarctos cinereus</i>	(koala)	3	1	33.3
VOMBATIDAE	<i>Vombatus ursinus</i>	(common wombat)	41	3	7.3
LEPORIDAE	<i>Oryctolagus cuniculus</i>	(rabbit)	25	—	—
MURIDAE	<i>Hydromys chrysogaster</i>	(water rat)	9	—	—
	<i>Rattus rattus</i>	(black rat)	10	—	—
	<i>Rattus fuscipes</i>	(bush rat)	100	2	2
	<i>Rattus lutreolus</i>	(eastern swamp rat)	10	—	—
CANIDAE	<i>Mastacomys fuscus</i>	(broad-toothed rat)	4	—	—
CERVIDAE	<i>Vulpes vulpes</i>	(fox)	20	—	—
	<i>Cervus timorensis</i>	(rusa deer)	12	4	25
	<i>Cervus unicolor</i>	(sambar deer)	6	—	—
	<i>Axis porcinus</i>	(hog deer)	11	—	—
	<i>Dama dama</i>	(fallow deer)	2	—	—

*Antibody titers: *Trichosurus vulpecula*: 1:64, 1:64, 1:64, 1:256 and 1:1024 to hardjo. *Macropus eugenii*: 1:64 to pomona. *Wallabia bicolor*: 1:64 and 1:256 to ballum. *Phascolarctos cinereus*: 1:32 to hardjo. *Vombatus ursinus*: 1:64 to australis, 1:256 to pomona and 1:256 to grippotyphosa. *Rattus fuscipes*: 1:32 to australis and 1:32 to ballum. *Cervus timorensis*: 1:128, 1:128, 1:256 and 1:1024 to hardjo.

Climatic differences between these two areas may provide an explanation for this.

Of the 41 wombats investigated in the present study, only 3 (7.3%) had a positive reaction in the MAT. Unlike other reports, there was no indication of widespread infection in this species with either *pomona* or *hardjo*.^{3,10,11} The high prevalence of reactions to *hardjo* in brush-tailed possums and rusa deer has been reported previously.³ It has been shown that serologic reactions in brush-tailed possums to members of the *Hebdomadis* serogroup were due to infection with *balcanica*, another member of the *Hebdomadis* serogroup. Persistent infection in brush-tailed possums was demonstrated, but transmission appeared to be only from possum to possum.^{3,4} The significance of titers to *hardjo* in rusa deer is unclear but, as these animals were farmed, there are increased opportunities for contact with cattle which, in Victoria, have a high

prevalence of serologic reactions and are the source of frequent isolations of this serovar.⁹

It would seem that leptospirosis is only common in brush-tailed possums and possibly wombats and rusa deer in southeastern Australia. The infecting serovars are members of the *Hebdomadis* serogroup. There is no evidence suggesting that these animals constitute either a human or veterinary health hazard in this area, although Munday¹⁰ postulated that the wombat may be involved in the transmission of *pomona* to domestic animals. Unfortunately, no samples were obtained from either *Rattus norvegicus*, postulated to be a maintenance host of leptospirosis^{1,5} or feral pigs, shown to have a high prevalence of *pomona* infection in New South Wales.⁸ The results of the present study support the earlier suggestion¹¹ that human leptospirosis in Victoria is attributable to occupational contact with domestic animals.

Acknowledgements

The authors wish to thank Kaylene Calvert for her excellent technical assistance and the numerous officers of the Victorian Departments of Agriculture and Fisheries and Wildlife who provided assistance in the collection of specimens. This study was supported by Dairying Research Committee grant DAV 22.

LITERATURE CITED

1. BATTEY, Y.M., D.J.W. SMITH and G. BARROW. 1964. The epidemiology of Leptospirosis in North Queensland. II. Further observations on the hosts in the Mossman district. *J. Hyg. Camb.* 62: 485-494.
2. COLE, J.R., C.R. SULZER and A.R. PURSELL. 1973. Improved microtechnique for the leptospiral microscopic agglutination test. *Appl. Microbiol.* 25: 976-980.
3. DURFEE, P.T. and P.J.A. PRESIDENTE. 1979. A serological survey of Australian wildlife for antibodies to leptospires of the *Hebdomadis* serogroup. *Aust. J. exp. Biol. med. Sci.* 57: 177-189.
4. — and —. 1979. A sero-epidemiological study of *Leptospira interrogans* serovar *balcanica* in four brush-tailed possum populations in Victoria, Australia. *Aust. J. exp. Biol. med. Sci.* 57: 191-201.
5. EMANUEL, M.L., I.M. MACKERRAS and D.J.W. SMITH. 1964. The epidemiology of leptospirosis in North Queensland. I. General survey of animal hosts. *J. Hyg. Camb.* 62: 451-484.

6. GALTON, M.M., C.R. SULZER, C.A. SANTA ROSA and M.J. FIELDS. 1965. Application of a microtechnique to the agglutination test for leptospiral antibodies. *Appl. Microbiol.* 13: 81-85.
7. GLAZEBROOK, J.S., R.S.F. CAMPBELL, G.W. HUTCHINSON and N.D. STALLMAN. 1978. Rodent zoonoses in North Queensland. The occurrence and distribution of zoonotic infections in North Queensland rodents. *Aust. J. exp. Biol. med. Sci.* 56: 147-156.
8. KEAST, J.C., I.R. LITTLEJOHNS, L.C. ROWAN and J.S. WANNAN. 1963. The role of the feral pig as a disease reservoir. *Aust. vet. J.* 39: 99.
9. MILNER, A.R., C.R. WILKS, I.R. MORGAN and N.E. ROSEN. 1980. *Leptospira* serogroup Hebdomadis infection as an Australian zoonosis. *Aust. vet. J.* 56: 70-73.
10. MUNDAY, B.L. 1972. A serological study of some infectious diseases of Tasmanian wildlife. *J. Wildl. Dis.* 8: 169-175.
11. ——— and A. CORBOULD. 1973. *Leptospira pomona* infection in wombats. *J. Wildl. Dis.* 9: 72-73.
12. SHENBERG, E., S. BIRNBAUM, E. RODRIG and M. TORTEN. 1977. Dynamic changes in the epidemiology of canicola fever in Israel. *Am. J. Epidemiol.* 105: 42-48.
13. SULLIVAN, N.D. 1974. Leptospirosis in animals and man. *Aust. vet. J.* 50: 216-223.
14. WILKS, C.R. and A.R. MILNER. 1979. Leptospirosis. *Med. J. Aust.* 1: 396-397.

Received for publication 12 February 1980
