



PANCREATIC INFECTIONS OF *Myxobolus osburni* HERRICK (MYXOZOA: MYXOSPOREA) IN THE PUMPKINSEED, *Lepomis gibbosus* (LINNAEUS) IN IOWA

Authors: INGRAM, K.M., and MITCHELL, L.G.

Source: Journal of Wildlife Diseases, 18(1) : 75-79

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-18.1.75>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

PANCREATIC INFECTIONS OF *Myxobolus osburni* HERRICK (MYXOZOA: MYXOSPOREA) IN THE PUMPKINSEED, *Lepomis gibbosus* (LINNAEUS) IN IOWA

K.M. INGRAM and L.G. MITCHELL, Department of Zoology, Iowa State University, Ames, Iowa 50011, USA.

Abstract: Cysts of *Myxobolus osburni* occurred throughout the pancreas in 119 of 150 (79.3%) pumpkinseed (*Lepomis gibbosus*) from West Lake Okoboji, northwest Iowa during the summer, 1980. None of 341 bluegill (*Lepomis macrochirus*) were infected. Lobate cysts (up to 3 mm diameter) adjacent to the gall bladder contained sporulating plasmodia and mature spore masses. Inflammatory responses in the infected pancreas included engorged blood vessels, endocrine cell nuclei pyknosis, leukocyte (mostly lymphocyte) infiltration, fibroblast proliferation, dark pigment deposition and fibrosis. Host reaction was most pronounced in pancreas containing unencapsulated spore aggregates. Spore variability and plasmodial structure are also described.

INTRODUCTION

The myxozoan genus *Myxobolus* consists of numerous species infecting a wide variety of organs and tissues in many families of fishes.^{1,5,8-12} *Myxobolus osburni* Herrick was described originally from the mesenteries and peritoneum of the smallmouth bass, *Micropterus dolomieu* (Lacépède), and the pumpkinseed, *Lepomis gibbosus* (Linnaeus) from Lake Erie.⁵ This species also was reported from the gall bladder of the bluegill, *Lepomis macrochirus* (Rafinesque), and the black crappie, *Pomoxis nigromaculatus* (Le Seur), from the Lake Okoboji area in northwest Iowa.¹¹ Cone and Anderson¹ described *M. osburni* infections "in or on" endocrine islets and within diffuse exocrine pancreatic tissue in the pumpkinseed. This is the first report of developmental stages of plasmodia and the first report of histopathologic changes associated with pancreatic infections of *M. osburni*. Spore variability and summer prevalence also are reported.

MATERIALS AND METHODS

Fish were collected by fyke net and wire funnel traps in Little Miller's Bay,

West Lake Okoboji, in northwest Iowa from June through September, 1980. Viscera from seven infected pumpkinseed were removed; five were preserved in Bouin's and two in Stieve's fixative. Tissue was paraffin embedded, serially sectioned at 6 to 7 μ m, and stained with hematoxylin and eosin. Spores were examined in smears of 12 individual plasmodia in 0.6% NaCl. All measurements, recorded in micrometers, are of unstained spores.

RESULTS AND DISCUSSION

One hundred nineteen of 150 (79.3%) pumpkinseed were infected with *Myxobolus osburni*. Both sexually mature and immature fish (std. length 7.5 - 13.0 cm) were infected. Prevalence was highest in sexually mature fish, and was the same in females (74 infected/89 examined; 83%) and males (35/42; 83%). These are the highest prevalence values yet reported for *M. osburni*.^{1,5} Prevalence was considerably lower in sexually immature pumpkinseed (10 infected/19 examined; 53%). These results are generally similar to those of Cone and Anderson² who found prevalence of *M. osburni* to increase with age of hosts. Previous

estimates of prevalence are probably low. In sections we have observed microscopic cysts in the pancreas that appeared uninfected in routine examination.

We did not find cysts in 341 bluegills (*Lepomis macrochirus*) examined macroscopically during the study period. Microscopic examination of the viscera of 10 bluegills did not reveal *M. osburni* infections.

The gall bladder and bile ducts contained a few scattered spores. The report of Otto and Jahn¹¹ of microscopic *M. osburni* trophozoites floating free in bile in one bluegill seems questionable for this histozoic myxozoan.

Spores. Spores (Fig. 1) closely resembled those described by Herrick.⁵ Seven to 10 sutural folds were distinct in most spores. Two cnidocyst discharge canals appeared as distinct anterior opacities. The angle of filament coiling was approximately 45° as drawn by Herrick,⁵ although some spores showed filaments coiled at about 90° to the long axis of the spores. Dimensions of the Iowa population were: length 9-11 (mean=10, n=650); breadth 10-13 (mean=11.5, n=650); thickness 6-8 (mean=7, n=79). The majority of spores were elliptical in front view (slightly broader than long) as drawn by Herrick.



FIGURE 1. *Myxobolus osburni* spore, unstained, 830X.

Cone and Anderson¹ reported longer, spherical spores: length 10-13 (mean=12, n=40); breadth 10-14 (mean=12, n=40); thickness 6-7 (mean=6.5, n=5). We have seen some spherical spores in plasmodia containing predominantly elliptical forms, indicating considerable intraspecific variation in this myxobolid. The total range of spore size recorded is 9-13 × 9.5-14 × 6-8.

Plasmodia. Creamy-white to yellowish lobate cysts (up to 3 mm in diameter) were observed in pancreatic tissue in mesenteries posteromedial to the gall bladder (Fig. 2). One (rarely two) of these large cysts was seen per infected pumpkinseed. Previous reports,^{1,5} described cysts containing only mature spores anterior to the gall bladder. In our sections, single or diffuse aggregates of microscopic cysts containing mature spores in exocrine and endocrine pancreas were in the vicinity of the gall bladder and bile duct.

Lobate cysts (Figs. 2, 3) of individual sporulating plasmodia and/or fully ripened micro- and macroscopic spherical or multilocular spore masses usually were encapsulated by connective tissue.

Immature (sporulating) plasmodia showed discrete striated eosinophilic ectoplasm. The endoplasm contained an outermost region of undifferentiated




FIGURE 2. Infected pumpkinseed showing large lobate cyst behind gall bladder. Approximately life size. Arrow indicates cyst.




FIGURE 3. Section of lobate cyst adjacent to gall bladder showing three sporulating plasmodia with core of mature spores (sp) in pancreatic islet tissue (pi). Arrow indicates thin border of exocrine pancreatic tissue. H&E. 35 \times .




FIGURE 4. Edges of three plasmodia showing striated ectoplasm (ec), sporulating endoplasm (en), and mature spores (sp). Note diffuse leukocytic infiltration between plasmodia. H&E. 260 \times .

nuclei, an intermediate zone of pan-sporoblasts and a central core of mature spores (Fig. 4).

Histopathology. Sporulating plasmodia within lobate cysts typically were surrounded by normal pancreatic islet cells in direct contact with plasmodial ectoplasm. The edges of developing plasmodia were bordered by layers of flattened islet cells, fibroblasts, or a thin connective tissue capsule. Several plasmodia seemed to have developed within a single islet, replacing endocrine cells. Enlarged islet capillaries adjacent to plasmodia suggested some cysts had developed from intracapillary infections. Mature spore masses (Fig. 5) were usually enclosed in a connective tissue capsule (up to 10 μ m thick).

Isolated spores and capillaries were common in these capsules.

A diffuse inflammatory response around most plasmodia and encapsulated spore masses was manifested as engorged small blood vessels, pyknotic endocrine cell nuclei and diffuse leukocytic infiltration (Fig. 4). Granulocytes and lymphocytes were common in enlarged capillaries. Melanin macrophage involvement was indicated by occasional dark pigment deposition in cyst capsules. Pancreas containing unencapsulated spore masses had developed granulomatous areas showing infiltration of lymphocytes and macrophages, proliferation of fibroblasts and fibrosis among endocrine and exocrine cells (Fig. 6).




FIGURE 5. Encapsulated spore mass surrounded by normal exocrine pancreas; pancreatic islet (pi). H&E. 270 \times .

Heckmann and Jensen found unencapsulated spore masses of *Henneguya sebasta* associated with necrosis, hyperplasia, hypertrophy, and proliferation of connective tissue in smooth muscle of the bulbus and truncus arteriosus in bocaccio, *Sebastes paucispinis* (Ayres).¹ These authors also reported severe atrophy and necrosis of skeletal muscle associated with cyst development and encapsulation of spores of *Kudoa clupeidae* in bocaccio. Dykova and Lom¹ described inflammatory responses leading to cyst degradation and granuloma formation in late stages of




FIGURE 6. Leukocytic infiltration (l) of pancreas containing unencapsulated spore mass (sp). Exocrine pancreatic cells (ex) containing zymogen granules. H&E. 230 \times .

gill-inhabiting species of *Henneguya*. Epithelioid granuloma formation has been reported surrounding spore masses of the cartilage-invading myxozoans, *Myxosoma cartilaginis* and *M. cerebralis*.^{6,7} All of these studies indicate that while growing plasmodia of myxozoans may cause tissue destruction, host reactions to these parasites become most pronounced after spore maturation. Experimental studies with other *Myxobolus* (= *Myxosoma*) species indicate that several weeks or months may elapse after plasmodia appear in tissue before cellular reactions appear.^{6,7} The occurrence of developing plasmodia and mature spore masses in granulation tissue within large lobate cysts of *M. osburni* indicates that chronic infections do not elicit host protective immunity in the pumpkinseed.

Acknowledgements

Partial support for this research was provided by the Iowa State University Graduate College. We wish to thank Thomas Rice and Mark Wehrspann for technical assistance. This study was conducted at the Iowa Lakeside Laboratory, Milford, Iowa.

LITERATURE CITED

1. CONE, D.K. and R.C. ANDERSON. 1977. Myxosporidan parasites of pumpkinseed (*Lepomis gibbosus* L.) from Ontario. J. Parasit. 63: 657-666.
2. ——— and ———. 1977. Parasites of pumpkinseed (*Lepomis gibbosus* L.) from Ryan Lake, Algonquin Park, Ontario. Can. J. Zool. 55: 1410-1421.
3. DYKOVA, I. and J. LOM. 1978. Histopathological changes in fish gills infected with myxosporidan parasites of the genus *Henneguya*. J. Fish Biol. 12: 197-202.
4. HECKMANN, R.A. and L.A. JENSEN. 1978. The histopathology and prevalence of *Henneguya* *sebasta* and *Kudoa* *clupeidae* in the rockfish, *Sebastes paucispinis* of southern California. J. Wildl. Dis. 14: 259-262.
5. HERRICK, J.A. 1936. Two new species of *Myxobolus* from fishes of Lake Erie. Trans. Am. Microsc. Soc. 55: 194-198.
6. HOFFMAN, G.L., C.E. DUNBAR and A. BRADFORD. 1962. Whirling disease of trouts caused by *Myxosoma cerebralis* in the United States. U.S. Fish Wildl. Serv., Spec. Sci. Rep. —Fish. 427: 1-15.
7. ———, R.E. PUTZ and C.E. DUNBAR. 1965. Studies on *Myxosoma cartilaginis* n.sp. (Protozoa: Myxosporidea) of centrarchid fish and a synopsis of the *Myxosoma* of North American freshwater fishes. J. Protozool. 12: 319-332.
8. KUDO, R.R. 1919. Studies on Myxosporidia. A synopsis of genera and species. Ill. Biol. Monogr. 5: 1-265.
9. LOM, J. 1969. On a new taxonomic character in Myxosporidia, as demonstrated in descriptions of two new species of *Myxobolus*. Folia Parasit. (Praha). 16: 97-103.
10. MITCHELL, L.G. 1978. Myxosporidan infections in some fishes of Iowa. J. Protozool. 25: 100-105.
11. OTTO, G.R. and T.L. JAHN. 1943. Internal myxosporidian infections of some fishes of the Okoboji region. Proc. Iowa Acad. Sci. 50: 323-335.
12. SHULMAN, S.S. and G.A. SHTEIN. 1962. Phylum Protozoa. pp. 5-235. In: *Key to Parasites of Freshwater Fish of the USSR*. Bykovskaya-Pavlovskaya et al., eds. Akad. Nauk USSR, Moscow-Leningrad (Israel Program for Scientific Translations. 1964. OTS, U.S. Dept. of Commerce).

Received for publication 13 February 1981