

A REVIEW OF DISEASES OF PARASITES OF THE KAFUE LECHWE (KOBUS LECHE KAFUENSIS)

Author: Stafford, K. J.

Source: Journal of Wildlife Diseases, 27(4) : 661-667

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-27.4.661>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A REVIEW OF DISEASES OF PARASITES OF THE KAFUE LECHWE (*KOBUS LECHWE KAFUENSIS*)

K. J. Stafford

Department of Clinical Studies, Samora Machel School of Veterinary Medicine,
UNZA, P.O. Box 32379, Lusaka, Zambia

ABSTRACT: The diseases of the Kafue lechwe (*Kobus lechwe kafuensis*) are reviewed in this paper. Kafue lechwe are an important natural resource for Zambia. Bovine tuberculosis is widespread within the lechwe population and they are host to many parasites, especially the warble *Strobiloestrous vanzyli*.

Key words: *Kobus lechwe kafuensis*, bovine tuberculosis, *Mycobacterium bovis*, endoparasites, *Strobiloestrous vanzyli*, ectoparasites, review.

INTRODUCTION

Kafue lechwe (*Kobus lechwe kafuensis*) are found in the natural state only within the Kafue Flats area of Zambia. The 65,000 lechwe (Howard et al., 1988) are a "natural resource of national importance" for Zambia (Vesey-Fitzgerald, 1965). They are utilized as a source of meat, hides and trophies. They are important as a tourist attraction and in maintaining the fertility of the Kafue river flats and fisheries (Mitchell and Uys, 1961; Howard, 1977). Lechwe have been domesticated (Bainbridge, 1970) and are being stocked onto game farms in Zambia. In the past lechwe were cropped for meat by traditional kills (Ansell, 1955) and controlled culling, utilizing an abattoir, was carried out for a short period in the late 1960's (Bainbridge, 1970; Pullan, 1983).

This review is to provide veterinarians and biologists with information on the diseases and parasites of Kafue lechwe.

INFECTIOUS DISEASES

Tuberculous lesions of bone were found in a Kafue lechwe (Leroux, 1958) and pulmonary tuberculosis caused by *Mycobacterium bovis* has been diagnosed in lechwe (Gallagher et al., 1972; Macadam et al., 1974; Clancey, 1977). The prevalence of tuberculosis in lechwe has varied (Table 1) but recent work would indicate that about 16% of adult male animals are affected (Stafford et al., 1990). Lechwe are semi-aquatic animals living in large groups

which become concentrated during the rains and early dry season (November to June) when their range is flooded (Gallagher et al., 1972). Crowding is important in the transmission of tuberculosis (Thoen and Hines, 1981) and Sayer and Van Lavierien (1975) speculated that the disease spread when the lechwe were concentrated at the beginning of the dry season and were suffering from "nutritional stress" (Gallagher et al., 1972). There are few predators in the area inhabited by the Kafue lechwe. Gallagher et al. (1972) estimated that tuberculosis killed 20% of the lechwe population each year and may be the major factor limiting population growth.

Lechwe are in contact with cattle when they graze together on the fresh grass which grows as the flood recedes. Cattle may have introduced tuberculosis into the lechwe population (Dillmann, 1976; Rottcher, 1976); $\leq 25\%$ of cattle from the Kafue Flats showed tubercular lesions on meat inspection (Anonymous, 1965) and $\leq 40\%$ may be affected (Rottcher, 1976). Infected lechwe may introduce the disease to cattle and other game species if they are moved onto cattle-game farms. When lechwe are killed for meat there is a potential public health problem as $\leq 80\%$ of carcasses may be tuberculous (Macadam et al., 1974; Dillmann, 1976; Krauss et al., 1986). The control of tuberculosis amongst free-ranging lechwe on the Kafue Flats would be extremely difficult. Lechwe being moved

TABLE 1. Tuberculosis in the Kafue lechwe.

Number examined	Number affected	Percent affected	Reference
125	45	36	Gallagher et al., 1972
300	90	30	Dillmann, 1976
86	29	34	Gallagher et al., 1972
7	7	100 ^a	Clancey, 1977
63	33	52	Clancey, 1977
141	46	33	Krauss et al., 1986
41	33	80 ^a	Krauss et al., 1986
147	5	3 ^b	Krauss et al., 1986
92	15	16	Stafford et al., 1990

^a Cachectic animals with suspected tuberculosis.^b Dead animals collected from Kafue Flats.

onto game farms could be tested for tuberculosis; however the efficacy of testing lechwe for tuberculosis either by skin testing (Macadam, 1973) or serological testing is unknown. Tuberculosis was not a problem among Black lechwe (*Kobus lechwe smithemani*); they inhabit an environ-

ment similar to the Kafue lechwe but do not come in contact with cattle (Grimsdell and Bell, 1975).

Serological surveys have been conducted to determine the disease status of lechwe (Table 2). Antibodies against brucella organisms, the *Leptospira interrogans* serovars *autumnalis autumnalis* and *ballum ballum*, *Pseudomonas pseudomallei*, *Francisella tularensis*, *Chlamydia psittaci*, *Coxiella burnetii*, infectious bovine rhinotracheitis/infectious pustular vaginitis virus, parainfluenza 111 virus and reoviruses 1,11,111 were found in <10% of lechwe (Table 2). Antibodies against bluetongue virus were found in 50% of a small number of lechwe. The significance of these findings to the health of lechwe and domestic animals in contact with them are unclear (Krauss et al., 1986) but may become more obvious as lechwe are moved onto cattle-game farms. Foot and Mouth

TABLE 2. Serological Surveys of Kafue lechwe.

Infectious agent	Number of samples	Results	Reference
Leptospirosis ^a	45	3 positive ^a	K. Stafford (unpubl. data)
Brucellosis ^b	200	positive ^c	Krauss et al., 1986
Brucellosis ^d	55	2 positive ^c	K. Stafford (unpubl. data)
<i>Mycobacterium paratuberculosis</i>	200 ^c	all negative	Krauss et al., 1986
<i>Anaplasma marginale</i>	31	all negative	Dillmann, 1976
<i>Anaplasma centrale</i>	31	all negative	Dillmann, 1976
<i>Pseudomonas pseudomallei</i>	200 ^c	positive ^e	Krauss et al. 1986
<i>Francisella tularensis</i>	200 ^c	positive ^e	Krauss et al. 1986
<i>Chlamydia psittaci</i>	200 ^c	positive ^e	Krauss et al. 1986
<i>Coxiella burnetii</i>	200 ^c	positive ^e	Krauss et al. 1986
Infectious Bovine Rhinotracheitis	10 ^c	positive ^e	Krauss et al. 1986
Parainfluenza 111	10 ^c	positive ^e	Krauss et al. 1986
Reovirus 1, 11, 111	5 ^c	positive ^e	Krauss et al. 1986
Bluetongue	10	5 positive	Krauss et al. 1986
<i>Babesia bigemina</i>	31	all negative	Dillmann, 1976
<i>Theileria parva</i>	31	all negative	Dillmann, 1976
<i>Theileria mutans</i>	31	all negative	Dillmann, 1976

^a The microagglutination test was used to test for antibodies against the following serovars: *Leptospira interrogans* serovar, *canticola canticola*, *icterohaemorrhagiae copenhageni*, *pomona pomona*, *grippotyphosa grippotyphosa*, *tarrasovi tarrasovi*, *australis australis*, *australis bratislava*, *sejroe hardjo*, *hebdomadis hebdomadis*, *mini mini*, *bataviae bataviae*, *javanica javanica*, *pyrogenes pyrogenes*, *autumnalis autumnalis*, *ballum ballum*, *celledoni celledoni*, *cynopteri cynopteri* and *shermani shermani*. Three were positive for *autumnalis autumnalis* and one for *Ballum ballum*.

^b Serum agglutination test.

^c The exact number tested was not stated.

^d Rose Bengal test and slide agglutination test.

^e Positive to both tests (d).

^f The exact number of positive sera was not stated.

TABLE 3. Nematode parasites of Kafue lechwe.

<i>Oesophagostomum lechwei</i>	(Leroux, 1955)
<i>Bunostomum cobí</i>	(Leroux, 1958)
<i>Bunostomum</i> sp.	(Leroux, 1932)
<i>Trichostrongylus colubriformis</i> a	(Leroux, 1930)
<i>Haemonchus bedfordi</i>	(Leroux, 1955)
<i>Haemonchus contortus</i> a	(Leroux, 1930)
<i>Cooperia curticei</i>	(Leroux, 1932)
<i>Dictyocaulus filaria</i>	(Fitzsimmons, 1961)
<i>Dictyocaulus viviparus</i>	(Robinette and Child, 1964)
<i>Thelazia rhodesi</i>	(Robinette and Child, 1964)
<i>Thelazia</i> sp.	(Leroux, 1955)
<i>Onchocerca</i> sp.	(Leroux, 1947, 1955)
<i>Setaria bicornata</i>	(Yeh, 1959)
<i>Setaria boulengeri</i>	(Yeh, 1959)
<i>Setaria hornbyi</i>	(Yeh, 1959)
<i>Setaria</i> sp.	(Leroux, 1955)
<i>Trichuris</i> sp.	(Ratcliffe, 1930)

a Round (1968) questioned this finding.

disease has not been diagnosed clinically in Kafue lechwe and the causative virus was not isolated from 50 animals sampled by Dillmann (1976). Lechwe are probably susceptible to the disease but their importance as a source of the disease for domestic animals is unknown (Howard, 1977). Heartwater disease caused by *Cowdria ruminantium* was diagnosed in lechwe with heavy loads of *Amblyomma variegatum* ticks held on a game farm (G. S. Pandey, pers. comm.). Abscesses were found in the testes of three lechwe (Robinette and Child, 1964) but the lesions were not examined bacteriologically.

HELMINTH PARASITES

Kafue lechwe are host to many parasites (Tables 3, 4, 5; Round, 1968). The effect of these parasites on lechwe is unknown but Robinette and Child (1964) doubted if internal parasites were primary cause of mortalities. Strongyle nematode eggs were found in 74 of 76 faecal samples taken from recently shot adult lechwe (K. Stafford, unpubl. data). The number of eggs per gram were <100 in 68 of these samples. Coccidial oocysts were found in many

TABLE 4. Trematode and cestode parasites of Kafue lechwe.

<i>Fasciola gigantica</i>	(Gallagher et al., 1972; Wright et al., 1979a)
<i>Fasciola</i> sp.	(Leroux, 1957)
<i>Cotylophoron cotylophoron</i>	(Leroux, 1932)
<i>Cotylophoron</i> sp.	(Robinette and Child, 1964)
<i>Carmerius</i> sp. ^a	(Wright et al., 1979b)
<i>Gastrothylax crumenifer</i>	(Leroux, 1932)
<i>Paramphistomum cervi</i> ^b	(Leroux, 1932)
<i>Gigantocotyle explanatum</i>	(Leroux, 1932)
<i>Gigantocotyle formosanum</i>	(Round, 1968)
<i>Gigantocotyle lerouxi</i>	(Yeh, 1957)
<i>Choerocotylodes onotragi</i>	(Prudhoe et al., 1964)
<i>Homalogaster paloniae</i>	(Leroux, 1955)
<i>Stephanopharynx compactus</i>	(Wright et al., 1979b)
<i>Calicophoron calicophoron</i>	(Wright et al., 1979a)
<i>Schistosoma leiperi</i>	(Leroux, 1955)
<i>Schistosoma margrebowiei</i>	(Leroux, 1933, 1955)
<i>Schistosoma matthei</i>	(Leroux, 1958)
<i>Schistosoma spindalis</i> ^c	(Ortlepp, 1961)
<i>Taenia</i> spp.	(Leroux, 1955, 1957; Robinette and Child, 1964)

^a May be *C. mancupatus* or *C. spastosis* (Wright et al., 1979b).

^b May be *P. microbothrium* or *P. phillierouxi* (Wright et al., 1979b; Round, 1968).

^c Round (1968) questioned this finding.

of these faecal samples. Coccidia have not previously been reported in lechwe. Lungworms of three different species were found in about 55% of culled lechwe (Dillmann, 1976) but only two species have been definitely identified (Table 3). *Thelazia rhodesi* found in small numbers in a

TABLE 5. Ectoparasites of Kafue lechwe.

<i>Strobiloestrous vanzyli</i>	(Zumpt, 1961; Howard, 1975, 1980, 1985)
<i>Damalinia</i> sp.	(Robinette and Child, 1964)
<i>Amblyomma variegatum</i>	(Colbo, 1973)
<i>Rhipicephalus evertsi</i>	(Colbo, 1973)
<i>Rhipicephalus sanguineus</i>	(Colbo, 1973)
<i>Rhipicephalus appendiculatus</i>	(Robinette and Child, 1964)
Muscoid headflies	(Robinette and Child, 1964)

few lechwe (Robinette and Child, 1964) may be the filarial worms referred to by Sayer and Van Lavieren (1975) as the cause of blindness in lechwe. *Onchocerca* sp. probably *O. gibsoni* were found in fibrous lesions in periarticular regions especially near the stifle joint (Leroux, 1947, 1955; Robinette and Child, 1964). Approximately 35% of the lechwe in Lochinvar National Park were affected by onchocerciasis which caused lameness in some animals (Dillmann, 1976). *Setaria* sp. were "abundant" in lechwe (Sayer and Van Lavieren, 1975; Robinette and Child, 1964; Dillmann, 1976) and three species have been identified (Table 3). *Fasciola gigantica* were identified in adult lechwe (Wright et al., 1979a). Female lechwe had a higher prevalence of liver fluke (22%) than males (6%) possibly due to the female habit of grazing more in the marsh along the water's edge (Gallagher et al., 1972). However *Fasciola* eggs were found in 68 of 76 (89%) faecal samples taken from male carcasses and examined by K. Stafford (unpubl. data). Amphistome flukes were regularly seen in lechwe (Table 4). *Schistosoma margrebowiei* have been found in young lechwe (Leroux, 1933, 1955) and *S. leiperi* in older animals (Wright et al., 1979a; Howard et al., 1982). Lechwe <1 yr of age had a mean infection of 24 *S. margrebowiei* and no *S. leiperi* while adult animals >4 yr of age had a mean infection of 273 *S. leiperi* and no *S. margrebowiei*. One adult lechwe had over 1,000 *S. leiperi* (Howard et al., 1982). *S. matthei* found in lechwe by Leroux (1958) was not identified by Howard et al. (1982) in lechwe sharing pasture with cattle, but was seen in waterbuck (*Kobus defassa*) living remote from cattle.

ECTOPARASITES

The oestrid larvae of *Strobiloestrous vanzyli*, the southern lechwe fly, were found in the subcutis of the back and hind-quarters of lechwe (Zumpt, 1961; Sayer and Van Lavieren, 1975; Howard, 1980). In a sample of 526 lechwe of all ages, 95%

of those <1-yr-old were affected with warbles (\bar{x} = 149 warbles/animal) while only 20% of older animals were affected (\bar{x} = 8.6 warbles/host) (Howard, 1985). The warbles are on the lechwe from November until September when the prepupae emerge, drop on the ground and pupate in the soil. Adult flies emerge within 1 mo (Howard, 1985). *S. trobiloestrous vanzyli* may not be distinct from *S. erikssoni* (Howard, 1980). Leroux (1942) reported finding similar warbles on lechwe and cattle and concluded that they were from the same species of fly. The large numbers of larvae seen on young lechwe led Mitchell and Uys (1961) to conclude that the warbles caused "heavy" mortalities but evidence to support this conclusion is lacking. A small unidentified muscoid fly was seen to be a nuisance to lechwe during the warmer months of October and November. Lechwe sought relief from the fly by rubbing their heads in grass or standing in a tight circle (Robinette and Child, 1964). Lice (*Damalinea* sp.) were common on lechwe (Robinette and Child, 1964). Various species of ticks have been found on lechwe (Table 5) (Ganagarajah, 1976). *Rhipicephalus appendiculatus* was found in small numbers (<5) only on the perianal region of 20 of 75 lechwe examined, the remaining 55 animals had no ticks (Robinette and Child, 1964).

MISCELLANEOUS PATHOLOGICAL CONDITIONS

A fibrosarcoma and other malignant but unidentified neoplasms were found in lechwe by Robinette and Child (1964).

DISCUSSION

The population of Kafue lechwe has dropped from 94,075 in 1970 to 65,000 in 1988 (Howard et al., 1988). This reduction in numbers has been due to a loss in available habitat, poaching and perhaps disease and malnutrition.

Tuberculosis and warbles are the major disease problems of lechwe. Tuberculosis was seen in most cachectic animals ex-

amined (Table 1). However it was only seen in 3% of 147 autopsies conducted on lechwe found dead (Krauss et al., 1986), the majority had died as a result of poaching injury. The significance of the effect of warbles and other internal parasites on the health of lechwe is unknown. The lameness seen in many lechwe may be due to onchocerciasis and/or poaching injuries.

To maintain the present population of lechwe attempts may have to be made to control tuberculosis. Increasing the available habitat and reducing contact between cattle and lechwe have been suggested as means of reducing mortalities due to tuberculosis and malnutrition (Rottcher, 1976). A suggestion by Dillmann (1976) that the culling of 5,000 lechwe/yr might reduce tuberculosis has not been confirmed. However the present level of tuberculosis is much lower than that seen in the seventies and this might be explained in part by the fact that lechwe numbers dropped to 41,000 in 1982 (Howard et al., 1988). Research should be directed towards examining the accuracy and feasibility of tuberculosis testing of lechwe in the wild, in transit and on game farms. The monitoring of lechwe held on game farms stocking cattle free from tuberculosis may demonstrate whether the disease can be maintained in a lechwe population isolated from tubercular cattle.

The limited number of serological surveys conducted suggests the need to evaluate further the prevalence of antibodies to bluetongue, Rift Valley Fever, Bovine Herpes virus and other diseases within the lechwe population. Lechwe are being moved onto farms which stock domestic ruminants. It is important the the transmission of disease from lechwe to domestic ruminants and vice versa, if it occurs, be monitored. Will the warbles of *Strobiloestrous vanzyli* affect cattle (Leroux, 1942) and *Schistosoma matthei* infest lechwe on cattle-game farms? Disease problems such as heartwater and heavy tick infections not reported from wild lechwe may become problems on game farms. The cropping

scheme proposed and supported by the World Wildlife Fund and initiated in 1989 is an opportunity to study further the disease conditions seen in adult lechwe. As the lechwe will be cropped for meat, post-mortem examination will have to be carried out to ensure the quality of the meat being sold. Blood obtained from the cropped animals could be used for serological studies. The cropping and farming of lechwe may be necessary to ensure the survival of the species in Zambia. Both these enterprises demand that lechwe are healthy, fit for human consumption and are not a source of disease for other wild and domestic stock held on game farms.

LITERATURE CITED

- ANONYMOUS. 1965. Annual Report of the Department of Veterinary and Tsetse Control Services. Ministry of Agriculture, Lusaka, Zambia, 32 pp.
- ANSELL, W. F. 1955. The declining red lechwe. *The Oryx* 3: 15-18.
- BAINBRIDGE, W. A. 1970. Report of the Chief Game Officer, Game and Fisheries Annual Report for the Year 1968. Ministry of Natural Resources and Tourism, Lusaka, Zambia, 72 pp.
- CLANCEY, J. K. 1977. The incidence of tuberculosis in lechwe (marsh antelope). *Tubercle* 58: 151-156.
- COLBO, M. H. 1973. Ticks of Zambian wild animals: A preliminary checklist. *The Puku* 7: 97-105.
- DILLMANN, J. S. 1976. Compiled final report. Veterinary Wildlife Research Section, Department of Veterinary Services, Ministry of Agriculture, Lusaka, Zambia, 395 pp.
- FITZSIMMONS, W. M. 1961. A possible dwarf form of *Dictyocaulus filaria* from a waterbuck, *Kobus leche* in northern Rhodesia. *Veterinary Record* 73: 1111-1112.
- GALLAGHER, J., I. MACADAM, J. SAYER, AND J. P. VAN LAVIEREN. 1972. Pulmonary tuberculosis in free living lechwe antelope in Zambia. *Tropical Animal Health and Production* 4: 204-213.
- GANAGARAJAH, M. 1976. A preliminary check list and host list of ticks of Zambia. Published by National Council for Scientific Research, Lusaka, Zambia, 16 pp.
- GRIMSDALL, J. J., AND R. H. BELL. 1975. Black lechwe research project, final report. Published by National Council for Scientific Research, Lusaka, Zambia, 175 pp.
- HOWARD, G. 1975. Differentially delayed development in the lechwe warble, *Strobiloestrous vanzyli* (Diptera:Oestridae). *Parasitology* 71(2): 10.

- . 1977. National Parks of the Kafue Basin. In *Development and ecology in the lower Kafue Basin in the 1970s*, G. Williams and G. Howard (eds.). The University of Zambia, Lusaka, Zambia, pp. 47–56.
- . 1980. Immature stages and affinities of the southern Lechwe warblefly, *Strobiloestrous vanzyli* Zumpt (Diptera: Oestridae). *Journal of Natural History* 14: 669–683.
- . 1985. Warble infestations of lechwe antelope. *Proceedings of the Fifth International Conference on Wildlife Diseases*, Uppsala, Sweden, 23 pp.
- , C. MALAMBO, AND R. C. JEFFERY. 1988. Kafue lechwe population census May/June, 1988. A report to the Director, National Parks and Wildlife Service, Lusaka, Zambia, 8 pp.
- , C. A. WRIGHT, AND V. R. SOUTHGATE. 1982. Schistosome infections of lechwe and waterbuck in Zambia—A preliminary report. In *Wildlife Diseases of the Pacific Basin and other countries*, M. E. Fowler (ed.). Wildlife Disease Association, Ames, Iowa, pp. 136–138.
- KRAUSS, H., D. ROTTCHER, D. WEISS, K. DANNER, AND O. J. HUBSCHLE. 1986. Wildlife as a potential source of infection in domestic animals—Studies on game in Zambia. *Animal Research and Development* 24: 41–58.
- LEROUX, P. L. 1930. Helminths of domestic stock in the Republic of South Africa. *Journal of South African Veterinary Medical Association* 1(4): 43–65.
- . 1932. List of helminths collected from mammals and birds in the Mazabuka area of northern Rhodesia. *Annual Report (1931) of the Department of Animal Health, Northern Rhodesia*, Appendix 3: 31–34.
- . 1933. A preliminary note on *Bilharzia margrebowiei*, a new parasite of ruminants and possibly man in northern Rhodesia. *Journal of Helminthology* 11: 57–62.
- . 1942. Report of the Associate Veterinary Research Officer. *Annual report of the Department of Animal Health, Northern Rhodesia*, 32 pp.
- . 1947. Onchocerca in the tendons of insertion of the M. rectus femoris of a Red Lechwe (*Onotragus lechwe*) a species of Kob, Kafue River, Mazabuka, northern Rhodesia. *Transcripts of the Royal Society of Tropical Medicine and Hygiene* 41: 9.
- . 1955. Game conservation and severe helminthiasis of stock in national game reserves. *Transcripts of the Royal Society of Tropical Medicine and Hygiene* 49: 293.
- . 1957. Report to the government of the Federation of Rhodesia and Nyasaland on the control of parasite diseases in livestock. *FAO Report number 696*, Rome, Italy, 47 pp.
- . 1958. Bone tuberculosis in a Red Lechwe (*Adenota lechwe*) in northern Rhodesia. *Transcripts of the Royal Society of Tropical Medicine and Hygiene* 52: 15.
- MACADAM, I. 1973. Handling lechwe for vaccination and tuberculosis testing. *Veterinary Record* 92: 461.
- , J. GALLAGHER, AND J. MCKAY. 1974. Experimental tuberculosis in lechwe antelope in Zambia. *Tropical Animal Health and Production* 6: 107–109.
- MITCHELL, B. L., AND J. M. UYS. 1961. The problem of the lechwe (*Kobus lechwe*) on the Kafue Flats. *The Oryx* 6: 171–183.
- ORTLEPP, R. J. 1961. South African helminths. *Orderstepoort Journal of Veterinary Science* 9: 311–336.
- PRUDHOE, S., L. YEW, AND L. F. KHALI. 1964. A new amphistome from ruminants in northern Rhodesia. *Journal of Helminthology* 38: 57–62.
- PULLAN, R. A. 1983. The use of wildlife in the development of Zambia. In *Natural resources in tropical countries*, Ooi Jin Bee (ed.). Singapore University Press, Singapore, Malaysia, pp. 267–325.
- RATCLIFFE, H. C. 1930. Animal parasites. In *Report of the Zoological Society of Philadelphia*, H. Fox (ed.). Philadelphia, Pennsylvania, pp. 57–60.
- ROBINETTE, W. L., AND G. F. CHILD. 1964. Notes on the biology of the Lechwe (*Kobus lechwe*). *The Puku* 2: 84–117.
- ROTTCHER, D. 1976. Research on wildlife diseases in Zambia. *Proceedings of the fourth regional conference on wildlife for East and Central Africa*, Ministry of Lords, Natural Resources and Tourism, Lusaka, Zambia, pp. 147–157.
- ROUND, M. C. 1968. Checklist of the helminth parasites of African mammals. Technical Communication, number 38, Commonwealth Bureau of Helminthology, The Commonwealth Agricultural Bureau, England, 25 pp.
- SAYER, J. A., AND L. P. VAN LAVIEREN. 1975. The ecology of the Kafue lechwe population of Zambia before the operation of hydroelectric dams on the Kafue River. *East African Wildlife Journal* 13: 9–37.
- STAFFORD, K., A. MWEENE, J. BAER, AND G. PANDEY. 1990. Tuberculosis in Kafue lechwe (*Kobus lechwe kafuensis*). *Sixth International Conference on Wildlife Diseases*. Berlin, German Democratic Republic, 76 pp.
- THOEN, C. O., AND E. M. HINES. 1981. Tuberculosis. In *Infectious diseases of wild animals*, J. W. Davis, L. H. Karstad and D. O. Trainer (eds.). Iowa State University Press, Ames, Iowa, pp. 263–274.
- VESEY FITZGERALD, L. D. 1965. Lechwe pastures. *The Puku* 3: 143–147.
- WRIGHT, C., V. SOUTHGATE, AND G. HOWARD. 1979a. Observations on the life cycle of *Schistosoma margrebowiei* and its possible interac-

- tions with *S. leiperi* in Zambia. *Journal of Natural History* 13: 499–506.
- , ———, AND ———. 1979b. A note on the life cycles of some amphistome flukes in Zambia. *Journal of Helminthology* 53: 251–252.
- YEH, L. S. 1957. On a new paramphistomid trematode *Gigantocotyle lerouxi* from the stomach of a Red Lechwe (*Onotragus lechwe*) from northern Rhodesia. *Parasitology* 47: 432–434.
- . 1959. On the identity of the filarial worms. *Transcripts of the Royal Society of Tropical Medicine and Hygiene* 52: 297.
- ZUMPT, F. 1961. The enigma of *Strobiloestrous brauer* (Diptera: Oestridae). *Proceedings of the Royal Entomological Society, London* 30: 93–102.

Received for publication 6 December 1989.