



Gnathostomiasis in Wild Boars from Japan

Authors: Ishiwata, Kenji, and Díaz Camacho, Sylvia Páz

Source: Journal of Wildlife Diseases, 34(1) : 155-157

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-34.1.155>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Gnathostomiasis in Wild Boars from Japan

Kenji Ishiwata,¹ Sylvia Páz Díaz Camacho,² Amrozi,³ Yoichiro Horii,³ Natsuki Nawa,⁴ and Yukifumi Nawa¹

¹ Department of Parasitology, Miyazaki Medical College, Miyazaki 889-16, Japan; ² Department of Public Health Research, Faculty of Chemical and Biological Sciences, University of Sinaloa, Culiacan, Mexico; ³ Department of Veterinary Internal Medicine, Faculty of Agriculture, Miyazaki University, Gakuen-Kibanadai 880-12, Japan; ⁴ Department of Oceanology, Faculty of Science, University of Ryukyus, Nishihara, Okinawa 903-01, Japan

ABSTRACT: *Gnathostoma doloresi*, a nematode parasite producing ulcerative and/or granulomatous lesions in the gastric wall of domestic pigs and wild boars (*Sus scrofa leucomystax*) throughout southeastern Asia, is an important zoonosis causing cutaneous larva migrans in humans. Patients infected with *G. doloresi* have been found yearly in the vicinity of Miyazaki Prefecture, Japan. Adult nematodes were found in 31 of 32 wild boars captured at two different geographic locations in the prefecture, a location in the high mountains and one at sea level. The parasite-free animal had gastric lesions typical of a previous infection with this parasite. The nematode intensities were higher in wild boars caught in the mountains than those caught at near sea level. The extremely high prevalence of *G. doloresi* in wild boars indicates that the life cycle of this parasite is well established in this area of Japan.

Key words: *Gnathostoma doloresi*, prevalence, survey, *Sus scrofa leucomystax*, wild boars.

Gnathostoma doloresi is a nematode parasite distributed throughout southeastern Asia which parasitizes the stomach wall of domestic pigs and wild boars (*Sus scrofa leucomystax*) causing ulcerative and/or granulomatous lesions (Miyazaki, 1960). The first intermediate hosts are copepods and the second intermediate/paratenic hosts are various species of freshwater fishes, salamanders, frogs, toads, snakes, mice and rats (Daengsvang, 1980; Nawa et al., 1993). In Japan, *G. doloresi* occurs where there are wild boars, their main definitive host. There is an especially high prevalence in Kyushu and Shikoku Districts of Japan (Miyazaki, 1960). Since Ogata et al. (1988; Nawa et al., 1989) first found human cases of cutaneous larva migrans due to *G. doloresi* infection in Miyazaki Prefecture, this parasite is known as one of the important food-borne zoonoses. Infection of humans occurs by ingestion of uncooked/under-

cooked meat of infected second intermediate/paratenic hosts. Because patients with gnathostomiasis have been found almost every year in the vicinity of Miyazaki Prefecture (Maruyama et al., 1996), we determined the prevalence of *G. doloresi* in wild boars captured in this area.

Thirty-two wild boars captured in two localities in Miyazaki Prefecture during November and December 1996 were examined. Twelve animals were caught at Shiromi (32°16'N, 131°15'E) located in a high mountainous area, and 20 were captured in Kushima (31°24'N, 131°20'E), an open area with a few low hills near the sea.

The stomachs of wild boars were incised along the greater curvature and nematodes attached to the gastric mucosa were carefully removed with forceps. Gastric contents were diluted in a large volume of water and strained through several layers of stainless steel mesh of different pore sizes to collect free nematodes. All specimens were identified as *G. doloresi* under a dissecting microscope according to their characteristic morphology of cuticular spines covering the whole body surface, swollen posterior body with irregularly folded cuticle, and eggs with two caps at each end of the shells. Body length of nematodes were measured from 11 animals. Representative specimens of *G. doloresi* are deposited in the Department of Parasitology (Miyazaki Medical College, Miyazaki, Japan; accession numbers GdA 961217, 961219, 961225, 961227, 970113).

All statistical analyses employed Stat-View SE+Graphics (Abacus Concepts, Inc., Berkeley, California, USA). The nematode intensities of the hosts in two different locations were compared by the Mann-Whitney *U* test. Significance was accepted

TABLE 1. *Gnathostoma doloresi* infection in wild boars from Miyazaki, Japan.

	Shiromi	Kushima	Total
Number of animals	12	20	32
Sex (Male/Female)	7/3 ^a	6/14	13/17 ^a
Body weight (Mean \pm SE)	58.5 \pm 6.8 kg ^a	51.9 \pm 3.6 kg	54.1 \pm 3.3 kg ^a
Range	40–100 kg	15–80 kg	15–100 kg
Prevalence (%)	12/12 (100)	19/20 (95)	31/32 (97)
Intensity (Mean \pm SE)	57.1 \pm 11.5 ^b	20.1 \pm 4.7 ^b	34.4 \pm 6.2
Range	16–148	5–91	5–148

^a Not identified for 2 animals.^b Significantly different from each other by Mann-Whitney's *U* test ($P < 0.01$).

when $P \leq 0.05$. Correlation between the nematode intensities and body weights of hosts or the average length of nematodes in individual hosts were examined using a simple regression test.

Twelve and 20 wild boars were from Shiromi and Kushima, respectively. An average body weight of animals ($n = 30$) was 54.1 kg ranging from 15 to 100 kg, and the sex ratio was male: female = 1:1.3. Of 32 wild boars examined, 31 (97%) were infected with *G. doloresi*. One uninfected animal had many burrows in the gastric wall indicating that it also had been infected previously. The mean intensity (\pm SE) was 34.4 ± 6.2 (range = 5–148). When the nematode intensity of the hosts in two different places were compared, that in Shiromi was 57.1 ± 11.5 , whereas that in Kushima was 20.1 ± 4.7 , and they were significantly different from each other ($P < 0.01$ by Mann-Whitney's *U* test).

The nematode intensities were not correlated with body weights of the hosts (correlation coefficient $r = 0.366$), suggesting that the parasites do not accumulate along with the age of hosts. To determine whether crowding effect is observed in *G. doloresi* infection in wild boars, length of individual nematodes was measured for those obtained from 11 animals. The average length of nematodes in individual hosts was not correlated to the nematode intensity ($r = 0.033$), indicating no such a effect in this study.

Most wild boars in Miyazaki Prefecture were infected with *G. doloresi* regardless

of geographic locality. A survey conducted in Shiromi, 9 yr previously (Nawa and Imai, 1989) also indicated a high prevalence (70%) of *G. doloresi* in wild boars. High prevalence ($>50\%$) of *G. doloresi* in wild boars in the mountainous areas of Shikoku and Kyushu Districts was reported by earlier studies (Miyazaki et al., 1953; Isobe, 1956; Miyazaki, 1960; Sakaguchi et al., 1985). In contrast, prevalence of *G. doloresi* in pigs from other southeastern Asian countries is $<10\%$ (Daengsvang, 1980). Because the life cycle of *G. doloresi* requires two or more intermediate/paratenic hosts depending on predator/prey relationships, ecosystems like those of Kyushu District may be exceptionally suitable for this species.

Although practically all wild boars were infected with *G. doloresi*, we noted a difference in intensities in hosts caught at Shiromi and Kushima. Shiromi is located in a mountainous area, 500 to 1,000 m above sea level, whereas Kushima is located in an open plain with a few low hills near the sea. The intermediate/paratenic hosts for *G. doloresi* may be distributed more densely in the mountainous areas. Conversely, wild boars living in mountainous areas may ingest more animal foods than those living in the plains because of the difference in the relative availability of animal and plant foods in two locations. This needs further clarification.

Nawa and Imai (1989) reported that wild boars of about 20 kg body weight had the highest nematode intensities com-

pared to larger or smaller animals. However, no such correlation was observed in the present study. This may simply be due to the number of animals examined. We examined over 30 animals in the present study, whereas only 10 animals were examined in the previous study. Adult nematodes do not accumulate in wild boars with time, suggesting that their survival may be fairly short and infection is continuously occurring. Alternatively, once hosts were infected with *G. doloresi*, they might become resistant to further infection. This also needs further clarification.

This work is supported in part by the Grant-in-Aid (#08041187) for International Scientific Research (Field Research) from the Ministry of Education, Culture, Sports and Science of Japan. The authors wish to thank K. Otsuka, S. Hamasuna, T. Kuramoto, H. Miyagoe, and T. Iwamitsu, for their enthusiastic collaboration in collecting wild boars.

LITERATURE CITED

- DAENGSVANG, S. 1980. *Gnathostoma doloresi* Tubangui, 1925. In A Monograph on the Genus *Gnathostoma* & Gnathostomiasis in Thailand. Southeast Asian Medical Information Center, Tokyo, Japan, pp. 61–70.
- ISOBE, C. 1956. Studies on *Gnathostoma* in Kumamoto Prefecture. Journal of Kumamoto Medical Society 30: 1183–1201.
- MARUYAMA, H., S. NODA, AND Y. NAWA. 1996. Emerging problems of parasitic diseases in southern Kyushu, Japan. Japanese Journal of Parasitology 45: 192–200.
- MIYAZAKI, I. 1960. On the genus *Gnathostoma* and human gnathostomiasis, with special reference to Japan. Experimental Parasitology 9: 338–370.
- , Y. ISHII, AND C. ISOBE. 1953. Incidence of infection among wild boars with *Gnathostoma doloresi* Tubangui. Igaku to Seibutsugaku (Medical Biology) 29: 65–68.
- NAWA, Y., AND J. IMAI. 1989. Current status of *Gnathostoma doloresi* infection in wild boars captured in Miyazaki Prefecture, Japan. Japanese Journal of Parasitology 38: 385–387.
- , ———, Y. HORII, K. OGATA, AND K. OTSUKA. 1993. *Gnathostoma doloresi* larvae found in *Lepomis macrochirus* RAFINESQUE, a freshwater fish (common name: blue-gill), captured in the central part of Miyazaki Prefecture, Japan. Japanese Journal of Parasitology 42: 40–43.
- , ———, K. OGATA, AND K. OTSUKA. 1989. The first record of a confirmed human case of *Gnathostoma doloresi* infection. The Journal of Parasitology 75: 166–169.
- OGATA, K., J. IMAI, AND Y. NAWA. 1988. Three confirmed and five suspected human cases of *Gnathostoma doloresi* infection found in Miyazaki Prefecture, Kyushu. Japanese Journal of Parasitology 37: 358–364.
- SAKAGUCHI, Y., T. MIMORI, H. HIRAI, M. KORENAGA, AND I. TADA. 1985. *Gnathostoma doloresi* infection in wild boars captured in Kumamoto Prefecture, Japan. Kumamoto Medical Journal 38: 147–152.

Received for publication 21 February 1997.