



Alternative Performance Targets for Integrating Cover Crops as a Proactive Herbicide-Resistance Management Tool

Authors: Bunchek, Jess M., Wallace, John M., Curran, William S., Mortensen, David A., VanGessel, Mark J., et al.

Source: Weed Science, 68(5) : 534-544

Published By: Weed Science Society of America

URL: <https://doi.org/10.1017/wsc.2020.49>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Research Article

Cite this article: Bunchek JM, Wallace JM, Curran WS, Mortensen DA, VanGessel MJ, Scott BA (2020) Alternative performance targets for integrating cover crops as a proactive herbicide-resistance management tool. *Weed Sci.* **68**: 534–544. doi: [10.1017/wsc.2020.49](https://doi.org/10.1017/wsc.2020.49)

Received: 17 February 2020
Revised: 21 May 2020
Accepted: 10 June 2020
First published online: 16 June 2020

Associate Editor:

Timothy L. Grey, University of Georgia

Keywords:

Erigeron canadensis; integrated weed management; no-till; population dynamics

Author for correspondence:

John M. Wallace, 116 ASI Building, University Park, PA 16802. (Email: jmw309@psu.edu)

Alternative performance targets for integrating cover crops as a proactive herbicide-resistance management tool

Jess M. Bunchek¹, John M. Wallace² , William S. Curran³, David A. Mortensen⁴, Mark J. VanGessel⁵ and Barbara A. Scott⁶

¹Graduate Research Assistant, Pennsylvania State University, University Park, PA, USA; ²Assistant Professor, Pennsylvania State University, University Park, PA, USA; ³Professor Emeritus, Pennsylvania State University, University Park, PA, USA; ⁴Professor, University of New Hampshire, Durham, NH, USA; ⁵Professor, University of Delaware, Georgetown, DE, USA and ⁶Associate Scientist, University of Delaware, Georgetown, DE, USA

Abstract

Intensified cover-cropping practices are increasingly viewed as a herbicide-resistance management tool but clear distinction between reactive and proactive resistance management performance targets is needed. We evaluated two proactive performance targets for integrating cover-cropping tactics, including (1) facilitation of reduced herbicide inputs and (2) reduced herbicide selection pressure. We conducted corn (*Zea mays* L.) and soybean [*Glycine max* (L.) Merr.] field experiments in Pennsylvania and Delaware using synthetic weed seedbanks of horseweed [*Conyza canadensis* (L.) Cronquist] and smooth pigweed (*Amaranthus hybridus* L.) to assess winter and summer annual population dynamics, respectively. The effect of alternative cover crops was evaluated across a range of herbicide inputs. Cover crop biomass production ranged from 2,000 to 8,500 kg ha⁻¹ in corn and 3,000 to 5,500 kg ha⁻¹ in soybean. Experimental results demonstrated that herbicide-based tactics were the primary drivers of total weed biomass production, with cover-cropping tactics providing an additive weed-suppression benefit. Substitution of cover crops for PRE or POST herbicide programs did not reduce total weed control levels or cash crop yields but did result in lower net returns due to higher input costs. Cover-cropping tactics significantly reduced *C. canadensis* populations in three of four cover crop treatments and decreased the number of large rosettes (>7.6-cm diameter) at the time of preplant herbicide exposure. Substitution of cover crops for PRE herbicides resulted in increased selection pressure on POST herbicides, but reduced the number of large individuals (>10 cm) at POST applications. Collectively, our findings suggest that cover crops can reduce the intensity of selection pressure on POST herbicides, but the magnitude of the effect varies based on weed life-history traits. Additional work is needed to describe proactive resistance management concepts and performance targets for integrating cover crops so producers can apply these concepts in site-specific, within-field management practices.

Introduction

The evolution and spread of glyphosate- and multiple herbicide-resistant weed species in conservation tillage systems has led to increased use of new herbicide-tolerant soybean [*Glycine max* (L.) Merr.] varieties and associated POST herbicides (dicamba, 2,4-D, glufosinate), as well as increased use of soil-applied herbicides (Kniss 2018; Norsworthy et al. 2012). Herbicide-resistance models indicate that applying herbicide mixtures containing multiple sites of action within crop phases and rotating sites of action across crop rotations are imperative for delaying further resistance evolution (Busi et al. 2018; Evans et al. 2018). In addition, integrated weed management (IWM) strategies that employ complementary chemical and nonchemical weed management tactics underpinned by ecological principles (Liebman et al. 2001) are critically needed for the long-term sustainability of weed management systems and preservation of currently effective herbicides (Mortensen et al. 2012).

Diversity is a foundation of both IWM and sustainable agroecosystem management. Crop rotation diversity can reduce weed populations and selection pressures that drive the evolution of herbicide resistance (Weisberger et al. 2019). Integrating fall-sown cover crops within annual grain rotations introduces additional crop diversity, extends the temporal period when a competitive crop is growing in a field, and is increasingly viewed as a best management practice for soil conservation (Hamilton et al. 2017; Wayman et al. 2016). At a mechanistic level, cover crops serve as a weed community assembly filter by competing with weeds for light, water, space, and nutrients (Smith et al. 2015). Fall-sown cover crops provide direct competition with weed species that have overlapping life cycles, and cover crop residue left on the soil surface after

© Weed Science Society of America, 2020. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.



termination can indirectly influence the recruitment and growth rates of summer annual weed species (Ryan et al. 2011b; Teasdale et al. 2005). Recent studies have demonstrated that integrating cover crops can increase suppression of glyphosate-resistant weeds, including fall- and spring-emerging horseweed [*Coryza canadensis* (L.) Cronquist] (Cholette et al. 2018; Pittman et al. 2019; Wallace et al. 2019) and summer annuals Palmer amaranth (*Amaranthus palmeri* S. Watson) and waterhemp [*Amaranthus tuberculatus* (Moq.) Sauer] (Loux et al. 2017; Montgomery et al. 2018; Wiggins et al. 2015).

Intensifying cover crop management via use of weed-suppressive monocultures or mixtures (Baraibar et al. 2018; Smith et al. 2015; Wortman et al. 2012), higher seeding rates (Ryan et al. 2011a), or delayed termination tactics (Nord et al. 2012; Wallace et al. 2019) will likely make cover cropping a more robust IWM tool. However, there are currently few guiding IWM principles for integrating cover crops with herbicide-based weed control tactics. We suggest that the lack of guidelines result, in part, from alternative and sometimes opposing management goals. Weed control programs have traditionally relied on a critical weed-free period management goal, but a zero seed rain management goal is increasingly recommended for multiple herbicide-resistant weeds in no-till systems with simplified crop rotations (Norsworthy et al. 2012). The contribution of cover crops to zero seed rain management of multiple herbicide-resistant weeds remains equivocal. For example, previous studies have concluded that multiple-pass and multiple site-of-action weed control programs are still needed, regardless of cover crop selection and management, to control multiple herbicide-resistant species such as *A. palmeri* (Loux et al. 2017). Consequently, when viewed as a prescriptive tool for achieving a zero seed rain management goal, integrating cover crops may be perceived by growers as contributing to additional weed control input costs, agronomic management complexity, or reduced yields due to incomplete termination of cover crops or unanticipated trade-offs related to soil fertility, crop establishment, and pest dynamics.

In contrast, cover crops are a promising tool for proactive resistance management, where additional management tactics are implemented before resistance is widespread in order to delay the evolution of resistance traits. The contribution of cover crops to proactive resistance management should be measured with different performance targets, including a measurable decrease in the intensity of selection pressure on herbicide-based weed control tactics and reduction in the number of herbicide applications or active ingredients in weed control programs. Increasing cover crop management intensity has been shown to decrease the intensity of herbicide selection pressure by reducing emerged weed population densities and the number of large individuals within the weed population at the time of herbicide exposure (Montgomery et al. 2018; Wallace et al. 2019; Wiggins et al. 2015). These two population-level outcomes can potentially delay the evolution of herbicide resistance by decreasing the proportion of the seedbank that is exposed to the selection pressure of a given herbicide (Comont et al. 2019) and by decreasing the survival rate, and therefore fitness advantage, of larger individuals that have a disproportionate effect on the gene pool of subsequent generations (Heywood 1986; Weiner 1985).

We conducted field experiments at two mid-Atlantic locations to evaluate high-residue cover crop management across a herbicide input gradient in no-till corn (*Zea mays* L.) and soybean production systems. Field experiments were designed to determine whether high-residue cover crop management: (1) facilitates reduced

herbicide inputs without compromising cash crop and weed control performance targets and (2) decreases the intensity of selection pressure at preplant burndown and POST herbicide application timings by reducing the density and the proportion of large individuals at the time of herbicide exposure. We evaluated weed population-level responses using artificial seedbanks of glyphosate-resistant *C. canadensis* and glyphosate-susceptible *A. hybridus*. Consequently, this study is best viewed as evaluation of proactive resistance management designed to slow future herbicide-resistance evolution by implementing IWM, rather than reactive management approaches for current, multiple herbicide-resistant weed species. Finally, we conducted a partial budget analysis to characterize the effects of cover crop-based IWM strategies on short-term profits.

Materials and Methods

Corn and soybean experiments were conducted from fall 2015 to 2017 at the University of Delaware's Carvel Research and Education Center near Georgetown, DE, and the Penn State Russell E. Larson Agricultural Experiment Center near Rock Springs, PA. The DE sites were on Pepperbox loamy sand (loamy, mixed, semiactive, mesic Aquic Arenic Paleudults) soils and the PA sites were on Hagerstown silt loam (fine, mixed, semiactive, mesic Typic Hapludalfs) soils. Average annual growing degree days (GDD) are higher at Georgetown, DE (2,440 GDD) than Rock Springs, PA (1,660 GDD), but annual precipitation is similar between locations, averaging (30-yr) between 100 and 115 cm (NOAACD 2019). The 2016 and 2017 growing season (April to September) approximated 30-yr averages at Rock Springs, PA, for both precipitation and GDDs. At Georgetown, DE, the only departure from 30-yr averages included lower accumulated precipitation in August of the 2016 growing season. Field experiments in corn were preceded by small grain production the previous year at each site, and soybean experiments were preceded by corn the previous year at each site (PA, corn silage; DE, corn grain).

Experimental Design and Field Operations

The interaction between alternative cover-cropping and herbicide control tactics was evaluated using a two-factor, randomized complete block with a split-plot design and four replications. Cover crop treatments were imposed in main plots (PA, 12 by 12 m; DE, 12 by 8 m) and herbicide treatments were imposed in split plots (PA, 3 by 12 m; DE, 3 by 8 m). In the corn experiment, cover crop treatments included: (1) a cover crop control, (2) cereal rye (*Secale cereale* L.) 'Aroostook' + crimson clover (*Trifolium incarnatum* L.) 'Dixie' seeded at 34 + 22 kg ha⁻¹, and (3) cereal rye + hairy vetch (*Vicia villosa* Roth) 'Auburn Early Cover' seeded at 34 + 34 kg ha⁻¹. In the soybean experiment, cover crop treatments included: (1) a cover crop control, (2) cereal rye seeded at 135 kg ha⁻¹, and (3) cereal rye + hairy vetch seeded at 101 + 22 kg ha⁻¹. Seeding rate selection was informed by cover crop management practices used in organic rotational no-till systems for optimizing weed suppression and nitrogen provisioning, with grass-dominant monocultures or mixtures preceding soybean and mixtures designed to increase legume proportion preceding corn (Wallace et al. 2017). Cover crops were seeded with a no-till drill in 19-cm-wide rows following a burndown glyphosate application (1.26 kg ae ha⁻¹) after small grain harvest (early September) in the corn experiment and after corn harvest (early to mid-October) in the soybean experiment (Table 1).

Table 1. Dates of field operations in corn and soybean experiment by site (DE, PA) and year (16, 17).^a

Field operation ^b	Corn experiment				Soybean experiment			
	DE16	DE17	PA16	PA17	DE16	DE17	PA16	PA17
Fall cover crop seeding	09/17	09/15	09/01	09/01	10/14	10/17	10/01	10/06
Preplant herbicide application	05/05	05/03	05/06	05/04	05/20	05/26	05/11	05/09
Cash crop planting	05/16	05/15	05/17	06/08	06/01	06/02	05/23	05/19
PRE herbicide application	05/17	05/13	05/17	06/09	06/02	06/02	05/23	05/19
POST herbicide application	06/29	06/28	06/16	07/06	07/08	06/28	06/27	07/03
Cash crop harvest	10/02	09/28	11/07	11/04	10/27	10/02	11/01	11/15

^aDE, University of Delaware's Carvel Research and Education Center near Georgetown, DE; PA, Penn State Russell E. Larson Agricultural Experiment Center near Rock Springs, PA. 16, 2015–2016; 17, 2016–2017.

^bHerbicide treatment combinations include substitutive (PRE only, POST only) and additive (PRE/POST) integrated weed management (IWM) strategies.

Table 2. Herbicide active ingredients and application rates used in corn and soybean herbicide treatments.

Strategy ^a	Corn		Soybean	
	Active ingredient	Application rate	Active ingredient	Application rate
PrePlant	Glyphosate + 2,4-D ester	1.26 kg ae ha ⁻¹ 0.56 kg ae ha ⁻¹	Glyphosate 2,4-D ester	1.26 kg ae ha ⁻¹ 0.56 kg ae ha ⁻¹
PRE	S-metolachlor + mesotrione	1.7 kg ai ha ⁻¹ 0.18 kg ai ha ⁻¹	S-metolachlor + flumetsulam	1.7 kg ai ha ⁻¹ 56 g ai ha ⁻¹
POST	Glufosinate + ammonium sulfate	0.59 kg ai ha ⁻¹ 3.36 kg ha ⁻¹	Glufosinate + ammonium sulfate	0.59 kg ai ha ⁻¹ 3.36 kg ha ⁻¹

^aHerbicide application timing for each strategy is presented in Table 1.

In each experiment, an artificial *C. canadensis* and *A. hybridus* seedbank was established in separate, permanently marked 0.5-m² microplots (0.5 by 1.0 m) in the middle of each split-plot treatment immediately after cover crop seeding each year. Mature *C. canadensis* and *A. hybridus* inflorescences were collected each season from local populations at each experimental site. Seeds were separated from inflorescences using a series of mesh screens in the laboratory, composited in a single seed lot, and dry stored at room temperature. Seeding rates were based on weight aliquots for approximately 5,000 *C. canadensis* seeds m⁻² and 500 *A. hybridus* seeds m⁻². Glyphosate resistance was not confirmed in *C. canadensis* seed lots, but *C. canadensis* biotypes were known to be resistant at both sites based on previous management experience.

Herbicide treatments in corn and soybean experiments included: (1) a preplant burndown application (PrePlant; herbicide control) timed to the late-boot growth stage (Zadoks 55) of cereal rye 10 d before cash crop planting; (2) PrePlant followed by (fb) a soil-applied PRE program (PRE) applied at cash crop planting; (3) PrePlant fb a POST program (POST) applied at the V3-V4 crop growth stage, approximately 35 to 42 d after cash crop planting; and (4) a three-pass herbicide program including the PrePlant, PRE, and POST applications (PRE/POST). Active ingredients used for herbicide treatments differed between corn and soybean experiments (Table 2). Herbicides were applied at 187 L ha⁻¹ at 276 kPa using a tractor-mounted sprayer equipped with AI11002VS spray tips (TeeJet® Technologies, Glendale Heights, IL).

Corn and soybean were planted in 76-cm-wide rows with a John Deere 1720 no-till planter (Deere & Company, Moline, IL) equipped with ZRX roller-crimpers and double-disk row cleaners (Dawn Equipment, Sycamore, IL) at the PA site. Integrated ZRX systems are designed to roll-crimp cover crops at planting with use of double-disk row cleaners that move residue away from the planted row to improve seed-to-soil contact. However, the amount of soil and cover crop mulch disturbance created by these row cleaners varies based on soil moisture conditions at the time of planting. Corn and soybean were planted in 76-cm-wide rows with

a Kinze 3000 planter (Kinze Manufacturing, Williamsburg, IA) at the DE site. Glufosinate-resistant corn 'DEKALB DKC48-56RIB' (DEKALB Genetics Corporation, DeKalb, IL 60115) was planted at 79,000 seeds ha⁻¹, with 45 kg ha⁻¹ N applied at planting. Glufosinate-resistant soybean 'Doebler's DB3217LL' (Doebler's Pennsylvania Hybrids, Williamsport, PA) was planted at 445,000 seeds ha⁻¹.

Data Collection

Cover crop performance was assessed by collecting living above-ground biomass (kg ha⁻¹) in two randomly placed 0.25-m² microplots at the split-plot level just before termination in the spring. Aboveground biomass subsamples were composited and separated by cover crop species, oven-dried (65 C) for 5 d, and weighed. Cash crop yield performance was assessed by machine harvesting the middle two rows from each split plot using a small plot combine. Corn and soybean yields were corrected to 15.5% and 13% moisture content, respectively, for statistical analyses.

Treatment effects on weed population dynamics were assessed using multiple indicators. First, *C. canadensis* and *A. hybridus* population densities were quantified within one-half of each microplot (0.25-m²) just before PrePlant and POST applications, respectively. Second, *C. canadensis* populations were subsampled at the PA site at the time of PrePlant applications to quantify treatment effects on the rosette size structure of the emerged populations. The diameter of the first 10 seedlings or rosettes encountered within the two middle cover crop interrows, from the front to back of the microplots, was measured to the nearest 0.5 cm. In plots with low *C. canadensis* density, plants were measured in the entire microplot or until 10 plants had been assessed. Similarly, *A. hybridus* populations were subsampled at the PA site at the time of POST application. The height of the first 10 *A. hybridus* plants was measured to the nearest 1 cm in cover crop interrows. Finally, aboveground weed biomass, including the resident emerged weed community, was collected in late August from each microplot at

both sites. Subsamples were separated by species, oven-dried (65 C) for 5 d, and weighed.

Statistical Analysis

Linear mixed-effect (LME) models were used to conduct null hypothesis tests of cover crop and herbicide treatment effects on cover crop biomass, grain yield, and weed biomass response variables in R using the NLME package (Pinheiro et al. 2015). Analyses were conducted by cash crop (corn and soybean) due to differences in cover crop and herbicide treatments. All models were fit with a nested random effects structure (year/site/block) to account for sources of variation associated with environmental and management differences (site, year) as well as spatial variation within site-years (block).

Fixed-effect parameters differed among LME models based on the observation timing of response variables. Total aboveground cover crop biomass (kg ha^{-1}) was averaged at the main plot level, because split-plot treatments (herbicide) had yet to be imposed. Cash crop grain yield (kg ha^{-1}) and total aboveground weed biomass (kg ha^{-1}) models were fit with cover crop, herbicide, and their interaction as fixed effects. Total aboveground weed biomass and cash crop yields were log transformed after adding a constant (1.0) to meet assumptions of normality and homogeneity of variance. Statistical significance of fixed-effect terms was assessed by conditional *F*-tests. Fitted models were used to estimate marginal means (\pm SE) for each response variable using the EMMEANS package (Lenth et al. 2018).

Population density (plants 0.25 m^{-2}) of *C. canadensis* and *A. hybridus* was assessed with generalized linear mixed-effects models (GLMM) using a negative binomial distribution and log link function (*glmer.nb*) in the LME4 package (Bates et al. 2016). Data from DE in 2015 to 2016 were excluded from this analysis due to low recruitment within synthetic weed seedbanks, leaving a random effects structure of block nested within site-year and three site-year treatment levels. Population density of *C. canadensis* at the preplant application timing was averaged at the main plot level, because split-plot treatments (herbicide) had yet to be imposed. Models of population density of *C. canadensis* and *A. hybridus* at the POST timing were fit with cover crop, herbicide, and their interaction as fixed effects. The herbicide factor was reduced to two treatment levels by pooling treatments with or without PRE applications to compare PRE and PrePlant-only treatment effects. Statistical significance of fixed effects and their interaction terms were assessed using log-likelihood ratio tests (Wald χ^2) in GLMMs (Bolker et al. 2008).

Mean separations of significant effects in both LME and GLMM models were conducted using Tukey's contrasts (*glht*) in the package MULTCOMP (Hothorn et al. 2008). In addition, marginal and conditional coefficients of determination (R^2_m and R^2_c , respectively) were calculated to describe the proportion of the variance in the response associated with fixed effects only (R^2_m) and random plus fixed-effect components (R^2_c) of the model using the package MuMIN (Nakagawa and Schielzeth 2013).

A limited data set resulting from variable recruitment levels across site-years precluded statistical analysis of *C. canadensis* and *A. hybridus* population size structure as described in Wallace et al. (2019). Frequency distribution plots were constructed to visually evaluate cover crop effects on weed size structure within populations. The frequency distribution of *C. canadensis* rosette size at the time of the PrePlant application was calculated at the main plot level and pooled across years. The frequency distribution of *A. hybridus* height

at POST application was calculated within split plots with and without PRE applications to isolate the effect of cover crops and then pooled across years.

Preplanned contrasts from fitted models and a partial budget analysis were utilized to better understand the effects of alternative IWM tactics that integrate cover crops on weed management and economic outcomes. In each set of analyses, pairwise contrasts were conducted between the PRE/POST treatment without cover crops, representing a "standard" herbicide-based weed management strategy, and additive (i.e., adding cover crops to herbicide-based program) or substitutive (i.e., substituting cover crops for either PRE or POST herbicide inputs) IWM strategies. Fitted models of (1) total weed biomass (August), (2) *C. canadensis* density at the PrePlant timing, and (3) *A. hybridus* density at the POST timing were used to estimate the marginal means of each treatment. We then calculated pairwise multiple comparisons using the Bonferroni correction for an overall error rate of 0.05 in the EMMEANS package. Pairwise comparisons of weed density data analyzed with a Poisson distribution are presented as the ratio of geometric (i.e., back-transformed) means and can be interpreted as the proportional difference between treatments. Weed biomass density is presented as the difference in percent weed control by expressing treatment differences as a proportion of the weedy check (PrePlant only, no cover crop). Finally, a partial budget analysis was conducted to compare net returns to management (revenues – input costs) for alternative IWM approaches. Revenues (yield \times price) were calculated using experimental yields (corn, 15.5% moisture; soybean, 13% moisture) and the national average 2016 to 2017 corn and soybean market prices: $\$0.13 \text{ kg}^{-1}$ and $\$0.34 \text{ kg}^{-1}$, respectively (USDA-NASS 2019). Input costs that differed among IWM approaches and were used in analyses included chemicals, cover crop seed, and field operation passes (Supplementary Table 1). Chemical and cover crop seed costs were obtained from mid-Atlantic distributors (Helena Chemical, Warriors Mark, PA 16877; King's Agriseeds, Lancaster, PA), and field operation costs were taken from 2016 Penn State University crop enterprise budgets (Harper 2016).

Results and Discussion

Cover Crop Performance

Mean total aboveground cover crop biomass did not differ ($P < 0.05$) between cover crop treatments in the corn and soybean experiments (Figure 1). Random effects associated with management and environmental conditions across sites and years contributed 80% and 50% of the total variation observed in total aboveground biomass production in corn and soybean experiments, respectively. Mean cover crop biomass ranged from 2,000 to 8,500 kg ha^{-1} across sites and years in the corn experiment and 3,000 to 5,500 kg ha^{-1} in the soybean experiment. For comparison, targeted weed-suppressive biomass levels for organic grain production, which is characteristically more management intensive, range from 5,000 to 9,000 kg ha^{-1} (Mirsky et al. 2012; Wallace et al. 2017). Observed variation in cover crop biomass production and the grass-legume ratio highlight how environmental and soil factors, rather than management practices such as cover crop species selection and seeding rates, drive biomass accumulation.

Weed Control and Yield Performance Targets

In the corn experiment, total aboveground weed biomass (August) showed a cover crop by herbicide treatment effect ($F(6, 165) = 2.5$,




Figure 1. Mean cover crop biomass (kg ha^{-1}) by site-year and cover crop treatment in (A) corn and (B) soybean experiments. Data (mean \pm SE) are presented as total cover crop biomass and by species in grass/legume mixtures. Cover crop treatments were not significantly different ($P > 0.05$) within corn and soybean experiments. See Table 1 for definitions of site-year abbreviations.

$P = 0.02$). Cover crops had no effect on total aboveground weed biomass in PRE, POST, and PRE/POST herbicide strategies, but rye/clover reduced weed biomass compared with no cover crop in the PrePlant-only treatment (Figure 2A). Within each cover crop treatment level, weed biomass was lower in PRE, POST, and PRE/POST treatments compared with PrePlant only. Pairwise comparisons between PRE, POST, and PRE/POST treatments within cover crop treatments did not differ, with the exception of rye/vetch, for which weed biomass was lower in the PRE/POST strategy compared with POST only. Weed management treatments explained 51% of the total variation observed in total aboveground weed biomass, whereas random effects associated with environmental and management conditions contributed 10% of total variation. Herbicide and cover crop treatments did not affect ($P > 0.05$) corn yield (Figure 2B). Corn yields approximated county-level averages, with exception of the 2016 experiment in DE, where drought conditions likely contributed to below-average corn yields.

In the soybean experiment, herbicide ($F(3, 164) = 90.1$, $P < 0.0001$) and cover crop ($F(2, 164) = 8.8$, $P = 0.0002$) treatments resulted in additive effects on aboveground weed biomass (August; Figure 2C). Each herbicide program (PRE, POST, PRE/POST) resulted in lower weed biomass than PrePlant only, and herbicide treatments with POST applications resulted in lower weed biomass than the PRE treatment. Weed biomass was lower in cover crop treatments compared with the control across herbicide strategy levels. Weed management treatments explained 44% of the total variation observed in total aboveground weed biomass, whereas random effects associated with environmental and management conditions contributed 27% of total variation. Herbicide treatments had a significant effect ($F(3,164) = 26.4$, $P < 0.0001$) on soybean yield (Figure 2D), with lower yields observed in the no-herbicide control compared with other herbicide programs (PRE, POST, PRE/POST). Soybean yields approximated county-level averages, with exception of the 2016 experiment in DE, where drought conditions likely contributed to below-average corn yields.

In summary, these results indicate that herbicide-based tactics were the primary drivers of weed suppression, and cover crops generally provided additional weed suppression that was smaller

in magnitude. Our results do not provide supporting evidence that surface residue mulch reduced herbicide efficacy or cash crop yields. Recent studies in corn and soybean production regions of the Midwest and Midsouth support the inference that herbicide inputs are the primary driver of cash crop yields and weed control performance, including multiple herbicide-resistant *Amaranthus* spp., when cover crops are integrated as a component of an IWM strategy (Loux et al. 2017; Montgomery et al. 2018; Wiggins et al. 2015).

Herbicide Selection Pressure on Weed Populations

In the corn experiment, cover crop main effects were significant ($\chi^2 = 29.5$, $P < 0.001$) for *C. canadensis* density at the PrePlant application timing. Both cover crop treatments reduced *C. canadensis* density compared with the control (Figure 3A). High levels of *C. canadensis* control efficacy were observed following the PrePlant application across treatments, which precluded reliable model fit of *C. canadensis* density at the POST application timing. A significant herbicide main effect ($\chi^2 = 46.2$, $P < 0.001$) was observed in analysis of *A. hybridus* density at the POST application timing, in which PRE residual herbicides reduced *A. hybridus* density compared with no PRE herbicides across cover crop treatments (Figure 3B). In PA experiments, 18% of *C. canadensis* rosettes exposed to PrePlant applications in the cover crop control were larger than 7.6 cm in diameter (Figure 3C), which is associated with decreased efficacy of glyphosate and auxinic herbicides on *C. canadensis* in spring preplant applications (Dinelli et al. 2006). Similarly, 3% of *A. hybridus* plants exposed to POST applications in the cover crop control exceeded 10 cm in height (Figure 3D), which is the maximum weed height commonly recommended on herbicide labels and a best management practice for maintaining herbicide efficacy. In comparison, no *C. canadensis* rosettes or *A. hybridus* plants exceeded a 5-cm diameter or 8-cm height, respectively, in treatments with cover crops.

In the soybean experiment, cover crop main effects were significant ($\chi^2 = 11.2$, $P < 0.01$) for *C. canadensis* density at the PrePlant timing. Rye/vetch reduced *C. canadensis* density compared with the control and rye only, which did not differ in pairwise




Figure 2. Mean total aboveground weed biomass (A and C) and cash crop yields (B and D) by herbicide and cover crop treatment in corn and soybean experiments. Weed biomass was collected in late August and included seeded weed species (*Conyza canadensis*, ERICA; *Amaranthus hybridus*; AMACH) and the resident weed community. Data are estimated population-level log-transformed means (\pm SE) of fixed effects. See Table 1 for definitions of site-year abbreviations.

comparisons (Figure 4A). High levels of *C. canadensis* control were also observed following PrePlant application in soybean. At the POST application timing, a significant herbicide main effect ($\chi^2 = 26.7$, $P < 0.001$) was observed, in which *C. canadensis* density was lower when herbicide strategies included a PRE application. Analysis of *A. hybridus* density at the POST application timing showed a cover crop by herbicide effect ($\chi^2 = 9.5$, $P < 0.01$), in which the PRE application resulted in lower *A. hybridus* density within the cover crop control but herbicide treatment differences were not observed within rye and rye/vetch treatments (Figure 4B). In PA soybean experiments, 25% of *C. canadensis* rosettes exposed to PrePlant applications in the cover crop control exceeded 7.6 cm in diameter, whereas no *C. canadensis* rosettes exceeded a diameter of 6 cm in treatments with cover crops (Figure 4C). In the cover crop control, 75% of *A. hybridus* plants exposed to POST applications exceeded 10 cm in height, as opposed to 15% in both treatments with cover crops (Figure 4D).

Contrasting Alternative Weed Management Strategies

To provide a deeper understanding of economic and management trade-offs associated with integrating cover crops, we performed multiple contrasts between the “standard” herbicide-based strategy (PRE/POST) without cover crops in our study and alternative IWM-based strategies that integrated cover crops, including (1) an “additive” strategy that aims to enhance weed suppression and diversify selection pressure by integrating cover crops with “standard” herbicide-based strategies and (2) a “substitutive” strategy that aims to reduce herbicide inputs by substituting cover-cropping tactics for PRE or POST herbicides.

Additive and substitutive strategies resulted in increased input costs relative to a herbicide-based strategy in corn (Figure 5) and soybean (Figure 6). Substituting cover-cropping tactics for PRE herbicides resulted in only marginal differences in input costs relative to the herbicide-based strategy when hairy vetch was excluded from the cover crop mixture. Revenues from corn and soybean were not significantly different ($P > 0.05$) in comparisons between IWM and herbicide-based strategies, but mean estimates for IWM strategies were lower than the herbicide-based strategy in 10 of 12 contrasts. Lower net returns to management were observed across all IWM strategies compared with the herbicide-based strategy, which was primarily a function of differences in input costs rather than differences in revenue. We recognize that this partial budget analysis is highly context dependent, but it is instructive for identifying factors that will shape economic decision making and outcomes when proactive IWM strategies are employed. First, we note that our PRE programs were designed for the expected weed community. Presence of multiple-resistant summer annual weeds within cropping systems would necessitate inclusion of more active ingredients and increase herbicide input costs. Next, cover crop seed costs can be highly variable depending on source (i.e., certified variety, bin run), seeding rate, and species mixture. Seeding rates used in this study were adopted from organic no-till practices that rely on cover crop residues as the primary weed-suppression tactic. Integrating cover crops with herbicide-based control tactics likely presents opportunities for reducing cover crop seeding rates and should be explored. Finally, due to widespread farmer interest in soil health and other associated ecosystem services (Schipanski et al. 2014), long-term net returns and nonmarket benefits are increasingly considered




Figure 3. Weed population dynamics in response to herbicide and cover crop treatments in corn experiment, including: (A) *Conyza canadensis* (ERICA) density just before PrePlant and POST herbicide applications; data are treatment means (\pm SE) by cover crop treatment and pooled across site-years. (B) *Amaranthus hybridus* (AMACH) density just before POST application timing in response to cover crop treatment and with or without a PRE application across site-years; horizontal line within box plots indicates median, whiskers extend to $1.5 \times$ interquartile range, and dots indicate outliers. (C) Relative frequency distribution of *C. canadensis* rosette diameter at the PrePlant application timing by cover crop treatment and pooled across Pennsylvania site-years. (D) Relative frequency distribution of *A. hybridus* height just before POST application timing by cover crop in treatments that excluded a PRE application and pooled across Pennsylvania site-years.

in economic decision making related to cover cropping, extending beyond a singular focus on weed management goals.

In our study, total weed control (%) was not significantly different ($P > 0.05$) in comparisons between each additive and substitutive strategy and the herbicide-based strategy in corn (Figure 5) and soybean (Figure 6). The absence of significant differences among strategies for both crop yield and weed control suggests that there are opportunities for reducing herbicide inputs as a proactive resistance management approach in cropping systems that are uninhabited by multiple herbicide-resistant summer annual weed species. However, general trends in our data suggest that substituting cover crops for herbicide inputs have the potential to sustain critical-period weed management goals but compromise zero seed rain management goals. Mean weed control levels were 3% to 15% lower when cover crops were substituted for POST herbicides and 0% to

9% lower when cover crops were substituted for PRE herbicides. In comparison, weed control levels within additive cover crop strategies were 6% lower to 2% higher than herbicide-based strategies.

The effect of IWM strategies on weed population densities at time of herbicide exposure produced similar trends in corn (Figure 5) and soybean (Figure 6) but differed between winter annual (*C. canadensis*) and summer annual (*A. hybridus*) weed species. With exception of the cereal rye monoculture treatment in soybean, cover-cropping tactics significantly reduced *C. canadensis* density at the PrePlant application timing compared with the herbicide-based strategy. In contrast, substituting cover crops for PRE herbicides resulted in higher *A. hybridus* population density ($P < 0.01$), and thus greater herbicide selection pressure at the POST application timing compared with the standard herbicide-based strategy.




Figure 4. Weed population dynamics in response to herbicide and cover crop treatments in soybean experiment, including: (A) *Conyza canadensis* (ERICA) density just before PrePlant and POST herbicide applications; data are treatment means (\pm SE) by cover crop treatment and pooled across site-years. (B) *Amaranthus hybridus* (AMACH) density just before POST application timing in response to cover crop treatment and with or without a PRE application across site-years; horizontal line within box plots indicates median, whiskers extend to $1.5 \times$ interquartile range, and dots indicate outliers. (C) Relative frequency distribution of *C. canadensis* rosette diameter at the PrePlant application timing by cover crop treatment and pooled across Pennsylvania site-years. (D) Relative frequency distribution of *A. hybridus* height just before POST application timing by cover crop in treatments that excluded a PRE application and pooled across Pennsylvania site-years.

Recent studies have consistently demonstrated significant *C. canadensis* population reductions within fall-sown cover crops compared with winter fallow (Cholette et al. 2018; Pittman et al. 2019; Wallace et al. 2019), but the impact of cover crop species and seeding rates on *C. canadensis* recruitment remains equivocal. Wallace et al. (2019) reported that fall cover crop biomass production and foliar ground cover were the most important predictors of spring *C. canadensis* population density, whereas spring biomass production was the primary driver of *C. canadensis* rosette size at the time of preplant herbicide exposure. Though results from our study and others indicate that cover cropping can contribute to herbicide-resistance management by reducing the total population and number of large *C. canadensis* individuals exposed to herbicide selection, additional management challenges may arise. For example, complete suppression of *C. canadensis* by cover crops has

not been observed in published studies, which highlights the need for effective herbicide-based control tactics within fall-sown cover crops. In this management scenario, herbicide applications will need to be optimized to ensure *C. canadensis* individuals surviving within cover crops are exposed to effective herbicide doses, because reduced herbicide deposition through cover crops could result in recurrent selection to low herbicide doses, which has been shown to result in evolution of resistance in certain weed species (Neve and Powles 2005).

Cover-cropping effects on the exposure of *A. hybridus* populations to POST herbicide applications varied in magnitude across our corn and soybean experiments, but results suggest that the presence of surface mulch likely lengthens management windows for on-label POST applications that target small weeds. Wiggins et al. (2015) and Montgomery et al. (2018) showed that




Figure 5. Pairwise comparisons of economic and weed management outcomes between the “standard” herbicide-based strategy (PRE/POST) without cover-cropping and alternative integrated weed management (IWM) strategies that integrate cover crops in corn production. IWM strategies include adding cover crops to herbicide-based strategies (additive) or substituting cover crops for PRE or POST herbicide inputs (substitutive). Input costs reflect differences among strategies in use of cover crop seed, herbicide, labor, and fuel. Pairwise comparisons are presented by cover crop tactic (i.e., species). Significant differences are denoted by asterisks (*** $P < 0.001$; ** $P < 0.01$); input costs and net returns were not subjected to significance tests.




Figure 6. Pairwise comparisons of economic and weed management outcomes between the “standard” herbicide-based strategy (PRE/POST) without cover-cropping and alternative integrated weed management (IWM) strategies that integrate cover crops in soybean production. IWM strategies include adding cover crops to herbicide-based strategies (additive) or substituting cover crops for PRE or POST herbicide inputs (substitutive). Input costs reflect differences among strategies in use of cover crop seed, herbicide, labor, and fuel. Pairwise comparisons are presented by cover crop tactic (i.e., species). Significant differences are denoted by asterisks (** $P < 0.01$); input costs and net returns were not subjected to significance tests.

integrating high-residue cover crops like crimson clover or hairy vetch extended the length of time during which POST applications targeting *A. palmeri* remained on label due to delayed recruitment or reduced growth rates from cover crop mulch interference. Extending management windows for *A. palmeri* is particularly critical given high relative growth rates, but such benefits are likely to extend for many summer annual species that are adapted to emerge with or in a cash crop.

In our study, PRE herbicides resulted in lower *A. hybridus* populations at the time of POST herbicide exposure compared with cover crop mulch alone. Cover crop mulch properties can have

significant impacts on the fate of residual herbicides by influencing interception, retention, and degradation processes in no-till systems (Alletto et al. 2010; Cassigneul et al. 2015). Previous studies suggest that integrating cover crop residues and soil-applied residual herbicides can result in synergistic or antagonistic effects on early-season weed suppression (Teasdale et al. 2003, 2005). Emerging cover-cropping practices that include delaying cover crop termination until or after cash crop planting increase the likelihood of reduced bioavailability of residual herbicides due to interception by cover crop residues. For example, Whalen et al. (2020) reported that the soil concentration and weed control

efficacy of sulfentrazone applied preplant or POST was higher in earlier-terminated (21 d preplant) cover crop systems compared with later-terminated (7 d preplant) systems.

In summary, developing complementary weed control tactics that reduce the intensity of herbicide selection pressure while maintaining weed control and cash crop yield performance targets is imperative in no-till grain crop production systems. Our study indicates that reducing herbicide selection pressure is a potential nonmarket benefit of intensified cover crop management that may be considered with other ecosystem services when assessing long-term net returns to management. In the near term, rotation of herbicide mixtures remains the primary recommendation for prevention of herbicide-resistance evolution (Busi et al. 2019), yet there is increasing concern that evolution of cross-resistance to multiple herbicide sites of action will reduce the long-term efficacy of this strategy (Liu et al. 2020; Nandula et al. 2019). Reducing herbicide inputs in favor of intensified cultural control practices offers an alternative path forward. Our study suggests that high-residue cover crop management offers opportunities for reducing herbicide inputs while meeting crop protection goals but should be integrated with other nonchemical control tactics, such as harvest weed seed control (Shergill et al. 2020), to achieve zero seed rain management goals. Finally, we suggest that herbicide-resistance management and economic returns will likely improve with advances in planter technologies for high-residue cover-cropping systems and improved decision tools for cover crop species and seeding rate selection. Further development of high-residue cover-cropping practices, coupled with a deeper understanding of weed population dynamics in high-residue cover crops, is needed to improve the performance of integrated cover-cropping and herbicide-based tactics to consistently realize IWM goals.

Acknowledgments. This research was funded by USDA-CPPM award 2014-70006-2251 and the Pennsylvania Soybean Board. No conflicts of interest have been declared. We are grateful for the assistance of Tosh Mazzone, Dayton Spackman, and many undergraduate employees.

Supplementary material. To view supplementary material for this article, please visit <https://doi.org/10.1017/wsc.2020.49>

References

- Alletto L, Coquet Y, Benoit P, Heddadj D, Barriuso E (2010) Tillage management effects on pesticide fate in soils. A review. *Agron Sustain Dev* 30:367–400
- Baraibar B, Hunter MC, Schipanski ME, Hamilton AV, Mortensen DA (2018) Weed Suppression in Cover Crop Monocultures and Mixtures. *Weed Sci* 66:121–133
- Bates D, Maechler M, Bolker B, Walker S, Christensen RH, Singmann H, Dai B, Grothendieck G, Green P (2016) lme4: Linear Mixed-Effect Models. <http://CRAN.R-project.org/package=lme4>. Accessed: March 26, 2018
- Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MH, White JS (2008) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends Ecol Evol* 24:127–135
- Busi R, Goggin DE, Heap IM, Horak MJ, Jugulam M, Masters RA, Napier RM, Riar DS, Satchivi NM, Torra J, Westra P, Wright TR (2018) Weed resistance to synthetic auxin herbicides. *Pest Manag Sci* 74:2265–2276
- Busi R, Powles SB, Beckie HJ, Renton M (2019) Rotations and mixtures of soil-applied herbicides delay resistance. *Pest Manag Sci* 76:487–496
- Cassaigne A, Alletto L, Benoit P, Bergheaud V, Etievant V, Dumey V, Le Gac AL, Chuette D, Rumpel C, Justes E (2015) Nature and decomposition degree of cover crops influence pesticide sorption: quantification and modeling. *Chemosphere* 119:1007–1014
- Cholette TB, Soltani N, Hooker DC, Robinson DE, Sikkema PH (2018) Suppression of glyphosate-resistant Canada fleabane (*Conyza canadensis*) in corn with cover crops seeded after wheat harvest the previous year. *Weed Technol* 32:244–250
- Comont D, Hicks H, Crook L, Hull R, Cocciantelli, Hadfield J, Childs D, Freckleton R, Neve P (2019) Evolutionary epidemiology predicts the emergence of glyphosate resistance in a major agricultural weed. *New Phytol* 223:1584–1594
- Dinelli G, Marotti I, Bonetti A, Minelli M, Catizone P, Barnes J (2006) Physiological and molecular insight on the mechanisms of resistance to glyphosate in *Conyza canadensis* (L.) Cronq. biotypes. *Pestic Biochem Physiol* 86:30–41
- Evans JA, Williams A, Hager AG, Mirsky SB, Tranel PJ, Davis AS (2018) Confronting herbicide resistance with cooperative management. *Pest Manag Sci* 74:2424–2431
- Hamilton AV, Mortensen DA, Kammerer Allen M (2017) The state of the cover crop nation and how to set realistic future goals for the popular conservation practice. *J Soil Water Conserv* 72:111A–115A
- Harper J (2016) Farm Management Enterprise Budgets. Penn State Agronomy Guide. University Park: Penn State. Pp 151–162
- Heywood, JS (1986) The effect of plant size variation on genetic drift in populations of annuals. *Am Nat* 137:851–861
- Hothorn T, Bretz F, Westfall P, Heiberger RM, Schuetzenmeister A, Scheibe S (2008) multcomp: Simultaneous Inference in General Parametric Models. R Package v. 3.2.2.4. <http://CRAN.R-project.org/package=multcomp>. Accessed March 26, 2018
- Kniss AR (2018) Genetically engineered herbicide-resistant crops and herbicide-resistant weed evolution in the United States. *Weed Sci* 66:260–273
- Lenth R, Singmann H, Love J, Buerkner PH (2018) emmeans: Estimated Marginal Means, aka LeastSquare Means. R Package v. 1.2.3. <https://cran.r-project.org/web/packages/emmeans/emmeans.pdf>. Accessed: March 26, 2018
- Liebman M, Mohler CL, Staver CP (2001) Ecological Management of Agricultural Weeds. New York: Cambridge University Press
- Liu C, Neve P, Glasgow L, Wuerffel RJ, Owen MDK, Kaundun SS (2020) Modeling the sustainability and economics of stacked herbicide-tolerant traits and early weed management strategy for waterhemp (*Amaranthus tuberculatus*) control. *Weed Sci* 68:179–185
- Loux MM, Dobbels AF, Bradley KW, Johnson WG, Young BG, Spaunhorst DJ, Norsworthy JK, Palhano M, Steckel LE (2017) Influence of cover crops on management of *Amaranthus* species in glyphosate- and glufosinate-resistant soybean. *Weed Technol* 31:487–495
- Mirsky SB, Ryan MR, Curran WS, Teasdale JR, Maul J, Spargo JT, Moyer J, Grantham AM, Weber D, Way T, Camargo GG (2012) Conservation tillage issues: cover crop-based organic rotational no-till grain production in the Mid-Atlantic, USA. *Renew Agric Food Syst* 27:31–40
- Montgomery GB, McClure AT, Hayes RM, Walker FR, Senseman SA, Steckel LE (2018) Dicamba-tolerant soybean combined cover crop to control Palmer amaranth. *Weed Technol* 32:109–115
- Mortensen DA, Egan JF, Maxwell BD, Ryan MR, Smith RG (2012) Navigating a critical juncture for sustainable weed management. *BioScience* 62:75–84
- Nakagawa S, Schielzeth H (2013) A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods Ecol Evol* 4:133–142
- Nandula VK, Riechers DE, Ferhatoglu Y, Barrett M, Duke SO, Dayan FE, Goldberg-Cavalleri A, Tetard-Jones C, Wortley DJ, Onkokesung N, Brazier-Hicks M, Edwards R, Gaines T, Iwakami S, Jugulam M, et al. (2019) Herbicide metabolism: crop selectivity, bioactivation, weed resistance, and regulation. *Weed Sci* 67:149–175
- [NOAA] National Oceanic and Atmospheric Administration Climate Data (2019) Climate Data Online. <https://www.ncdc.noaa.gov/cdo-web>. Accessed: December 2019
- Neve P, Powles S (2005) Recurrent selection with reduced herbicide rates result in the rapid evolution of herbicide resistance in *Lolium rigidum*. *Theor Appl Genet* 110:1154–1166
- Nord EA, Ryan MR, Curran WS, Mortensen DA, Mirsky SB (2012) Effects of management type and timing on weed suppression in soybean no-till planted into roll-crimped cereal rye. *Weed Sci* 60:624–633

- Norsworthy JK, Ward SM, Shaw DR, Llewellyn RS, Nichols RL, Webster TM, Bradley KW, Frisvold G, Powles S, Burgos NR (2012) Reducing the risks of herbicide resistance; best management practices and recommendations. *Weed Sci* 60:31–62
- Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team (2015) nlme: Linear and Nonlinear Mixed Effects Models. R Package v. 3.1.2.4. <http://CRAN.R-project.org/package=nlme>. Accessed: March 26, 2018
- Pittman KB, Barney JN, Flessner ML (2019) Horseweed (*Conyza canadensis*) suppression from cover crop mixtures and fall-applied residual herbicides. *Weed Technol* 33:303–311
- Ryan MR, Curran WS, Grantham AM, Hunsberger LK, Mirsky SB, Mortensen DA, Nord EA, Wilson DO (2011a) Effects of seeding rate and poultry litter on weed suppression from a rolled cereal rye cover crop. *Weed Sci* 59:438–444
- Ryan MR, Mirsky SB, Mortensen DA, Teasdale JR (2011b) Potential synergistic effects of cereal rye biomass and soybean planting density on weed suppression. *Weed Sci* 59:238–246
- Schipanski ME, Barbercheck M, Douglas MR, Finney DM, Haider K, Kaye JP, Kemanian AR, Mortensen DA, Ryan MR, Tooker J, White C (2014) A framework for evaluating ecosystem services provided by cover crops in agroecosystems. *Agric Syst* 125:12–22
- Shergill LS, Bejleri K, Davis A, Mirsky SB (2020) Fate of weed seeds after impact mill processing in midwestern and mid-Atlantic United States. *Weed Sci* 68:92–97
- Smith RG, Atwood LW, Pollnac FW, Warren ND (2015) Cover crop species as distinct biotic filters in weed community assembly. *Weed Sci* 63:282–29
- Teasdale JR, Pillai R, Collins RT (2005) Synergism between cover crop residue and herbicide activity on emergence and early growth of weeds. *Weed Sci* 53:521–527
- Teasdale JR, Shelton DR, Sadeghi AM, Isensee AR (2003) Influence of hairy vetch residue on atrazine and metolachlor soil solution concentration and weed emergence. *Weed Sci* 51:628–634
- [USDA-NASS] United States Department of Agriculture National Agricultural Statistics Service (2019) Agriculture Prices Program. https://www.nass.usda.gov/Surveys/Guide_to_NASS_Surveys/Prices/index.php. Accessed: March 26, 2018
- Wallace JM, Curran WS, Mortensen DA (2019) Cover crop effects on horseweed (*Erigeron canadensis*) density and size inequality at the time of herbicide exposure. *Weed Sci* 67:327–338
- Wallace JM, Keene CL, Curran W, Mirsky S, Ryan MR, VanGessel MJ (2017) Integrated weed management strategies in cover crop-based, organic rotational no-till corn and soybean in the Mid-Atlantic region. *Weed Sci* 66:1–15
- Wayman S, Kucek LK, Mirsky SB, Ackroyd V, Cordeau S, Ryan MR (2016) Organic and conventional farmers differ in their perspectives on cover crop use and breeding. *Renew Agric Food Syst* 32:376–386
- Weiner J (1985) Size hierarchies in experimental populations of annual plants. *Ecology* 66:743–752
- Weisberger D, Nichols V, Liebman M (2019) Does diversifying crop rotations suppress weeds? A meta-analysis. *PLoS ONE* 14:e0219847
- Whalen DM, Shergill LS, Kinne LP, Bish MD (2020) Integration of residual herbicides with cover crop termination in soybean. *Weed Technol* 34:11–18
- Wiggins MS, McClure MA, Hayes RM, Steckel LE (2015) Integrating cover crops and POST herbicides for glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) control in corn. *Weed Technol* 29:412–418
- Wortman SE, Francis CA, Bernards ML, Drijber RA, Lindquist JL (2012) Optimizing cover crop benefits with diverse mixtures and an alternative termination method. *Agron J* 104:1425–1435