



## **Unraveling the Mechanism of Resistance in a Glufosinate-Resistant Palmer Amaranth (*Amaranthus palmeri*) Accession**

Authors: Carvalho-Moore, Pamela, Norsworthy, Jason K., González-Torralva, Fidel, Hwang, Jeong-In, Patel, Jinesh D., et al.

Source: Weed Science, 70(4) : 370-379

Published By: Weed Science Society of America

URL: <https://doi.org/10.1017/wsc.2022.31>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Research Article

**Cite this article:** Carvalho-Moore P, Norsworthy JK, González-Torralva F, Hwang J-I, Patel JD, Barber LT, Butts TR, McElroy JS (2022) Unraveling the mechanism of resistance in a glufosinate-resistant Palmer amaranth (*Amaranthus palmeri*) accession. *Weed Sci.* **70**: 370–379. doi: [10.1017/wsc.2022.31](https://doi.org/10.1017/wsc.2022.31)

Received: 6 March 2022

Revised: 4 May 2022

Accepted: 24 May 2022

First published online: 3 June 2022

**Associate Editor:**

Mithila Jugulam, Kansas State University

**Keywords:**

Glutamine synthetase; herbicide resistance; gene amplification

**Author for correspondence:**

Pamela Carvalho-Moore, University of Arkansas, 1354 W Altheimer Drive, Fayetteville, AR 72704. Email: [pcarvalh@uark.edu](mailto:pcarvalh@uark.edu)

# Unraveling the mechanism of resistance in a glufosinate-resistant Palmer amaranth (*Amaranthus palmeri*) accession

Pamela Carvalho-Moore<sup>1</sup> , Jason K. Norsworthy<sup>2</sup> , Fidel González-Torralva<sup>3</sup> , Jeong-In Hwang<sup>3</sup> , Jinesh D. Patel<sup>4</sup> , L. Tom Barber<sup>5</sup>, Thomas R. Butts<sup>6</sup>  and J. Scott McElroy<sup>7</sup> 

<sup>1</sup>Graduate Research Assistant, Department of Crop, Soil, and Environmental Sciences, University of Arkansas, Fayetteville, AR, USA; <sup>2</sup>Professor and Elms Farming Chair of Weed Science, Department of Crop, Soil, and Environmental Sciences, University of Arkansas, Fayetteville, AR, USA; <sup>3</sup>Postdoctoral Fellow, Department of Crop, Soil, and Environmental Sciences, University of Arkansas, Fayetteville, AR, USA; <sup>4</sup>Research Associate, Department of Crop, Soil, and Environmental Sciences, Auburn University, Auburn, AL, USA; <sup>5</sup>Professor and Extension Weed Scientist, Department of Crop, Soil, and Environmental Sciences, University of Arkansas, Lonoke, AR, USA; <sup>6</sup>Assistant Professor and Extension Weed Scientist, Department of Crop, Soil, and Environmental Sciences, University of Arkansas, Lonoke, AR, USA and <sup>7</sup>Professor, Department of Crop, Soil, and Environmental Sciences, Auburn University, Auburn, AL, USA

**Abstract**

Glufosinate resistance in Palmer amaranth (*Amaranthus palmeri* S. Watson) was recently detected in three accessions from Arkansas, USA. *Amaranthus palmeri* is the first and only broadleaf weed species resistant to this herbicide, and the resistance mechanism is still unclear. A previous study characterized the glufosinate resistance level in the accessions from Arkansas. A highly glufosinate-resistant accession was further used to investigate the mechanism conferring glufosinate resistance in *A. palmeri*. Experiments were designed to sequence the herbicide target enzyme cytosolic and chloroplastic glutamine synthetase isoforms (*GS1* and *GS2*, respectively) and quantify copy number and expression. Absorption, translocation, and metabolism of glufosinate using the <sup>14</sup>C-labeled herbicide were also evaluated in the resistant and susceptible accessions. The glufosinate-resistant accession had an increase in copy number and expression of *GS2* compared with susceptible plants. All accessions showed only one *GS1* copy and no differences in expression. No mutations were identified in *GS1* or *GS2*. Absorption (54% to 60%) and metabolism (13% to 21%) were not different between the glufosinate-resistant and glufosinate-susceptible accessions. Most residues of glufosinate (94% to 98%) were present in the treated leaf. Glufosinate translocation to tissues above the treated leaf and in the roots was not different among accessions. However, glufosinate translocation to tissues below the treated leaf (not including roots) was greater in the resistant *A. palmeri* (2%) compared with the susceptible (less than 1%) accessions. The findings of this paper strongly indicate that gene amplification and increased expression of the chloroplastic glutamine synthetase enzyme are the mechanisms conferring glufosinate resistance in the *A. palmeri* accession investigated. Thus far, no additional resistance mechanism was observed, but further investigations are ongoing.

**Introduction**

Glufosinate-ammonium is a broad-spectrum, nonselective herbicide labeled to control dicotyledons and certain monocotyledons. In the United States, this herbicide was first registered in 1993 by AgrEvo under the commercial names Finale<sup>®</sup> and Rely<sup>®</sup> (Bijman 2001; Hoerlein 1994). Glufosinate is one of the foundational postemergence herbicides in LibertyLink<sup>®</sup> (BASF, Florham Park, NJ, USA), Enlist<sup>™</sup> (Corteva Agriscience, Indianapolis, IN, USA), and XtendFlex<sup>®</sup> (Bayer CropScience, Pittsburgh, PA, USA) systems, in which crops contain the glufosinate-resistance trait. Glufosinate controls plants by inhibiting the glutamine synthetase enzyme, which synthesizes glutamine from glutamate with photorespiratory ammonia. Because the herbicide is a chemical analogue to glutamate, it competes with this amino acid to bind to the enzyme. Inhibition of glutamine synthetase eventually leads to ammonia accumulation, amino acid depletion, detrimental accumulation of reactive oxygen species, lipid peroxidation, and, ultimately, cell death (Bayer et al. 1972; Takano et al. 2019, 2020; Wendler et al. 1990; Wild and Manderscheid 1984). In plants, glutamine synthetase has two major isoforms in different compartments: one located in the cytosol (*GS1*) and the other located in the chloroplasts (*GS2*) (Mann et al. 1979). *GS1* is associated to nitrogen assimilation that will generate glutamine for nitrogen transport inside plants, while *GS2* is involved in

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



the reassimilation of ammonium from the photorespiratory pathway and other plant processes (Edwards et al. 1990; Kamachi et al. 1991; Wallsgrove et al. 1987).

Compared with other frequently used nonselective herbicides such as glyphosate and paraquat, glufosinate has a relatively lower number of resistant weed species. Currently, only five weed species have been confirmed resistant to glufosinate. These weeds are goosegrass [*Eleusine indica* (L.) Gaertn.], Italian ryegrass [*Lolium perenne* L. ssp. *multiflorum* (Lam.) Husnot], perennial ryegrass (*Lolium perenne* L.), rigid ryegrass (*Lolium rigidum* Gaudin), and Palmer amaranth (*Amaranthus palmeri* S. Watson). *Amaranthus palmeri* is the first and only broadleaf weed resistant to glufosinate (Avila-Garcia and Mallory-Smith 2011; Ghanizadeh et al. 2015; Heap 2022b; Jalaludin et al. 2010; Priess et al. 2022; Seng et al. 2010; Travlos et al. 2018).

According to the Herbicide Resistance Action Committee, confirmation of herbicide resistance starts with the seed collection from weeds surviving the lethal dose of a specific product, followed by herbicide screening and a dose-response experiment with whole plants in a greenhouse (Heap 2005). In 2019 and 2020, Arkansas farmers from Crittenden and Mississippi counties noticed that glufosinate applications failed to control *A. palmeri* plants. Seeds from these fields were collected, and glufosinate resistance was confirmed through dose-response experiments (Priess et al. 2022). In addition to glufosinate, *A. palmeri* is resistant to herbicides targeting eight other sites of action, which makes this weed extremely challenging to control (Heap 2022b). Although the resistance has been confirmed, the resistance mechanism in glufosinate-resistant *A. palmeri* remains unrevealed.

The possible mechanisms of resistance are divided into target-site and non-target site mechanisms. Target-site resistance encompasses any alteration in the target enzyme that will prevent herbicide binding, such as amino acid/nucleotide change. Gene amplification and overexpression of the targeted protein are also considered to be target-site resistance mechanisms. Non-target site resistance is any plant mechanism that reduces the quantity of herbicide reaching the target site (Délye et al. 2013; Powles and Yu 2010). It is crucial to understand the basis of herbicide resistance and the biology of a weed population. With this knowledge, scientists can design and apply proper strategies to limit the spread of resistance (Norsworthy et al. 2012). Glufosinate resistance was characterized in three *A. palmeri* accessions from Arkansas by Priess et al. (2022), and a high resistance level was observed in one of these accessions (resistance/susceptibility ratio = 24). In this study, this highly glufosinate-resistant accession was further used to investigate the mechanism conferring resistance to glufosinate.

## Materials and Methods

One *A. palmeri* accession previously confirmed to have high glufosinate resistance (resistance/susceptibility ratio = 24) was selected to conduct the experiments described. The seeds from the resistant accession (Glu-R1) were collected in 2020 from a cotton (*Gossypium hirsutum* L.) field where glufosinate applications failed to provide control (Priess et al. 2022). Additionally, two susceptible *A. palmeri* accessions collected in South Carolina in 1986 (S1) and in Arkansas in 2001 (S2) were used for comparison. Seedlings were established in a greenhouse at 25 ± 5 °C and 16-h day at the Milo J. Shult Agricultural Research and Extension Center in Fayetteville, AR, USA.

## Glutamine Synthetase Sequencing

Illumina sequencing was conducted on RNA from Glu-R1 survivors and S1 nontreated plants to identify the presence of GS1 and GS2 mutations possibly correlated with target-site resistance. Approximately 1 g of leaf tissue was collected from glufosinate-resistant and glufosinate-susceptible *A. palmeri* accessions, frozen using liquid nitrogen, and then ground to fine powder. The finely powdered leaf tissue of each accession was processed using the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) to extract RNA. The quality and quantity of RNA were determined using a NanoDrop spectrophotometer (ThermoFisher Scientific, Waltham, MA, USA). RNA samples were immediately stored at -80 °C until further processing for transcriptomic sequencing. RNA samples of the Glu-R1 and S1 accessions of *A. palmeri* were analyzed using the Agilent Bioanalyzer 2100 instrument (Agilent Technologies, Palo Alto, CA, USA) to determine the RNA integrity number. Prepared libraries were run on Illumina NovaSeq 6000 instrument (Novogene, Beijing, China) to produce 150-bp paired-end reads. The sequence read data were evaluated to determine the percentage of reads containing adapters, reads containing N > 10% (N represents the base that cannot be determined), and reads of low quality (Qscore ≤ 5) before releasing the data.

Paired-end reads were assembled using Trinity (<https://github.com/trinityrnaseq>), with standard flags of Trimmomatic (Bolger et al. 2014). Assembled contigs were annotated using Trinotate (<https://trinotate.github.io>), and peptide sequences were produced using TransDecoder (<http://transdecoder.github.io>). Glutamine synthetase nucleotide and peptide sequences were extracted from fasta files using the TrinotateExtractor (<http://github.com/mcelrjo/trinotateExtractor>) based on Blast annotation in the Trinotate output file. Protein sequences were aligned within each accession to eliminate redundant sequences and were sorted within each accession to refine annotation to specific glutamine synthetase orthologues based on Blast annotation. The Glu-R1 and S1 protein sequences derived from TransDecoder were aligned for GS1 and GS2 to identify any amino acid differences. Protein sequences were aligned to reference protein sequences of glutamine synthetase cytosolic and chloroplastic isozymes from *A. palmeri* (NCBI accession GFQG01042326.1 and Heap [2022a], respectively). The GS1 and GS2 isozymes of cantaloupe (*Cucumis melo* L.) (NCBI accessions NP\_001284433.1 and NP\_001284439.1, respectively) were included as an unrelated species to demonstrate the sequence homology across diverse taxa. Illumina sequencing reads for Glu-R1 and S1 were submitted as NCBI BioProject PRJNA831848.

Read mapping was performed to identify numerical differences in expression between Glu-R1 and S1 accessions. Illumina sequencing of the Glu-R1 biotype generated 22,582,008 paired-end reads, while sequencing of the S1 biotype generated 10,883,584 paired-end reads. Reads were mapped using the CLC Genomics Workbench (CLCbio, Seoul, Republic of Korea) read mapping tool with the following settings: no masking, match score 1, mismatch cost 2, linear gap cost for insertions and deletions, insertion cost 3, deletion cost 3, length fraction of 0.5, similarity fraction of 0.8, and no global alignment. Read counts, average coverage of reads, and the percent of the total reads mapped are presented in Supplementary Table 1.

## Glutamine Synthetase Gene Copy Number Quantification

Gene copy number assay was conducted with nontreated plants from the susceptible accessions (S1 and S2), and glufosinate survivors from the resistant accession (Glu-R1) sprayed with glufosinate

**Table 1.** Primer pairs used to quantify relative copy number and gene expression by real-time polymerase chain reaction in *Amaranthus palmeri* accessions.

| Targeted gene <sup>a</sup> | Primer sequence <sup>b</sup>                                       | Amplicon —bp— | Efficiency —%— |
|----------------------------|--------------------------------------------------------------------|---------------|----------------|
| <i>Gs1a</i>                | F 5'-CTGGAATGGTGTCTGGTCT-3'<br>R 5'-TGTGCTCTTTGTGCCTCAAC-3'        | 116           | 94.3           |
| <i>Gs2a</i>                | F 5'-ATCGTGGTTGCTCTATCCGTG-3'<br>R 5'-GTTTCTGCGAGCAACCTGTT-3'      | 121           | 102.9          |
| <i>Gs1b</i>                | F 5'-AACCATGGTACGGTATCGAACAGG-3'<br>R 5'-AGGCAAGCCTTGAGTGTGAATC-3' | 172           | — <sup>c</sup> |
| <i>Gs2b</i>                | F 5'-AAGGATCCATTCCGTGGTGG-3'<br>R 5'-TCTCAGAAACAACCTTTGGGTCG-3'    | 133           | — <sup>c</sup> |
| <i>CCR</i>                 | F 5'-CGACGGAAATAGCAACAAAGTG-3'<br>R 5'-GTCCTTGACGGTGGCGTTAAC-3'    | 116           | — <sup>c</sup> |
| <i>PPAN</i>                | F 5'-TGCTTCATTTTGAGGGTTCG-3'<br>R 5'-GACATCGAGGCTCAACTGTG-3'       | 113           | — <sup>c</sup> |

<sup>a</sup>*Gs1a* (cytoplasmatic) and *Gs2a* (chloroplastic) glutamine synthetase in gene copy number experiment; *Gs1b* (cytoplasmatic) and *Gs2b* (chloroplastic) glutamine synthetase in gene expression experiment; *CCR*, Cinnamoyl-CoA reductase; *PPAN*, peter Pan-like.

<sup>b</sup>F, forward; R, reverse.

<sup>c</sup>Primer information is presented in Takano and Dayan (2021) and González-Torralva and Norsworthy (2021).

(Interline<sup>®</sup>, UPL Limited, King of Prussia, PA, USA) at 656 g ai ha<sup>-1</sup> at the 5- to 7-leaf stage. Around 100 mg of leaf tissue was collected per plant from four plants of each accession, and genomic DNA was extracted using the E.Z.N.A.<sup>®</sup> Plant DNA Kit (Omega Bio-Tek, Norcross, GA, USA) according to the manufacturer's directions. After extraction, DNA was quantified using a NanoDrop spectrophotometer (ThermoFisher Scientific, Waltham, MA, USA) and diluted with deionized water to 10 ng µl<sup>-1</sup>. A quantitative real-time polymerase chain reaction (qPCR) was conducted to quantify the cytoplasmatic (*GS1*) and chloroplastic (*GS2*) glutamine synthetase copy number. The primers *GS1a* and *GS2a* were designed to quantify gene copy number for *GS1* or *GS2*, respectively (Table 1).

The qPCR reaction mixture (20 µl) consisted of 10 µl of 2× SsoAdvanced Universal SYBR Green Supermix (Bio-Rad Laboratories, Hercules, CA, USA), 0.8 µl each of 10 µM forward and reverse primers (Table 1), 5.9 µl of deionized water, and 25 ng of genomic DNA. The assay was conducted in a CFX96 Real-Time System (Bio-Rad Laboratories) under the following conditions: 98 C for 3 min, 40 cycles of 98 C for 10 s, and 61 C for 30 s. Melting curves were created by increasing the temperature from 65 C to 95 C, 0.5 C every 5 s to ensure specific amplification. Each biological sample (total of four per accession) had two technical replicates in each primer pair, and the experiment was repeated in time. No-template controls (DNA substituted by deionized water) were included in each plate. Quantification cycles (Cq) were produced by CFX Maestro software (Bio-Rad Laboratories), and genomic copy numbers of *GS1* and *GS2* were calculated using a modified version of the 2<sup>-ΔΔCt</sup> method (Gaines et al. 2010; Livak and Schmittgen 2001). Fold increase in *GS1* and *GS2* was assessed relative to two reference genes (single gene copy) previously used in *A. palmeri*, Cinnamoyl-CoA reductase (*CCR*), and Peter Pan-like (*PPAN*) (González-Torralva and Norsworthy 2021; Salas et al. 2012).

### Glutamine Synthetase Gene Expression

The gene expression assay was conducted with nontreated plants from the same susceptible and glufosinate-resistant accessions

described earlier. Approximately 100 mg of leaf tissue was collected per plant from three plants of each accession and immediately frozen in liquid nitrogen. RNA was extracted using the Monarch Total RNA Miniprep Kit (New England Biolabs, Ipswich, MA, USA) according to the manufacturer's instructions. After extraction and quantification in a spectrophotometer, a total of 1 µg RNA per sample was reverse transcribed into complementary DNA (cDNA) using the iScript Reverse Transcription Supermix (Bio-Rad Laboratories). This cDNA was used to quantify the cytoplasmatic (*GS1*) and chloroplastic (*GS2*) glutamine synthetase expression using primer pairs *GS1b* and *GS2b*, respectively (Table 1).

The qPCR reaction mixture followed the same methodology described in "Glutamine Synthetase Gene Copy Number Quantification" with forward and reverse primers described in Table 1. The primer pairs *GS1b* and *GS2b* amplifying *GS1* and *GS2*, respectively, were obtained from a previous study targeting these genes in *A. palmeri* (Takano and Dayan 2021). The assay was conducted under the following conditions: 30 s at 98 C, 40 cycles of 98 C for 10 s, and 60 C for 30 s. After the cycle was completed, the temperature increased by 0.5 C every 5 s from 65 C to 95 C to generate melting curves. Two technical replicates were used for each biological sample (total of three per accession), and the experiment was repeated in time. No-template controls were included on each plate. Quantification cycles (Cq) were produced by CFX Maestro software (Bio-Rad Laboratories). The Ct values of *GS1* and *GS2* in resistant and susceptible plants were normalized against the reference genes *CCR* and *PPAN* (Table 1), and relative expression was calculated against susceptible accessions using the 2<sup>-ΔΔCt</sup> method (Livak and Schmittgen 2001).

### Absorption and Translocation of Glufosinate

Absorption and translocation experiments of <sup>14</sup>C-labeled glufosinate were conducted in resistant (Glu-R1) and susceptible (S1 and S2) accessions. *Amaranthus palmeri* seedlings were transplanted into 7-cm-diameter plastic pots filled with potting mix (Sun Gro<sup>®</sup> Horticulture, Agawam, MA, USA). Each accession had three replications that consisted of one plant per pot in each replication. The experiment was organized in a completely randomized design and repeated twice. At the 6- to 8-leaf stage, plants received an overspray of nonradioactive glufosinate at 656 g ai ha<sup>-1</sup>. The overspray was applied using a spray chamber equipped with 1100067 nozzles (TeeJet<sup>®</sup> Technologies, Springfield, IL, USA) calibrated to deliver 187 L ha<sup>-1</sup> at 1.6 km h<sup>-1</sup>. Immediately after nonradioactive herbicide application, plants were spotted with [<sup>14</sup>C]glufosinate. A working solution of [<sup>14</sup>C]glufosinate was prepared in the overspray solution. Four 0.5-µl droplets, each containing 1 kBq of [<sup>14</sup>C]glufosinate, were placed on the second fully expanded leaf of each treated plant. Therefore, a total of 4 kBq of radiolabeled herbicide was spotted per plant. The treated plants were maintained in a greenhouse set to 25 C under a 14-h light/10-h dark photoperiod. There was no function inside the greenhouse to regulate the relative humidity, and this factor was relatively low (≤50%) throughout both runs. Nontreated plants were maintained as a control.

At 48 h after <sup>14</sup>C-labeled herbicide treatment, plants were collected and divided into four sections: treated leaf (TL), above treated leaf (ATL), below treated leaf (BTL), and roots. Roots were washed to remove soil residues. The treated leaf was collected from each plant and rinsed with 5 ml of methanol. The rinsate was gathered in a 20-ml scintillation vial containing 10 ml of scintillation



cocktail (Ultima Gold™, PerkinElmer, Waltham, MA, USA) and analyzed with a Tri-Carb 2900TR Liquid Scintillation Analyzer (LSA; PerkinElmer). Absorption was calculated by subtracting the [ $^{14}\text{C}$ ]glufosinate activity in the rinsate of the treated leaf at the sampling time from the [ $^{14}\text{C}$ ]glufosinate activity in the rinsate of the treated leaf at the initial time. The initial [ $^{14}\text{C}$ ]glufosinate recovery was 95%, and it was obtained by washing treated leaves soon after spotting treatment.

Each plant section was individually placed into a paper envelope and then dried in a freeze-dryer (Botanique Preservation Equipment, Phoenix, AZ, USA) at  $-50\text{ }^{\circ}\text{C}$  for 72 h. The dried plant parts were individually combusted to  $^{14}\text{CO}_2$  by a biological oxidizer (Model OX-700, R.J. Harvey Instruments, Tappan, NY, USA) at  $900\text{ }^{\circ}\text{C}$  for 3 min. The  $^{14}\text{CO}_2$  gas was entrapped into 15 ml of  $^{14}\text{C}$ -trapping cocktail (R.J. Harvey Instrument). The  $^{14}\text{C}$  activity in the vials was analyzed using the LSA. Translocation was calculated as the proportion of the [ $^{14}\text{C}$ ]glufosinate measured in each plant part relative to the total [ $^{14}\text{C}$ ]glufosinate absorbed after 48 h.

### Metabolism of Glufosinate

Metabolism experiments were conducted twice on the same dates on which the absorption/translocation experiment were being performed. *Amaranthus palmeri* plant sample preparation, non-radiolabeled herbicide spray, and  $^{14}\text{C}$ -labeled herbicide treatment were done following the same methods as used in the absorption/translocation experiment. At 48 h after  $^{14}\text{C}$ -labeled herbicide treatment, the treated leaf was thoroughly rinsed with 15 to 20 ml of methanol to eliminate the  $^{14}\text{C}$ -labeled herbicide persisting on the leaf surface. Following the leaf wash, each plant was placed into a paper envelope without dissection and then freeze-dried for 72 h.

Extraction of [ $^{14}\text{C}$ ]glufosinate and its metabolites from metabolism samples was performed based on the analytical method proposed by Küpper et al. (2018) and Meyer et al. (2020). The dried plant sample was cut into small pieces ( $<2\text{ mm}$ ) and then transferred into a 2-ml Eppendorf tube containing 600  $\mu\text{l}$  of 90% methanol in water. Subsequently, the sample was thoroughly ground using a polypropylene pellet pestle. After 30 s of vortexing, the sample was left at  $4\text{ }^{\circ}\text{C}$  in a refrigerator for 1 h and then centrifuged at  $8000\times g$  for 6 min. The supernatant was transferred to an evaporating flask. The residue remaining in the tube was additionally extracted with 600  $\mu\text{l}$  of 90% acetonitrile in water, followed by 600  $\mu\text{l}$  of 10% methanol in water. All supernatants of additional extracts were combined on the same evaporating flask with the initial supernatant and then evaporated to  $<1\text{ ml}$  using an Xcelvap evaporator (Horizon Technologies, Lake Forest, CA, USA). The evaporated sample was reconstituted to 1 ml with methanol and filtered using a 0.2- $\mu\text{l}$  polytetrafluoroethylene (PTFE) syringe filter. The quantification of [ $^{14}\text{C}$ ]glufosinate in the samples was conducted using high-performance liquid chromatography (HPLC; Prominence-i LC-2030C 3D, Shimadzu, Kyoto, Japan) linked to a LabLogic Beta-Ram Model 4B radiation detector (RAD; LabLogic Systems, Tampa, FL, USA).

A 25- $\mu\text{l}$  aliquot from the final sample solution was injected into the HPLC–RAD system. The mobile phases consisted of 50 mM ammonium acetate in water (A) and HPLC reagent grade water (B). Solvents were run for a 1-min plateau at 15% solvent A, a 5-min linear gradient from 15% to 30% of solvent A plateauing for 2 min, followed by a linear gradient returning to 15% solvent A in 5 min. The column was then flushed with 15% solvent A for 2 min. A SeQuant ZIC-pHILIC 5  $\mu\text{M}$  polymeric LC column (100 cm in length by 4.6 mm in inner diameter, Merck KGaA,

Darmstadt, Germany) was used to separate [ $^{14}\text{C}$ ]glufosinate from the sample matrices. The column oven temperature was kept at  $40\text{ }^{\circ}\text{C}$ , and the flow rate of mobile phases was  $0.5\text{ ml min}^{-1}$ . The recovery in the blank *A. palmeri* sample treated with 1.0 kBq of [ $^{14}\text{C}$ ]glufosinate was  $>85\%$ . Metabolism (%) was calculated by subtracting the  $^{14}\text{C}$  activity of the parent compound from the  $^{14}\text{C}$  activity absorbed at 48 h after [ $^{14}\text{C}$ ]glufosinate treatment.

### Data Analysis


All data collected were subjected to ANOVA in JMP Pro v. 15 (SAS Institute, Cary, NC, USA). The experimental runs were not significant across the experiments and were thus set as a random effect in the subsequent statistical model statement. If significant, means from gene copy number and gene expression assays were separated using Fisher's protected LSD at  $\alpha = 0.05$ . Results of absorption, translocation, and metabolism were also subjected to ANOVA and separated using Tukey's honestly significant difference (HSD) at  $\alpha = 0.05$ .

## Results and Discussion

### Glutamine Synthetase Sequencing, Relative Copy Number, and Expression

Protein isoforms were compared to identify any amino acid substitutions present in the resistant (Glu-R1) in comparison to the susceptible (S1) *A. palmeri* accession and reference sequences (*A. palmeri* and *C. melo*). Based on amino acid sequences, one isoform of cytosolic resistant (Glu-R1) and susceptible (S1) was identified and aligned with the reference cytosolic GS (Supplementary Figure 1). The sequences did not assemble correctly past Trp-306 due to poor alignment quality past this point of the sequencing. While differences between the *C. melo* reference and Glu-R1 and S1 were observed, no differences were observed between the *A. palmeri* accessions (reference, Glu-R1, and S1). For chloroplastic GS, only one isoform was identified (Supplementary Figure 2). The Glu-R1 GS2 isoform (GSChl\_Glu-R1) contained one amino acid substitution (Gly-20-Ser) not present in GS2 from S1 or the *A. palmeri* reference. However, Blastp alignment inspection of this region identified that this position shows high amino acid polymorphism across different species. Due to this lack of evolutionary conservation, it is unlikely that this substitution would constitute a target-site resistance, but additional studies such as cloning vectors need to be conducted to prove this hypothesis. Little numerical difference was observed in GS1 between Glu-R1 and S1 reads (Supplementary Table 1). For GS1, 0.013% to 0.035% of total reads mapped to either accession with no obvious numerical differences. For GS2, greater numerical difference was observed between the biotype reads. Glu-R1 reads mapped to 0.079% to 0.242% of the total reads, while S1 reads only mapped to 0.018% of total reads. The mapping difference translates to 4.4 to 13.4 times greater expression for Glu-R1 than S1.

Sequence data for both GS1 and GS2 suggest that a point mutation does not contribute to glufosinate resistance in the Glu-R1 accession. Working with a glufosinate-resistant *L. perenne* accession, Avila-Garcia et al. (2012) identified an amino acid substitution of aspartate for asparagine in the GS2 gene, which was initially considered the resistance mechanism. However, further investigation led to the conclusion that this alteration in GS2 could not account for glufosinate resistance in this specific accession (Brunharo et al. 2019). In a recent investigation




**Figure 1.** Cytoplasmic glutamine synthetase copy number (A) and expression (B) relative to *Cinnamoyl-CoA reductase* (CCR) and *peter Pan-like* (PPAN) reference genes in glufosinate-resistant (Glu-R1) and glufosinate-susceptible (S1 and S2) accessions. Error bars represent standard errors of the means ( $n = 8$  and  $n = 6$ ). Means were subjected to ANOVA, and P-values were generated using JMP Pro v. 15 (SAS Institute, Cary, NC).

conducted with glufosinate-resistant *E. indica*, Zhang et al. (2022) identified the substitution of a serine for glycine at the 59th position of GS1 in the resistant accession, which is likely the resistance mechanism in this accession. This substitution was also encountered in glufosinate-resistant *E. indica* from different regions and countries. Mutation in the target enzyme might confer glufosinate resistance in the other *A. palmeri* accessions not tested in this study.


Gene copy number and gene expression assays were conducted to detect any differences in the cytoplasmic and chloroplastic glutamine synthetase (GS1 and GS2, respectively) among the *A. palmeri* biotypes. Regarding the GS1 copy number relative to the reference primers, there was no difference between susceptible and resistant plants, and all accessions had approximately one GS1 copy (Figure 1A). Similarly, GS1 expression was not different within the accessions (Figure 1B). Based on these results, glufosinate resistance in *A. palmeri* did not involve gene amplification or increased expression of cytosolic glutamine synthetase. The cytosolic isoform of the GS enzyme is essential in the assimilation and transport of nitrogen throughout the plant. GS1 and GS2 enzyme

activities greatly vary among species, plant sections, and plant stages (Bernard and Habash 2009; Brugiére et al. 2000; Habash et al. 2001; McNally et al. 1983; Miflin and Habash 2002; Woo et al. 1982). After applications of glufosinate to *A. palmeri* seedlings from Colorado, higher GS1 expression in old leaf or root tissues compared with young leaf tissues was found. Glufosinate has no soil activity and is recommended to be applied when *A. palmeri* plants are small and young; therefore, gene amplification or overexpression of GS1 might not impact resistance due to low GS1 expression in this plant stage.

GS2 copy number and expression were different within the glufosinate-resistant and glufosinate-susceptible accessions. Calculated against CCR and PPAN reference genes, accession Glu-R1 had 85 and 86 copies, respectively, while the two susceptible accessions had 2 GS2 copies (Figure 2A). For gene expression, accession Glu-R1 showed 15 and 31 times relative GS2 expression increase relative to CCR and PPAN, respectively (Figure 2B). The susceptible accessions showed no increase in GS2 expression. In a dose-response experiment, accession Glu-R1 showed 24-fold glufosinate resistance (Priess et al. 2022). The resistance fold



**Figure 2.** Chloroplastic glutamine synthetase copy number (A) and expression (B) relative to *Cinnamoyl-CoA reductase* (CCR) and *peter Pan-like* (PPAN) reference genes in glufosinate-resistant (Glu-R1) and glufosinate-susceptible (S1 and S2) accessions. Error bars represent standard errors of the means ( $n = 8$  and  $n = 6$ ). Means were subjected to ANOVA, and P-values were generated using JMP Pro v. 15 (SAS Institute, Cary, NC). Means displayed with different uppercase letters are different according to Fisher's protected LSD test at  $\alpha = 0.05$ .



**Figure 3.** Absorption of  $[^{14}\text{C}]$ glufosinate by glufosinate-resistant (Glu-R1) and glufosinate-susceptible (S1 and S2) accessions assessed at 48 h after application of radiolabeled herbicide. Error bars represent standard errors of the means ( $n = 6$ ). Means were subjected to ANOVA, and P-values were generated using JMP Pro v. 15 (SAS Institute, Cary, NC).  
 Downloaded From: <https://bioone.org/journals/Weed-Science> on 27 Jul 2024  
 Terms of Use: <https://bioone.org/terms-of-use>

obtained by Priess et al. (2022) and the relative GS2 expression obtained in this study for accession Glu-R1 are similar, suggesting that high copy number and expression may confer high resistance. Therefore, gene amplification and overexpression of GS2 enzyme is likely the mechanism conferring glufosinate resistance in the *A. palmeri* accessions investigated in this study. Further experiments, such as examinations of enzyme activity and inheritance of this trait, remain to be conducted. This is the first report of increase in GS2 copy number or expression in a glufosinate-resistant species.


Herbicide resistance by gene amplification and overexpression has been previously reported in *A. palmeri*. Resistance to glyphosate can be due to an increased copy number of the target enzyme, 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS; Gaines et al. 2010; Singh et al. 2018). Working with glyphosate-resistant *A. palmeri* accessions, Ribeiro et al. (2014) found that EPSPS relative copy number had a positive correlation with EPSPS expression, results similar to those obtained in the present study. It is important to note that Glu-R1 also exhibits resistance to glyphosate (Priess et al. 2022). Likewise, increased EPSPS relative copy number was observed in this accession (data not shown). Other glyphosate-resistant weeds, such as kochia [*Bassia scoparia* (L.) A.J. Scott.] and *E. indica*, also exhibit gene amplification as the resistance mechanism (Chen et al. 2015; Gaines et al. 2016; Godar et al. 2015). Gene amplification has also been observed in large crabgrass [*Digitaria sanguinalis* (L.) Scop.] resistant to acetyl-CoA carboxylase inhibitors (Laforest et al. 2017). Although seldom identified as a herbicide-resistance mechanism, increase in gene copy number has been reported in several cases of fungicide or insecticide resistance (Anthony et al. 1998; Cattell et al. 2021; Elmore et al. 2015; Heckel 2022; Puinean et al. 2010).

Interestingly, the gene amplification data strongly suggest that the GS2 gene in *A. palmeri* has two copies natively in susceptible plants (Figure 2A). A future experiment involving other *A. palmeri* accessions from different regions might further investigate this finding. GS2 duplication has been observed before in black cottonwood [*Populus balsamifera* L. ssp. *trichocarpa* (Torr. & A. Gray ex Hook.) Brayshaw], barrelclover (*Medicago truncatula* Gaertn.), and some green algae species (Castro-Rodríguez et al. 2011; Ghoshroy et al. 2010; Seabra et al. 2010).

#### <sup>14</sup>C]Glufosinate Absorption, Translocation, and Metabolism

No difference in glufosinate absorption was observed among resistant and susceptible accessions (Figure 3). Absorption ranged from 54% to 60% in the accessions, similar to the magnitude (59% to 85%) reported in previous studies with *A. palmeri* (Everman et al. 2009b; Meyer et al. 2020).


Most of the [<sup>14</sup>C]glufosinate absorbed remained in the treated leaf: 94% for Glu-R1 and 98% for susceptible accessions ( $P < 0.05$ ) (Figure 4). Translocation of [<sup>14</sup>C]glufosinate to tissues above the treated leaf (1% to 1.9%) and roots (<1% to 1.4%) showed no difference among accessions. However, glufosinate translocation to tissues below the treated leaf (not including roots) was slightly greater in the Glu-R1 accession (2%) than in the susceptible accessions (<1%). Although the herbicide translocation to tissues below the treated leaf was statistically significant among the accessions, this negligible difference (1%) seems to be insufficient to explain the mechanism of glufosinate resistance. The low translocation (<6%) of [<sup>14</sup>C]glufosinate observed in this study was likely due to the localized phytotoxicity and rapid tissue necrosis caused by glufosinate, which possibly restrain translocation to other plants



**Figure 4.** Translocation of [<sup>14</sup>C]glufosinate by glufosinate-resistant (Glu-R1) and susceptible (S1 and S2) accessions assessed at 48 h after application by plant section. ABT, above-treated leaf; TL, treated leaf; BTL, below treated leaf. Error bars represent standard errors of the means ( $n = 6$ ). Means were subjected to ANOVA, and P-values were generated using JMP Pro v. 15 (SAS Institute, Cary, NC). Means displayed with different uppercase letters are different according to Tukey's honestly significant difference (HSD) test at  $\alpha = 0.05$ .

parts (Beriault et al. 1999; Steckel et al. 1997). Low glufosinate translocation has been previously observed in *A. palmeri* and other weed species such as pitted morningglory (*Ipomoea lacunosa* L.) (Everman et al. 2009a, 2009b; Meyer et al. 2020; Steckel et al. 1997).





**Figure 5.** Metabolism of [ $^{14}\text{C}$ ]glufosinate by glufosinate-resistant (Glu-R1) and glufosinate-susceptible (S1 and S2) accessions assessed at 48 h after application. Error bars represent standard errors of the means ( $n = 6$ ). Means were subjected to ANOVA, and P-values were generated using JMP Pro v. 15 (SAS Institute, Cary, NC).

Metabolism did not differ between the glufosinate-resistant and glufosinate-susceptible accessions, and the accessions metabolized 13% to 21% of the absorbed glufosinate at 48 h after treatment (Figure 5; Supplementary Figure 3). Previous studies reported total glufosinate metabolites in *A. palmeri* ranging from 31% to 62% of the total  $^{14}\text{C}$  absorbed (Everman et al. 2009b; Meyer et al. 2020). Metabolism varied from 20% to 30% in the broadleaf weeds common lambsquarters (*Chenopodium album* L.) and sicklepod [*Senna obtusifolia* (L.) Irwin & Barneby], respectively (Everman et al. 2009a; Pline et al. 1999). Metabolism results observed in this study demonstrate that a mechanism enhancing herbicide metabolism was not involved in evolution of glufosinate resistance in the tested *A. palmeri*.

The overall results of uptake, translocation, and metabolism show that the resistance evolution to glufosinate in *A. palmeri* is not attributable to the non-target site resistance mechanisms investigated in this study. A similar result was reported in glufosinate-resistant *E. indica*, in which the resistant accession had no changes in uptake, translocation, or enhanced metabolism (Jalaludin et al. 2017). In contrast, one glufosinate-resistant *L. perenne* population from Oregon showed increased metabolism compared with the susceptible standard (Brunharo et al. 2019). Another non-target site resistance mechanism is herbicide degradation by glutathione conjugation (Powles and Yu 2010). However, in a previous study conducted with the addition of 4-chloro-7-nitrobenzofurazan (NBD-Cl), a glutathione S-transferase inhibitor, to glufosinate, there was no difference in mortality when resistant *A. palmeri* was treated with only glufosinate or glufosinate plus NBD-Cl (Carvalho-Moore et al. 2021). Other investigations into areas such as the influence of the addition of cytochrome P450 inhibitors or reactive oxygen species accumulation in different biotypes remain to be conducted.

In conclusion, the results obtained strongly indicate that glufosinate resistance in the investigated *A. palmeri* accession from Arkansas is likely a result of increased chloroplastic glutamine synthetase copy number and overexpression. No alterations were observed in the cytosolic glutamine synthetase isoform. There was no change observed in absorption, translocation, or metabolism;

therefore, it can be concluded that these three non-target site resistance mechanisms do not influence the glufosinate resistance level in the resistant accession assessed in this study. Future efforts will focus on the heritability of this mechanism across generations, correlation between gene expression and enzyme activity, and alternative control methods targeting this problematic accession. Along with resistance mechanism investigations, glufosinate screenings with *A. palmeri* field accessions that have survived one or more applications of glufosinate have been conducted yearly for the past 4 yr at the University of Arkansas System Division of Agriculture. These screenings aim to identify other potentially glufosinate-resistant accessions and provide farmers rapid identification of problematic areas, consequently minimizing the dissemination of a resistant accession to neighboring fields.

**Supplementary material.** To view supplementary material for this article, please visit <https://doi.org/10.1017/wsc.2022.31>

**Acknowledgments.** Funding for this research was provided by the Arkansas Soybean Promotion Board and BASF Corporation. No conflicts of interest have been declared.

## References

- Anthony N, Unruh T, Ganser D (1998) Duplication of the Rdl GABA receptor subunit gene in an insecticide-resistant aphid, *Myzus persicae*. *Mol Gen Genet* 260:165–175
- Avila-Garcia WV, Mallory-Smith C (2011) Glyphosate-resistant Italian ryegrass (*Lolium perenne*) populations also exhibit resistance to glufosinate. *Weed Sci* 59:305–309
- Avila-Garcia WV, Sanchez-Olguin E, Hulting AG, Mallory-Smith C (2012) Target-site mutation associated with glufosinate resistance in Italian ryegrass (*Lolium perenne* L. ssp. *multiflorum*). *Pest Manag Sci* 68:1248–1254
- Bayer E, Gugel KH, Hägele K, Hagenmaier H, Jessipow S, König WA, Zähler H (1972) Stoffwechselprodukte von Mikroorganismen. 98. Mitteilung. Phosphinothricin und Phosphinothricyl-Alanyl-Alanin. *Helv Chim Acta* 55:224–239
- Beriault JN, Horsman GP, Devine MD (1999) Phloem transport of D, L-glufosinate and acetyl-L-glufosinate in glufosinate-resistant and-susceptible *Brassica napus*. *Plant Physiol* 121:619–628

- Bernard SM, Habash DZ (2009) The importance of cytosolic glutamine synthetase in nitrogen assimilation and recycling. *New Phytol* 182:608–620
- Bijman J (2001) AgrEvo: From crop protection to crop production. *AgBioForum* 4:20–25
- Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120.
- Brugiere N, Dubois F, Masclaux C, Sangwan RS, Hirel B (2000) Immunolocalization of glutamine synthetase in senescing tobacco (*Nicotiana tabacum* L.) leaves suggests that ammonia assimilation is progressively shifted to the mesophyll cytosol. *Planta* 211:519–527
- Brunharo CA, Takano HK, Mallory-Smith CA, Dayan FE, Hanson BD (2019) Role of glutamine synthetase isogenes and herbicide metabolism in the mechanism of resistance to glufosinate in *Lolium perenne* L. spp. *multiflorum* biotypes from Oregon. *J Agric Food Chem* 67:8431–8440
- Carvalho-Moore P, Norsworthy JK, Gonzalez-Torralva F, Barber T, Piveta LB, Zaccaro-Gruener (2021) Does the addition of glutathione S-transferase inhibitor overcome glufosinate resistance in Palmer amaranth? Page 6 in *Proceedings of the Arkansas Crop Protection Association Conference*. Fayetteville: Arkansas Crop Protection Association
- Castro-Rodríguez V, García-Gutiérrez A, Canales J, Avila C, Kirby EG, Cánovas FM (2011) The glutamine synthetase gene family in *Populus*. *BMC Plant Biol* 11:1–16
- Cattel J, Haberkorn C, Laporte F, Gaude T, Cumer T, Renaud J, Sutherland IW, Hertz JC, Bonneville JM, Arnaud V, Fustec B, Boyer S, Marcombe S, David JF (2021) A genomic amplification affecting a carboxylesterase gene cluster confers organophosphate resistance in the mosquito *Aedes aegypti*: from genomic characterization to high-throughput field detection. *Evol Appl* 14:1009–1022
- Chen J, Huang H, Zhang C, Wei S, Huang Z, Chen J, Wang X (2015) Mutations and amplification of *EPSPS* gene confer resistance to glyphosate in goosegrass (*Eleusine indica*). *Planta* 242:859–868
- Délye C, Jasieniuk M, Le Corre V (2013) Deciphering the evolution of herbicide resistance in weeds. *Trends Genet* 29:649–658
- Edwards JW, Walker EL, Coruzzi GM (1990) Cell-specific expression in transgenic plants reveals nonoverlapping roles for chloroplast and cytosolic glutamine synthetase. *Proc Natl Acad Sci USA* 87:3459–3463
- Elmore MH, McGary KL, Wisecaver JH, Slot JC, Geiser DM, Sink S, O'Donnell K, Rokas A (2015) Clustering of two genes putatively involved in cyanate detoxification evolved recently and independently in multiple fungal lineages. *Genome Biol Evol* 7:789–800
- Everman WJ, Mayhew CR, Burton JD, York AC, Wilcut JW (2009a) Absorption, translocation, and metabolism of  $^{14}\text{C}$ -glufosinate in glufosinate-resistant corn, goosegrass (*Eleusine indica*), large crabgrass (*Digitaria sanguinalis*), and sicklepod (*Senna obtusifolia*). *Weed Sci* 57:1–5
- Everman WJ, Thomas WE, Burton JD, York AC, Wilcut JW (2009b) Absorption, translocation, and metabolism of glufosinate in transgenic and nontransgenic cotton, Palmer amaranth (*Amaranthus palmeri*), and pitted morningglory (*Ipomoea lacunosa*). *Weed Sci* 57:357–361
- Gaines TA, Barker AL, Patterson EL, Westra EP, Wilson RG (2016) EPSPS gene copy number and whole-plant glyphosate resistance level in *Kochia scoparia*. *PLoS ONE* 11:e0168295
- Gaines TA, Zhang W, Wang D, Bukun B, Chisholm ST, Shaner DL, Nissen SJ, Patzoldt WL, Tranel PJ, Culpepper AS, Grey TL, Webster TM, Vencill WK, Sammons RD, Jiang J, et al. (2010) Gene amplification confers glyphosate resistance in *Amaranthus palmeri*. *Proc Natl Acad Sci USA* 107:1029–1034
- Ghanizadeh H, Harrington KC, James TK (2015) Glyphosate-resistant *Lolium multiflorum* and *Lolium perenne* populations from New Zealand are also resistant to glufosinate and amitrole. *Crop Prot* 78:1–4
- Ghoshroy S, Binder M, Tartar A, Robertson DL (2010) Molecular evolution of glutamine synthetase II: phylogenetic evidence of a non-endosymbiotic gene transfer event early in plant evolution. *BMC Evol Biol* 10:1–12
- Godar AS, Stahlman PW, Jugulam M, Dille JA (2015) Glyphosate-resistant kochia (*Kochia scoparia*) in Kansas: EPSPS gene copy number in relation to resistance levels. *Weed Sci*. 63:587–595
- González-Torralva F, Norsworthy JK (2021) Understanding resistance mechanisms to trifluralin in an Arkansas Palmer amaranth population. *Genes* 12:1225
- Habash DZ, Massiah AJ, Rong HL, Wallsgrove RM, Leigh RA (2001) The role of cytosolic glutamine synthetase in wheat. *Ann Appl Biol* 138:83–89
- Heap I (2005) Criteria for confirmation of herbicide-resistant weeds. International Survey of Herbicide-Resistant Weeds. <https://hracglobal.com/herbicide-resistance/confirming-resistance>. Accessed: February 23, 2022
- Heap I (2022a) cDNA Sequence for GS in *Amaranthus palmeri*. The International Herbicide-Resistant Weed Database. [roadsideweeds.com/sequence/sequencedetails.aspx?SequenceID=125](https://roadsideweeds.com/sequence/sequencedetails.aspx?SequenceID=125). Accessed: February 18, 2022
- Heap I (2022b) The International Herbicide-Resistant Weed Database. [www.weedsdatabase.org](https://weedsdatabase.org). Accessed: February 18, 2022
- Heckel DG (2022) Perspectives on gene copy number variation and pesticide resistance. *Pest Manag Sci* 78:12–18
- Hoerlein G (1994) Glufosinate (phosphinothricin), a natural amino acid with unexpected herbicidal properties. *Rev Environ Contam Toxicol* 138:73–145
- Jalaludin A, Ngim J, Bakar BH, Alias Z (2010) Preliminary findings of potentially resistant goosegrass (*Eleusine indica*) to glufosinate-ammonium in Malaysia. *Weed Biol Manag* 10:256–260
- Jalaludin A, Yu Q, Zoellner P, Beffa R, Powles SB (2017) Characterisation of glufosinate resistance mechanisms in *Eleusine indica*. *Pest Manag Sci* 73:1091–1100
- Kamachi K, Yamaya T, Mae T, Ojima K (1991) A role for glutamine synthetase in the remobilization of leaf nitrogen during natural senescence in rice leaves. *Plant Physiol* 96:411–417
- Küpper A, Peter F, Zöllner P, Lorentz L, Tranel PJ, Beffa R, Gaines TA (2018) Tembotrione detoxification in 4-hydroxyphenylpyruvate dioxygenase (HPPD) inhibitor-resistant Palmer amaranth (*Amaranthus palmeri* S. Wats.). *Pest Manag Sci* 74:2325–2334
- Laforest M, Soufiane B, Simard MJ, Obeid K, Page E, Nurse RE (2017) Acetyl-CoA carboxylase overexpression in herbicide-resistant large crabgrass (*Digitaria sanguinalis*). *Pest Manag Sci* 73:2227–2235
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta\text{CT}}$  method. *Methods* 25:402–408
- Mann AF, Fentem PA, Stewart GR (1979) Identification of two forms of glutamine synthetase in barley (*Hordeum vulgare*). *Biochem Biophys Res Commun* 88:515–521
- McNally SF, Hirel B, Gadal P, Mann AF, Stewart GR (1983) Glutamine synthetases of higher plants: evidence for a specific isoform content related to their possible physiological role and their compartmentation within the leaf. *Plant Physiol* 72:22–25
- Meyer CJ, Peter F, Norsworthy JK, Beffa R (2020) Uptake, translocation, and metabolism of glyphosate, glufosinate, and dicamba mixtures in *Echinochloa crus-galli* and *Amaranthus palmeri*. *Pest Manag Sci* 76:3078–3087
- Mifflin BJ, Habash DZ (2002) The role of glutamine synthetase and glutamate dehydrogenase in nitrogen assimilation and possibilities for improvement in the nitrogen utilization of crops. *J Exp Bot* 53:979–987
- Norsworthy JK, Ward SM, Shaw DR, Llewellyn RS, Nichols RL, Webster TM, Bradley KW, Frisvold G, Powles SB, Burgos NR, Witt WW, Barrett M (2012) Reducing the risks of herbicide resistance: best management practices and recommendations. *Weed Sci* 60 (SP1):31–62
- Pline WA, Wu J, Hatzios KK (1999) Absorption, translocation, and metabolism of glufosinate in five weed species as influenced by ammonium sulfate and pelargonic acid. *Weed Sci* 47:636–643
- Powles SB, Yu Q (2010) Evolution in action: plants resistant to herbicides. *Annu Rev Plant Biol* 61:317–347
- Priess GL, Norsworthy JK, Godara N, Mauromoustakos A, Butts TR, Roberts TL, Barber T (2022) Confirmation of glufosinate-resistant Palmer amaranth and response to other herbicides. *Weed Technol*. doi:10.1017/wet.2022.21
- Puinean AM, Foster SP, Oliphant L, Denholm I, Field LM, Millar NS, Williamson MS, Bass C (2010) Amplification of a cytochrome P450 gene is associated with resistance to neonicotinoid insecticides in the aphid *Myzus persicae*. *PLoS Genetics* 6:e1000999
- Ribeiro DN, Pan Z, Duke SO, Nandula VK, Baldwin BS, Shaw DR, Dayan FE (2014) Involvement of facultative apomixis in inheritance of EPSPS gene amplification in glyphosate-resistant *Amaranthus palmeri*. *Planta* 239:199–212

- Salas RA, Dayan FE, Pan Z, Watson SB, Dickson JW, Scott RC, Burgos NR (2012) *EPSPS* gene amplification in glyphosate-resistant Italian ryegrass (*Lolium perenne* ssp. *multiflorum*) from Arkansas. *Pest Manag Sci* 68:1223–1230
- Seabra AR, Vieira CP, Cullimore JV, Carvalho HG (2010) *Medicago truncatula* contains a second gene encoding a plastid located glutamine synthetase exclusively expressed in developing seeds. *BMC Plant Biol* 10:1–16
- Seng CT, Lun LV, San CT, Sahid IB (2010) Initial report of glufosinate and paraquat multiple resistance that evolved in a biotype of goosegrass (*Eleusine indica*) in Malaysia. *Weed Biol Manag* 10:229–233
- Singh S, Singh V, Lawton-Rauh A, Bagavathiannan MV, Roma-Burgos N (2018) *EPSPS* gene amplification primarily confers glyphosate resistance among Arkansas Palmer amaranth (*Amaranthus palmeri*) populations. *Weed Sci* 66:293–300
- Steckel GJ, Hart SE, Wax LM (1997) Absorption and translocation of glufosinate on four weed species. *Weed Sci* 45:378–381
- Takano HK, Beffa R, Preston C, Westra P, Dayan FE (2019) Reactive oxygen species trigger the fast action of glufosinate. *Planta* 249:1837–1849
- Takano HK, Beffa R, Preston C, Westra P, Dayan FE (2020) A novel insight into the mode of action of glufosinate: how reactive oxygen species are formed. *Photosynth Res* 144:361–372
- Takano HK, Dayan FE (2021) Biochemical basis for the time-of-day effect on glufosinate efficacy against *Amaranthus palmeri*. *Plants* 10:1–9
- Travlos I, Cheimona N, De Prado R, Jhala A, Chachalis D, Tani E (2018) First case of glufosinate-resistant rigid ryegrass (*Lolium rigidum* Gaud.) in Greece. *Agronomy* 8:35
- Wallsgrave RM, Turner JC, Hall NP, Kendall AC, Bright SW (1987) Barley mutants lacking chloroplast glutamine synthetase—biochemical and genetic analysis. *Plant Physiol* 83:155–158
- Wendler C, Barniske M, Wild A (1990) Effect of phosphinothricin (glufosinate) on photosynthesis and photorespiration of C3 and C4 plants. *Photosynth Res* 24:55–61
- Wild A, Manderscheid R (1984) The effect of phosphinothricin on the assimilation of ammonia in plants. *Z Naturforsch C* 39:500–504
- Woo KC, Morot-Gaudry JF, Summons RE, Osmond CB (1982) Evidence for the glutamine synthetase/glutamate synthase pathway during the photorespiratory nitrogen cycle in spinach leaves. *Plant Physiol* 70: 1514–1517
- Zhang C, Yu Q, Han H, Yu C, Nyporko A, Tian X, Beckie H, Powles S (2022) A naturally evolved mutation (Ser59Gly) in glutamine synthetase confers glufosinate resistance in plants. *J Exp Bot* 73:2251–2262