



In-Field Assessment of EPSPS Amplification on Fitness Cost in Mixed Glyphosate-Resistant and Glyphosate-Sensitive Populations of Palmer Amaranth (*Amaranthus palmeri*)

Authors: Cahoon, Charles W., Jordan, David L., Tranel, Patrick J., York, Alan C., Riggins, Chance, et al.

Source: Weed Science, 70(6) : 663-668

Published By: Weed Science Society of America

URL: <https://doi.org/10.1017/wsc.2022.60>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

In-field assessment of *EPSPS* amplification on fitness cost in mixed glyphosate-resistant and glyphosate-sensitive populations of Palmer amaranth (*Amaranthus palmeri*)

Research Article

Cite this article: Cahoon CW, Jordan DL, Tranel PJ, York AC, Riggins C, Seagroves R, Inman M, Everman W, Leon R (2022) In-field assessment of *EPSPS* amplification on fitness cost in mixed glyphosate-resistant and glyphosate-sensitive populations of Palmer amaranth (*Amaranthus palmeri*). *Weed Sci.* **70**: 663–668. doi: [10.1017/wsc.2022.60](https://doi.org/10.1017/wsc.2022.60)

Received: 9 May 2022

Revised: 20 September 2022

Accepted: 20 September 2022

First published online: 24 October 2022

Associate Editor:










Ian Burke, Washington State University

Keywords:

EPSPS amplification; resistance

Author for correspondence:

Charles W. Cahoon, Department of Crop and Soil Sciences, North Carolina State University, Campus Box 7620, Raleigh, NC 27695. Email: cwcahoo@ncsu.edu

Charles W. Cahoon¹ , David L. Jordan² , Patrick J. Tranel³ , Alan C. York⁴ , Chance Riggins⁵ , Richard Seagroves⁶ , Matthew Inman⁷ , Wesley Everman¹  and Ramon Leon⁸ 

¹Associate Professor, Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC, USA;

²William Neal Reynolds Professor, Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC, USA;

³Ainsworth Professor and Associate Head, Department of Crop Sciences, University of Illinois, Urbana, IL, USA;

⁴William Neal Reynolds Professor Emeritus, Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC, USA;

⁵Research Assistant Professor, Department of Crop Sciences, University of Illinois, Urbana, IL, USA;

⁶Research Specialist, Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC, USA;

⁷Graduate Research Assistant, Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC, USA and

⁸Professor and University Faculty Scholar, Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC, USA

Abstract

Comparing fitness of herbicide-resistant and herbicide-susceptible weed biotypes is important for managing herbicide resistance. Previous research suggests there is little to no fitness penalty from amplification of the 5-enolpyruvylshikimate-3-phosphate synthase (*EPSPS*) gene (a mechanism of glyphosate resistance) in Palmer amaranth (*Amaranthus palmeri* S. Watson) in controlled studies in the greenhouse or growth chamber. A field study was conducted in North Carolina at three locations naturally infested with *A. palmeri* to determine vegetative, reproductive, and germination fitness of plants with and without *EPSPS* amplification grown season-long with cotton (*Gossypium hirsutum* L.). Seed number was not correlated with *EPSPS* copy number. However, when plants were binned into two groups, those having an *EPSPS* copy number ≥ 2 (relative to reference genes) and those having an *EPSPS* copy number < 2 , plant fresh weight and seed number were 1.4 and 1.6 times greater, respectively, for plants with fewer than 2 *EPSPS* copies. *Amaranthus palmeri* height and seed germination, and yield of cotton, did not differ when comparing the two binned groups. These data suggest that *A. palmeri* plants with *EPSPS* amplification are relatively less fit in the absence of glyphosate, but this reduced fitness does not translate into differences in interference with cotton.

Introduction

Glyphosate resistance in Palmer amaranth (*Amaranthus palmeri* S. Watson) was first confirmed in Georgia in 2005 (Culpepper et al. 2006) and continues to threaten numerous crops in the United States (Heap 2022; Webster 2013). Characteristics of *A. palmeri* that contribute to its ability to dominate fields include high photosynthetic capacity via the C_4 photosynthetic pathway (Ehleringer 1983), rapid growth (Horak and Loughin 2000; Sellers et al. 2003), drought tolerance mechanisms (Place et al. 2008; Wright et al. 1999), shade-adaptive capabilities (Jha and Norsworthy 2009; Jha et al. 2008), prolonged germination (Steckel et al. 2004; Ward et al. 2013), immense fecundity (Schwartz et al. 2016; Webster and Grey 2015), and wide genetic variation (Chandi et al. 2013). These traits give *A. palmeri* a competitive advantage over most crops (Bensch et al. 2003; Burke et al. 2007; MacRae et al. 2013; Massinga et al. 2001; Monks and Oliver 1988). For example, 8 *A. palmeri* plants m^{-1} row are capable of reducing cotton (*Gossypium hirsutum* L.) yield by as much as 92% (MacRae et al. 2013; Morgan et al. 2001; Rowland et al. 1999). When coupled with widespread herbicide resistance, these characteristics cause significant yield loss and increased management costs (Culpepper et al. 2010; Klingaman and Oliver 1994; Morgan et al. 2001).

Amaranthus palmeri biotypes resistant to glyphosate (Group 9) and acetolactate synthase (ALS)-inhibiting (Group 2) herbicides are now commonplace (Heap 2022; Nandula et al. 2012; Poirier et al. 2014; Sosnoskie et al. 2011). The weed has also evolved resistance to triazine (Group 5) and 4-hydroxyphenylpyruvate dioxygenase (HPPD)-inhibiting (Group 27) herbicides (Jhala et al. 2014; Kohrt et al. 2017). More alarming are the recent confirmations of *A. palmeri* biotypes resistant to protoporphyrinogen oxidase (PPO)-inhibiting (Group 14) herbicides, very-long-chain fatty-acid synthesis-inhibiting (Group 15) herbicides, synthetic auxin

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



(Group 4) herbicides, and glufosinate (Group 10) (Brabham et al. 2019; Heap 2022; Salas et al. 2016). Several mechanisms confer glyphosate resistance, including target-site mutation, gene amplification, vacuolar sequestration, reduced cellular absorption, and hypersensitive response (Sammons and Gaines 2014). Specific to gene amplification, *A. palmeri* evolved resistance to glyphosate via amplification of the 5-enolpyruvylshikimate-3-phosphate synthase (*EPSPS*) gene. Researchers who first discovered this mechanism of resistance reported glyphosate-resistant (GR) biotypes from Georgia to have 5- to greater than 160-fold more copies of the *EPSPS* gene than glyphosate-susceptible (GS) biotypes (Gaines et al. 2010).

In some cases, herbicide resistance comes at a penalty, often referred to as a fitness cost. For example, Yannicari et al. (2016) reported GR biotypes of perennial ryegrass (*Lolium perenne* L.) were shorter, had less leaf area and shoot biomass, and produced fewer seeds than GS biotypes. Triazine-resistant smooth pigweed (*Amaranthus hybridus* L.) is less efficient at photosynthesis and is shorter than susceptible wild-type species (Ahrens and Stoller 1983; Jordan 1996). Wu et al. (2018) reported a significant fitness cost in waterhemp [*Amaranthus tuberculatus* (Moq.) Sauer] associated with ALS resistance but not with resistance to atrazine, PPO-inhibiting herbicides, HPPD-inhibiting herbicides, or glyphosate. However, the researchers did note that *A. tuberculatus* resistant to glyphosate via amplification of the *EPSPS* gene decreased in frequency, while glyphosate resistance due to Pro-106-Ser *EPSPS* codon substitution increased over the course of the six generation (3-yr) experiment. Numerous greenhouse experiments have been conducted to determine whether glyphosate resistance imparts a fitness penalty in *A. palmeri*. Giacomini et al. (2014) reported no significant relationship between both final plant biomass and seed production with *EPSPS* copy number, although a negative relationship between *EPSPS* copy number and seed production was noted for one of the four pseudo- F_2 families tested. Similarly, researchers from Argentina and Australia found no correlation between *EPSPS* copy number and plant height or biomass for this weed (Vila-Aiub et al. 2014).

Determination of fitness costs associated with herbicide resistance is challenging due to the need to control for genetic background and because of the potential environmental dependence for manifestation of the fitness costs. (Keshtkar et al. 2019; Vila-Aiub et al. 2011). Furthermore, control of genetic background is necessary to directly relate pleiotropic effects on fitness cost with genes that confer resistance (Bergelson and Purrington 1996; Vila-Aiub et al. 2011). Otherwise, the differences in fitness between a resistant and susceptible biotype might simply be explained by differences in other traits. It is critical to keep in mind that the question about fitness cost due to resistance trait is not just theoretical, and ultimately, it is intended to determine the likelihood of the trait persisting or increasing in agroecosystems. The reality is that under field conditions, and especially in a dioecious species, genome architecture can change constantly, so the genetic context in which the resistance trait acts will likely differ from population to population and even from plant to plant (Leon et al. 2021). Therefore, studying multiple populations can help our understanding of the impact of the herbicide-resistance trait on the species and the range of fitness costs, if they exist, that can influence the spread of resistant biotypes (Bravo et al. 2017, 2018; Keshtkar et al. 2019). Although studies have indicated no fitness penalty associated with *EPSPS* amplification in *A. palmeri* under controlled conditions, fitness costs under field conditions have not been

documented. The objective of this experiment was to study fitness costs of *EPSPS* amplification in *A. palmeri* under field conditions.

Materials and Methods

The experiment was conducted in North Carolina in 2014 near Clayton at the Central Crops Research Station (35.67°N, 78.51°W) and in a grower's field near Mount Olive (35.20°N, 77.96°W). The experiment was also conducted near Rocky Mount, NC, in 2015 at the Upper Coastal Plain Experiment Station (35.89°N, 77.67°W). Soil at Clayton was a Dothan loamy sand (fine-loamy, kaolinitic, thermic Plinthic Kandiudults) with 0.22% humic matter and pH 5.5. Soil at Mount Olive was a Lakeland sand (thermic, coated Typic Quartzipsammits) with 0.41% humic matter and pH 5.1. A Goldsboro fine sandy loam (fine-loamy, siliceous, subactive, thermic Aquic Paleudults) with 0.56% humic matter and pH 6.2 was present at Rocky Mount. Cotton cultivar 'Stoneville 4946GLB2' (Bayer CropScience, Research Triangle Park, NC, USA) was planted on May 3, 2014, and May 12, 2014, in Mount Olive and Clayton, respectively. At Rocky Mount, cotton cultivar 'Dyna-Gro 3385B2XF' (Loveland Products, Loveland, CO, USA) was planted on May 6, 2015. Cotton was seeded into conventionally tilled raised beds in Clayton and Rocky Mount and a strip-tillage system (Edmisten et al. 2022) in a desiccated wheat (*Triticum aestivum* L.) cover crop in Mount Olive. Cotton was planted at a seeding rate designed to achieve in-row populations of 15 plants m^{-2} with row spacing of 97 cm in Clayton and Mount Olive and 91 cm in Rocky Mount.

Plant Material

Each field site was dedicated to weed science research and naturally infested with a segregating field population of *A. palmeri* at densities greater than 100 plant m^{-2} and a long history (>15 yr) of recurrent glyphosate use. Expression of glyphosate resistance varied across populations and was approximately 40% to 60% at Clayton and Rocky Mount and 67% to 80% at Mount Olive. Level of glyphosate resistance was estimated from the population's response to glyphosate in previous experiments at each field site. As mentioned previously, control of genetic background is necessary to associate fitness cost with genes that confer resistance (Bergelson and Purrington 1996; Vila-Aiub et al. 2011); genetic background in this experiment was controlled by utilizing these three field sites naturally infested with *A. palmeri* segregating for *EPSPS* amplification.

One hundred to 175 *A. palmeri* plants per population were randomly selected at 3 wk after cotton planting and were covered with plastic cups to allow a broadcast application of glufosinate (Liberty® 280 herbicide, Bayer CropScience) at 656 g ai ha^{-1} plus acetochlor (Warrant® herbicide, Monsanto, St Louis, MO, USA) at 1,260 g ai ha^{-1} to control all other weeds. Cups were removed within 30 min of application. Other than the selected *A. palmeri* plants, weeds were removed by hand weeding throughout the season. Experimental fields were approximately 0.2 to 0.4 ha in size. The final plant density of *A. palmeri* was 1 plant 19 m^{-2} , which constituted a single plot; plots were spaced approximately 2.6 m apart. At 6 wk after planting, a single recently emerged leaf, approximately 1 cm^{-2} in size, was removed from each *A. palmeri* plant and stored on dry ice before being shipped to the University of Illinois to determine *EPSPS* copy number relative to a 1-copy reference gene (*CPS*, which encodes the large subunit of

carbamoylphosphate synthetase) number using quantitative PCR as described by Chatham et al. (2015). *Amaranthus palmeri* plants with an *EPSPS* copy number <2 were considered GS, whereas plants with *EPSPS* copy number ≥ 2 were considered GR.

Amaranthus palmeri height was recorded at 15 wk after planting, with aboveground fresh weight determined at physiological maturity in mid-September of each year before seed rain. Cotton yield within 1 m of the *A. palmeri* plant and between 1 and 2 m from the *A. palmeri* plant was determined by hand picking cotton when bolls were fully open in October or November. *Amaranthus palmeri* reproductive inflorescences from female plants were removed, dried, and threshed to determine seed production.

Germination Assay

Seed production per plant was determined by counting seeds in 0.06- to 0.081-g subsamples to calculate 100-seed weight of the cleaned seed. One hundred seeds from each female plant were seeded in separate 25 by 53 cm greenhouse trays containing potting soil mix (Sun Gro® Fafard 4P Mix, Agawam, MA, USA). The greenhouse was maintained at 35 ± 5 C with metal-halide lighting ($400 \mu\text{mol m}^{-2} \text{s}^{-1}$; Hubbell Lighting, Greenville, SC, USA) supplementing natural light for 14 h daily; greenhouse trays were lightly irrigated twice daily. To determine germination percentage, the number of emerged *A. palmeri* seedlings was recorded at 14 d after seeding.

Statistical Analysis

Data for *EPSPS* copy number and *A. palmeri* height, plant biomass, seed number, and germination percentage were subjected to ANOVA using PROC GLIMMIX in SAS (SAS Software v. 9.4, SAS Institute, Cary, NC, USA) to test for main effect of population, glyphosate-resistance binary designation (yes or no), and the interaction of population and resistance designation. Seed cotton yield within 1 m of the weed and from 1 to 2 m away from the weed was also subjected to ANOVA using PROC GLIMMIX in SAS along with cotton yield from a weed-free section. Significance between data for GR and GS biotypes was determined using a standard *t*-test at $P \leq 0.05$. Differences in cotton yield in presence of GR and GS biotypes and the weed-free control were separated using Fisher's protected LSD test at $P \leq 0.05$. Regression coefficients were constructed using the PROC REG in SAS for the following relationships: *EPSPS* copy number versus plant height, *EPSPS* copy number versus biomass, *EPSPS* copy number versus seed number, and *EPSPS* copy number versus seed germination at $P \leq 0.05$. Additional regression coefficients included seed production versus plant height, seed number versus biomass, and plant height versus biomass.

Results and Discussion

Average *EPSPS* copy number for *A. palmeri* plants with ≥ 2 *EPSPS* copies was 41.3, 35.9, and 36.2 at Mount Olive, Clayton, and Rocky Mount, respectively (Table 1). *EPSPS* copy number for *A. palmeri* plants with <2 *EPSPS* copies from these respective populations was 0.8, 0.5, and 0.7, respectively. Fewer female *A. palmeri* plants having <2 *EPSPS* copies relative to females having ≥ 2 *EPSPS* copies were present at the end of the season at Mount Olive, while roughly the same number were present at Clayton and Rocky Mount (Table 1). During the period of time when these experiments were conducted, fields often had varying frequencies of GR *A. palmeri*.

Table 1. Number of female *Amaranthus palmeri* plants and average *EPSPS* copy number and range at Mount Olive, Clayton, and Rocky Mount, NC, USA.

Population	Binary designation	Average <i>EPSPS</i> copy number	Range of <i>EPSPS</i> copy number	Number of plants
Mount Olive	≥ 2 <i>EPSPS</i> copies	41.3	6.9–128.0	13
Mount Olive	<2 <i>EPSPS</i> copies	0.8	0.3–1.3	3
Clayton	≥ 2 <i>EPSPS</i> copies	35.9	3.9–56.1	9
Clayton	<2 <i>EPSPS</i> copies	0.5	0.3–0.7	7
Rocky Mount	≥ 2 <i>EPSPS</i> copies	36.2	21.6–47.4	7
Rocky Mount	<2 <i>EPSPS</i> copies	0.7	0.2–1.2	7

Table 2. Regression coefficients for *Amaranthus palmeri* *EPSPS* copy number, height at 15 wk after planting, plant fresh weight at physiological maturity, seed production, and germination.^a

Correlation variables	P > F	Regression coefficient
<i>EPSPS</i> copy number vs. <i>A. palmeri</i> height	0.2721	0.04
<i>EPSPS</i> copy number vs. <i>A. palmeri</i> biomass	0.7590	0.01
<i>EPSPS</i> copy number vs. <i>A. palmeri</i> seed number	0.8184	-0.01
<i>EPSPS</i> copy number vs. <i>A. palmeri</i> seed germination	0.3374	0.04
<i>A. palmeri</i> height vs. <i>A. palmeri</i> biomass	0.9199	0.01
<i>A. palmeri</i> height vs. <i>A. palmeri</i> seed number	0.9643	0.01
<i>A. palmeri</i> biomass vs. <i>A. palmeri</i> seed number	<0.0001	0.53

^aData are pooled over three populations.

EPSPS copy number was not correlated with *A. palmeri* height, plant biomass, seed number, or seed germination (Table 2). These results are consistent with previous research (Giacomini et al. 2014; Vila-Aiub et al. 2014) showing no relationship between *EPSPS* copy number and biology of this weed, including seed production. While *A. palmeri* height was not correlated with plant biomass or seed number, plant biomass and seed number were correlated (Table 2).

We considered the lack of correlation between *EPSPS* copy number and fitness-associated traits may have occurred because *EPSPS* amplification affects those traits more in a qualitative rather than a quantitative fashion. Stated differently, we hypothesized that plants with, for example, 10 or 100 *EPSPS* copies would incur fitness costs relative to wild-type plants, but the relative fitness among plants expressing increased copy number would be similar. Therefore, we treated *EPSPS* copy number as a binary variable, with those having an *EPSPS* copy number ≥ 2 (relative to reference genes) and those having an *EPSPS* copy number <2 making up the two groups. Analyzed this way, the main effect of *EPSPS* copy number was significant for *A. palmeri* plant biomass and seed number (Table 3). The interaction of *EPSPS* copy number and population was not significant for these measurements. *Amaranthus palmeri* height and seed germination were not affected by *EPSPS* copy number or the interaction of *EPSPS* copy number and population.

When pooled over the three populations, *EPSPS* copy number was 37.8 for plants having ≥ 2 *EPSPS* copies and 0.6 for plants having <2 *EPSPS* copies (Table 4). *Amaranthus palmeri* height for plants having ≥ 2 *EPSPS* copies (141 cm) and plants having <2 *EPSPS* copies (144 cm) were not different. *Amaranthus palmeri* plant biomass and seed number were 1.4 and 1.6 times greater for plants having <2 *EPSPS* copies ($n = 17$) compared with plants

Table 3. Analysis of variance for *EPSPS* copy number, height at 15 wk after planting, biomass, number of seed, and germination of seed for *Amaranthus palmeri* plants based on binary designation of glyphosate resistance.

Source of variation	Degrees of freedom	<i>EPSPS</i> copy number		Height		Biomass		Seed number		Germination	
		F-value	P > F	F-value	P > F	F-value	P > F	F-value	P > F	F-value	P > F
Population	2	0.1	0.9221	100.9	<0.0001	2.1	0.1380	1.8	0.1844	0.1	0.7114
Binary designation ^a	1	38.7	<0.0001	0.1	0.7305	4.2	0.0478	6.4	0.0157	1.4	0.2502
Population × binary designation	2	0.1	0.9333	0.5	0.6277	0.3	0.7238	0.5	0.6268	1.0	0.3282
Error	40	—	—	—	—	—	—	—	—	—	—
Coefficient of variation (%)		72.4	—	16.8	—	56.7	—	61.5	—	33.5	—

^a*EPSPS* copy number ≥ 2 or *EPSPS* copy number < 2.

Table 4. *EPSPS* copy number, height at 15 wk after planting, plant biomass, seed production, and seed germination for *Amaranthus palmeri* plants based on binary designation of glyphosate resistance.^a

Binary designation ^b	<i>EPSPS</i> copy number ^b	Height	Biomass	Seed number	Seed germination
	no. plant ⁻¹	cm	g plant ⁻¹	no. plant ⁻¹	%
<i>EPSPS</i> copy number ≥ 2	37.8 (4.1)	141 (10)	4,503 (350)	342,175 (38,610)	37 (3)
<i>EPSPS</i> copy number < 2	0.6 (0.1)*	144 (16)	6,420 (1,020)*	557,543 (131,140)*	31 (2)

^aSEs for means in parentheses.

^bData are pooled over three populations.

*Significance at P < 0.05 using a paired t-test.

Table 5. Cotton yield with season-long interference from *Amaranthus palmeri* plants based on binary designation of glyphosate resistance in sections 0 to 1 m and 1 to 2 m from the weed.

Binary designation ^a	Cotton yield ^b	
	0 to 1 m from the weed ^c	1 to 2 m from the weed
	g m ⁻¹	
<i>EPSPS</i> copy number ≥ 2	121 (13) b	168 (27) b
<i>EPSPS</i> copy number < 2	115 (23) b	162 (23) b
Weed free	191 (21) a	191 (21) a

^aData are pooled over three populations.

^bSEs for means in parentheses.

^cMeans within a column followed by the same letter are not significantly different based on Fisher's Protected LSD test at P ≤ 0.05.

having ≥ 2 copies ($n = 29$). Seed size was similar for the two *A. palmeri* biotypes, with 100-seed weights of 32.9 and 36.4 mg for plants having < 2 *EPSPS* copies and plants having ≥ 2 copies, respectively (data not shown). Seed germination did not differ for *A. palmeri* plants having < 2 *EPSPS* copies or those having ≥ 2 copies, and no difference in cotton yield was observed after season-long interference with the two biotypes regardless of distance from the *A. palmeri* plant (Table 5). Cotton yield in the presence of a single *A. palmeri* plant was 62% of weed-free cotton within 1 m of the weed and 87% at a distance of 1 to 2 m from the weed.

These data are the only published results in the peer-reviewed literature with a mixed population of *A. palmeri* segregating for *EPSPS* amplification under field conditions in cotton demonstrating a fitness penalty for *EPSPS* copy number associated with seed production. Evaluation of fitness, especially in dioecious species, is difficult due to the changing genome architecture, which can differ from population to population (Leon et al. 2021). Furthermore, we considered the possibilities of a quantitative or qualitative relationship between fitness cost and *EPSPS* amplification. Because the frequencies of glyphosate-resistance mechanisms changed over time, Wu et al. (2018) hypothesized *A. tuberculatus* plants resistant to glyphosate via Pro-106-Ser *EPSPS* codon substitution were more

fit than biotypes resistant to the herbicide by amplification of the *EPSPS* gene, which suggests a fitness cost may exist. While differences in plant biomass and seed production were noted between plants having *EPSPS* copy number < 2 and those having ≥ 2 copies, interference with cotton and possibly other crops will not be affected by resistance. While there is a significant difference in seed production between plants having *EPSPS* copy number < 2 and those having ≥ 2 copies, the amount of seed produced by plants having ≥ 2 copies remains substantial.

Another important consideration is that in a species such as *A. palmeri* with high genetic variability and obligate outcrossing, the genetic architectures in which the resistance allele(s) operate are also variable within populations. Furthermore, other traits that affect fitness are also under selection, and they could additively (or synergistically/antagonistically) increase or mitigate the fitness change caused by the resistance allele(s) (Leon et al. 2021; Leon and van der Laet 2021). In fact, Bravo et al. (2017, 2018) documented in commercial farms the existence of *A. palmeri* populations in which GR biotypes were taller and produced more biomass than GS biotypes. They also found that the differences in growth were at least partially due to increased nutrient-use efficiency in the former compared with the latter. Later, Leon and van der Laet (2021) provided evidence that those differences were the result of selection pressure in the different production systems from which the populations originated. Therefore, interpreting the presence or absence of fitness costs in *A. palmeri* and other dioecious or outcrossing species with high genetic diversity must be done cautiously, because the final fitness that weeds exhibit in the field is highly determined by genes other than those controlling herbicide resistance.

It is not clear why we did not observe a quantitative relationship between fitness cost and *EPSPS* amplification but did observe a significant fitness cost when *EPSPS* amplification was analyzed as a qualitative trait. Intuitively, one would expect more copies would result in more fitness cost due to greater resource drain on the plant or greater perturbation of the shikimate pathway and, therefore, a quantitative relationship. However, one possibility is that increased copies of *EPSPS* do not confer a fitness cost. Instead, perhaps one

or more genes that are present within the *EPSPS* amplicon and that might play a role in fostering DNA amplification (Molin et al. 2020) act in a qualitative fashion to impart a fitness cost.

A practical goal for determining reproductive fitness costs of herbicide resistance is predicting the utility of herbicide rotation as a herbicide-resistance mitigation strategy. In this study, we did not compare fitness between *A. palmeri* males having <2 *EPSPS* copies and males having ≥ 2 copies. Also, because increased *EPSPS* copy number is not inherited as a single-gene trait (Gaines et al. 2010), it is difficult to extrapolate how our observed fitness cost in *A. palmeri* seed production would decrease the frequency of GR individuals over time if the population was not exposed to glyphosate. However, if one simply assumes that all GR and GS females produce only GR and GS progeny, respectively, and one further assumes no impact by relative male fitness and no seed dormancy, then even after 5 yr, a population starting with 75% GR individuals would still have about 20% GR individuals. With seed dormancy, the decrease in GR frequency would occur even more slowly. However, a return to glyphosate use would dramatically increase glyphosate resistance, but perhaps it could be an effective option for managing a population of GR *A. palmeri* if used in rotation only once every 5 to 10 yr.

Acknowledgments. The authors express sincere appreciation to Jung Eun Song for processing leaf samples via quantitative PCR at the University of Illinois. This research was funded in part by funds administered through the North Carolina Cotton Growers State Support Committee and Cotton Incorporated. No conflicts of interest have been declared.

References

- Ahrens WH, Stoller EW (1983) Competition, growth rate, and CO₂ fixation in triazine-susceptible and -resistant smooth pigweed (*Amaranthus hybridus*). *Weed Sci* 31:438–444
- Bensch CN, Horak MJ, Peterson D (2003) Interference of redroot pigweed (*Amaranthus retroflexus*), Palmer amaranth (*A. palmeri*), and common waterhemp (*A. rudis*) in soybean. *Weed Sci* 51:37–43
- Bergelson J, Purrington CB (1996) Surveying patterns in the cost of resistance in plants. *Am Nat* 148:536–558
- Brabham C., Norsworthy JK, Houston MM, Varanasi VK, Barber T (2019) Confirmation of S-metolachlor resistance in Palmer amaranth (*Amaranthus palmeri*). *Weed Technol* 33:720–726
- Bravo W, Leon RG, Ferrell JA, Mulvaney MJ, Wood CW (2017) Differentiation of life-history traits among Palmer amaranth populations (*Amaranthus palmeri*) and its relation to cropping systems and glyphosate sensitivity. *Weed Sci* 65:339–349
- Bravo W, Leon RG, Ferrell JA, Mulvaney MJ, Wood CW (2018) Evolutionary adaptations of Palmer amaranth (*Amaranthus palmeri*) to nitrogen fertilization and crop rotation history affect morphology and nutrient-use efficiency. *Weed Sci* 66:180–189
- Burke IC, Schroeder M, Thomas WE, Wilcut JW (2007) Palmer amaranth interference and seed production in peanut. *Weed Technol* 21:367–371
- Chandi A, Milla-Lewis SR, Jordan DL, York AC, Burton JD, Zuleta MC, Whitaker JR, Culpepper AS (2013) Use of AFLP markers to assess genetic diversity in Palmer amaranth (*Amaranthus palmeri*) populations from North Carolina and Georgia. *Weed Sci* 61:136–145
- Chatham LA, Bradley KW, Kruger GR, Martin JR, Owen MDK, Peterson DE, Mithila J, Tranel PJ (2015) A multi-state study of the association between glyphosate resistance and EPSPS gene amplification in waterhemp (*Amaranthus tuberculatus*). *Weed Sci* 63:569–577
- Culpepper AS, Grey TL, Vencill WK, Kichler JM, Webster TM, Brown SM, York AC, Davis JW, Hanna WW (2006) Glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) confirmed in Georgia. *Weed Sci* 54:620–626
- Culpepper AS, Webster TM, Sosnoskie LM, York AC (2010) Glyphosate-resistant Palmer amaranth in the United States. Pages 195–212 in Nandula VK, ed. *Glyphosate Resistance in Crops and Weeds: History, Development, and Management*. Hoboken, NJ: Wiley
- Edmisten K, Collins GD, Vann R (2022) Cotton production with conservation tillage. Pages 175–184 in 2022 Cotton Information. Publ. AG-417. Raleigh, NC: North Carolina Cooperative Extension Service
- Ehleringer J (1983) Ecophysiology of *Amaranthus palmeri*, a Sonoran Desert summer annual. *Oecologia* 57:107–112
- Gaines TA, Zhang W, Wang D, Bukun B, Chisholm ST, Shaner DL, Nissen SJ, Patzoldt WL, Tranel PJ, Culpepper AS, Grey TL, Webster TM, Vencill WK, Sammons RD, Jiang J, et al. (2010) Gene amplification confers glyphosate resistance in *Amaranthus palmeri*. *Proc Natl Acad Sci USA* 107:1029–1034
- Giacomini D, Westra P, Ward S (2014) Impact of genetic background in fitness cost studies: an example from glyphosate-resistant Palmer amaranth. *Weed Sci* 62:29–37
- Heap I (2022) The International Herbicide-Resistant Weed Database. <http://www.weedscience.org>. Accessed: March 3, 2022
- Horak MJ, Loughin TM (2000) Growth analysis of four *Amaranthus* species. *Weed Sci* 48:347–355
- Jha P, Norsworthy JK (2009) Soybean canopy and tillage effects on emergence of Palmer amaranth (*Amaranthus palmeri*) from a natural seed bank. *Weed Sci* 57:644–651
- Jha P, Norsworthy JK, Riley MB, Bielenberg DB, Bridges D Jr (2008) Acclimation of Palmer amaranth (*Amaranthus palmeri*) to shading. *Weed Sci* 56:729–734
- Jhala A, Sandell LD, Rana N, Kruger GR, Knezevic SZ (2014) Confirmation and control of triazine and hydroxyphenylpyruvate dioxygenase-inhibiting herbicide-resistant Palmer amaranth (*Amaranthus palmeri*) in Nebraska. *Weed Technol* 28:28–38
- Jordan N (1996) Effects of the triazine-resistance mutation on fitness in *Amaranthus hybridus* (smooth pigweed). *J Appl Ecol* 33:141–150
- Keshkar E, Abdolshahi R, Sasanfar H, Zand E, Beffa R, Dayan FE, Kudsk P (2019) Assessing fitness costs from a herbicide-resistance management perspective: a review and insight. *Weed Sci* 67:137–148
- Klingaman TE, Oliver LR (1994) Palmer amaranth (*Amaranthus palmeri*) interference in soybeans (*Glycine max*). *Weed Sci* 42:523–527
- Kohrt JR, Sprague CL, Nadakuduti SS, Douches D (2017) Confirmation of a three-way (glyphosate, ALS, and atrazine) herbicide-resistant population of Palmer amaranth (*Amaranthus palmeri*) in Michigan. *Weed Sci* 65:327–338
- Leon RG, Dunne JC, Gould F (2021) The role of population and quantitative genetics and modern sequencing technologies to understand evolved herbicide resistance and weed fitness. *Pest Manag Sci* 77:12–21
- Leon RG, van der Laat R (2021) Population and quantitative genetic analyses of life-history trait adaptations in *Amaranthus palmeri* S. Watson. *Weed Res* 61:342–349
- MacRae AW, Webster TM, Sosnoskie LM, Culpepper AS, Kichler JM (2013) Cotton yield loss potential in response to length of Palmer amaranth (*Amaranthus palmeri*) interference. *J Cotton Sci* 17:227–232
- Massinga RA, Currie RS, Horak MJ, Boyer J Jr (2001) Interference of Palmer amaranth in corn. *Weed Sci* 49:202–208
- Molin WT, Yaguchi A, Blenner M, Sasaki CA (2020) The EccDNA replicon: a heritable, extranuclear vehicle that enables gene amplification and glyphosate resistance in *Amaranthus palmeri*. *Plant Cell* 32:2132–2140
- Monks DM, Oliver LR (1988) Interactions between soybean (*Glycine max*) cultivars and selected weeds. *Weed Sci* 36:770–774
- Morgan GD, Baumann PA, Chandler JM (2001) Competitive impact of Palmer amaranth (*Amaranthus palmeri*) on cotton (*Gossypium hirsutum*) development and yield. *Weed Technol* 15:408–412
- Nandula VK, Reddy KN, Koger CH, Poston DH, Rimando AM, Duke SO, Bond JA, Ribeiro DN (2012) Multiple resistance to glyphosate and pyriithiobac in Palmer amaranth (*Amaranthus palmeri*) from Mississippi and response to flumiclorac. *Weed Sci* 60:179–188
- Place G, Bowman D, Burton M, Rufy T (2008) Root penetration through a high bulk density soil layer: differential response of a crop and weed species. *Plant Soil* 307:179–190
- Poirier AH, York AC, Jordan DL, Chandi A, Everman WJ, Whitaker JR (2014) Distribution of glyphosate- and thifensulfuron-resistant Palmer amaranth

- (*Amaranthus palmeri*) in North Carolina. *Int J Agron*. doi:10.1155/2014/747810
- Rowland MW, Murray DS, Verhalen LM (1999) Full-season Palmer amaranth (*Amaranthus palmeri*) interference with cotton (*Gossypium hirsutum*). *Weed Sci* 45:305–309
- Salas RA, Burgos NR, Tranel PJ, Singh S, Glasgow L, Scott RC, Nichols RL (2016) Resistance to PPO-inhibiting herbicides in Palmer amaranth from Arkansas. *Pest Manag Sci* 72:864–869
- Sammons RD, Gaines TA (2014) Glyphosate resistance: state of knowledge. *Pest Manag Sci* 70:1367–1377
- Schwartz LM, Norsworthy JK, Young BG, Bradley KW, Kruger GR (2016) Tall waterhemp (*Amaranthus tuberculatus*) and Palmer amaranth (*Amaranthus palmeri*) seed production and retention at soybean maturity. *Weed Technol* 30:284–290
- Sellers BA, Smeda RJ, Johnson WG, Kendig JA, Ellersieck MR (2003) Comparative growth of six *Amaranthus* species in Missouri. *Weed Sci* 51:329–333
- Sosnoskie LM, Kichler JM, Wallace RD, Culpepper AS (2011) Multiple resistance in Palmer amaranth to glyphosate and pyriithiobac confirmed in Georgia. *Weed Technol* 59:321–325
- Steckel LE, Sprague CL, Stoller EW, Wax LM (2004) Temperature effects on germination of nine *Amaranthus* species. *Weed Sci* 52:217–221
- Vila-Aiub MM, Goh SS, Gaines TA, Han H, Busi R, Yu Q, Powles SB (2014) No fitness cost of glyphosate resistance endowed by massive EPSPS gene amplification in *Amaranthus palmeri*. *Planta* 239:793–801
- Vila-Aiub MM, Neve P, Roux F (2011) A unified approach to the estimation and interpretation of resistance costs in plants. *Heredity* 107:386–394
- Ward SM, Webster TM, Steckel LE (2013) Palmer amaranth (*Amaranthus palmeri*): a review. *Weed Technol* 27:12–27
- Webster TM (2013) Weed survey—southern states: broadleaf crops subsection. Pages 275–287 in *Proceedings of the Southern Weed Science Society 67th Annual Meeting*. Las Cruces, NM: Southern Weed Science Society
- Webster TM, Grey TL (2015) Glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) morphology, growth, and seed production in Georgia. *Weed Sci* 63:264–272
- Wright SR, Jennette MW, Coble HD, Ruffly TW (1999) Root morphology of young *Glycine max*, *Senna obtusifolia*, and *Amaranthus palmeri*. *Weed Sci* 47:706–711
- Wu C, Davis AS, Tranel PJ (2018) Limited fitness costs of multiple herbicide-resistance traits in *Amaranthus tuberculatus* facilitate resistance evolution. *Pest Manag Sci* 74:293–301
- Yannicari M, Vila-Aiub M, Istilart C, Acciaresi H, Castro AM (2016) Glyphosate resistance in perennial ryegrass (*Lolium perenne* L.) is associated with a fitness penalty. *Weed Sci* 64:71–79