

# Introduction to a Special Section: Life History Characteristics of Elasmobranch Fishes from the Western North Atlantic Ocean

Authors: Hoffmayer, Eric R., and Sulikowski, James A.

Source: Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science, 5(5): 125-126

Published By: American Fisheries Society

URL: https://doi.org/10.1080/19425120.2013.799619

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science 5:125–126, 2013 © American Fisheries Society 2013 ISSN: 1942-5120 online DOI: 10.1080/19425120.2013.799619

### SPECIAL SECTION: ELASMOBRANCH LIFE HISTORY

# Introduction to a Special Section: Life History Characteristics of Elasmobranch Fishes from the Western North Atlantic Ocean

## Eric R. Hoffmayer and William B. Driggers III

National Marine Fisheries Service, Southeast Fisheries Science Center, Mississippi Laboratories, Post Office Drawer 1207, Pascagoula, Mississippi, USA

#### James A. Sulikowski

Marine Science Department, University of New England, 11 Hills Beach Road, Biddeford, Maine 04005, USA

Humans have harvested elasmobranchs in the waters of the western North Atlantic Ocean since at least the 16th century (Borhegyi 1961). In that time, fisheries for elasmobranchs have waxed and waned. However, it is now widely acknowledged that most elasmobranchs share k-selected life history characteristics, such as relatively slow growth, late maturity, and low fecundity, that make sustainable fisheries possible only through careful management (Walker 1998). In 1993, the National Marine Fisheries Service implemented a fishery management plan for selected shark species inhabiting territorial waters of the United States within the western North Atlantic Ocean (Stone et al. 1998). As a result, stock assessments have been conducted to monitor the status of the managed species. However, accurate biological information, such as reproductive rates, fecundity, and age at maturity, are required for these assessments to be reliable (Cortés 1998).

With the exception of the little information gleaned from the anatomical investigations, developmental studies, and faunal synopses of the late 1800s and early 1900s (e.g., Jordan and Evermann 1896), virtually nothing was known about the life histories of elasmobranchs until the 20th century. In the early to mid-1900s, a small group of biologists, most notably Henry Bigelow, Eugene Gudger, William Schroeder, and Stewart Springer, began reporting life history information from anecdotal sources and direct observations. The first detailed studies providing information on the life history of specific elasmobranch species in the western North Atlantic Ocean appeared in the 1950s (e.g., Springer 1950; Backus et al. 1956). Since that time there has been a progressively growing body of literature on the life histories of elasmobranchs in this region, and we now have at least a rudimentary understanding of the biology for some of the more commonly encountered and/or commercially important species.

Through the generous support of the Mississippi Chapter of the American Fisheries Society, the University of New England, and the University of Southern Mississippi Gulf Coast Research Laboratory, a symposium entitled "Life History Characteristics of Elasmobranch Fishes from the Western North Atlantic Ocean" was held on 28 January 2012 at the 20th meeting of the Southern Division of the American Fisheries Society in Biloxi, Mississippi. Representatives from over 15 universities, federal and state agencies, and private organizations participated in the symposium. Its objective was to provide a forum in which biologists could share recent findings regarding the age, growth, reproduction, and habitat utilization of various elasmobranchs inhabiting waters off the East Coast of the United States and within the northern Gulf of Mexico. What follows are eight contributions to our knowledge of the life histories of elasmobranch fishes in the western North Atlantic Ocean that were presented at the symposium.

We thank our coeditors Walter Bubley (Texas Department of Parks and Wildlife), Ken Goldman (Alaska Fish and Game Commission), and John Mandelman (New England Aquarium) for their efforts. Additionally, we thank the editor-in-chief of *Marine and Coastal Fisheries*, Donald Noakes, for his assistance throughout this process. Finally, we dedicate this special section to the memory of our friend and colleague Loraine F. Hale, who will be widely remembered as one who furthered our understanding of the biology of elasmobranch fishes and who was a pleasure to have known.

#### REFERENCES

Backus, R. H., S. Springer, and E. L. Arnold Jr. 1956. A contribution to the natural history of the White-tip Shark, *Pterolamiops longimanus* (Poey). Deep-Sea Research 3:178–188.

- Borhegyi, S. F. D. 1961. Shark teeth, stingray spines, and shark fishing in ancient Mexico and Central America. Southwestern Journal of Anthropology 17:273–296.
- Cortés, E. 1998. Demographic analysis as an aid in shark stock assessment and management. Fisheries Research 39:199–208.
- Jordan, D. S., and B. W. Evermann. 1896. The fishes of North and Middle America: a descriptive catalogue of the species of fish-like vertebrates found in the waters of North America, north of the isthmus of Panama. U.S. Government Printing Office, Bulletin of the United States National Museum 47, Washington, D.C.
- Springer, S. 1950. Natural history notes on the Lemon Shark, *Negaprion brevirostris*. Texas Journal of Science 3:349–359.
- Stone, R. B., C. M. Bailey, S. A. McLaughlin, P. M. Mace, and M. B Schuze. 1998. Federal management of US Atlantic shark fisheries. Fisheries Research 39:215–221.
- Walker, T. I. 1998. Can shark resources be harvested sustainably? A question revisited with a review of shark fisheries. Marine and Freshwater Research 49:553–572.