

Three new species of the genus *Notophthiracarus* from New Zealand (Acari: Oribatida: Phthiracaridae)

Authors: Liu, Dong, and Zhang, Zhi-Qiang

Source: Systematic and Applied Acarology, 19(2) : 189-196

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.19.2.9>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Three new species of the genus *Notophthiracarus* from New Zealand (Acari: Oribatida: Phthiracaridae)

DONG LIU¹ & ZHI-QIANG ZHANG^{2,3*}

¹ Key Laboratory of Wetland Ecology and Environment, Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun 130102, P. R. China. E-mail: liudong@iga.ac.cn

² Landcare Research, 231 Morrin Road, Auckland, New Zealand.

³ Centre for Biodiversity & Biosecurity, School of Biological Sciences, University of Auckland, Auckland, New Zealand

* Corresponding author. E-mail: zhangz@landcareresearch.co.nz

Abstract

Three New Zealand species of *Notophthiracarus* (Oribatida: Phthiracaridae) are described as new to science: *Notophthiracarus motumuka* **sp. nov.** from the Lady Alice Island, Hen & Chickens Islands, *Notophthiracarus tamaki* **sp. nov.** from the Tamaki Estuary Tohuna Torea, Auckland, and *Notophthiracarus rimi* **sp. nov.** from the Red Island, Mercury Islands. All holotype specimens are deposited at New Zealand Arthropod Collection, Landcare Research and some paratypes are also deposited in Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences.

Key words: Soil mites, Oribatida, Phthiracaridae, *Notophthiracarus*, new species, New Zealand

Introduction

The genus *Notophthiracarus* Ramsay, 1966 is one of the largest genera within the family Phthiracaridae and is widespread in the regions of the world except Nearctic Region (Subías 2014, Niedbała 2002). It is mainly distinguished from others by genital setae arranged in a single row (distance between g_6 and g_5 longer than that between g_5 and g_4 or g_3 and g_4), two setae (an_1 and an_2) near the paraxial margin of ano-adanal plate, and seta d on tibiae IV short and coupled with solenidions (Niedbała 1994). According to Niedbała (2012) and data summarized by various authors, 43% known phthiracarid mites (83 species) belong to the genus *Notophthiracarus* in Australian Region. *Notophthiracarus* is also the most diverse group of phthiracarid species in New Zealand, representing 24 species (Liu & Zhang, 2013). In this paper we report three additional new species of *Notophthiracarus* from New Zealand.

Material and methods

Measurements and descriptions are based on specimens mounted in temporary cavity slides that were studied using a light microscope equipped with a drawing attachment. Terminology generally follows Niedbała (1992, 2000). The unit of measurement is micrometre (μm).

All holotype specimens are deposited in the New Zealand Arthropod Collection, Landcare Research, Auckland (NZAC). Paratype specimens are split between NZAC and Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun (NIGA).

Descriptions of new species

Notophthiracarus motumuka sp. nov.

(Figs. 1–8)

Material examined: Holotype: adult (NZAC, in alcohol, 82/7), New Zealand: ND, Hen & Chickens Is. Lady Alice Island, Main Ridge above Grave Bay, from litter, 1 Jan. 1982, leg. R. Hay. Paratype: one adult (NIGA, in alcohol, 82/7), same data as holotype.

Etymology. Named after the type locality—the Lady Alice Island. Motu Muka is the Māori name for the Lady Alice Island, which is a large one (also known as the Big Chicken) among the Hen & Chicken Islands. It is here used as a noun in apposition.

Description. *Measurements.* Holotype: Prodorsum: length 240, width 155, height 95, setae: ss 38, ro 33, in 85; notogaster: length 428, width 270, height 287; setae: c_1 78, d_1 70, e_1 75, h_1 70, ps_1 63; ventral region: ad_1 90, ad_2 98, ad_3 19, an_1 45, an_2 48; genito-aggenital plate 100×102, ano-adanal plate 80×175. Paratype: Prodorsum: length 250, width 160, height 97; notogaster: length 435, width 275, height 290.

Integument. Colour yellowish. Surface of body covered with small and rounded foveoles with some distances between them.

Prodorsum (Figs. 1–2). Median crista and posterior furrows absent; lateral carinae reaching sinus; sigillar fields distinct, dorsal field narrow, longer than lateral fields; sensilli (ss) with narrow stalk, and fusiform head, covered with small spines; interlamellar setae (in) long, erect and stout, densely covered with small spines in distal half; rostral setae (ro) rough and semi-erect, much thinner than interlamellar setae; lamellar (le) and exobothridial (ex) setae vestigial; comparative length: $in > ss > ro$; mutual distance of setae: $in-in/ro-ro=4$.


Notogaster (Fig. 1). 15 pairs of setae ($c_1/c_1-d_1=0.78$) present, similar in shape with interlamellar setae; setae h_3 shortest and thinnest; setae c_1 and c_3 near anterior border, setae c_2 much further; vestigial setae f_1 positioned posterior to setae h_1 ; three pairs of lyrifissures ia , im and ih present.

Gnathosoma (Figs. 4–6). Subcapitulum normal (Fig. 4); setae h , m , and a simple and smooth; setae h shorter than distance between them; adoral setae typical of family; palp (Fig. 5) 4-segmented, with femur and genu fused; palpal setation: 0-2-2-7(1); supracoxal seta simple and smooth; chelicera (Fig. 6) typical of family.

Ano-genital region (Figs. 1, 3). Genital setae (g) with formula: 5: 4; ano-adanal plates each with five pairs of rough setae (an and ad), setae ad_1 and ad_2 long, thick and hooked distally, setae an_1 and an_2 thinner but straight, setae ad_3 shortest and thinnest, but distinctly longer and thicker than genital setae; comparative length: $ad_2 > ad_1 > an_2 > an_1 > ad_3$.

Legs (Figs. 7–8). Setal counts for leg segments (without tarsi): I: 1-4-2(2)-4(1); II: 1-3-2(1)-3(1), III: 2-2-1(1)-2(1), IV: 2-1-1-2(1); chaetotaxy of legs complete; setae d on femora I inserted at level of setae l'' ; setae a'' on tarsi I and setae ft'' on tarsi II curved distally; setae a'' on tarsi II curved distally; setae s and pv' on tarsi IV present; setae s on tarsi I and II present.

Remark. This new species is more close to *Notophthiracarus repostus* Niedbala, 1989 in sharing the following features: posterior furrows of prodorsum absent, similar shape of sensilli, interlamellar and notogastral setae, lamellar and exobothridial setae vestigial, vestigial setae f_1 positioned posterior to setae h_1 , formula of genital setae: 5: 4, similar shape of setae ad_2 and ad_3 , and chaetotaxy of legs complete. However, the new species can be easily distinguished from the latter species by the following eight characters (a versus b): in *N. motumuka* sp. nov., (1a) lateral carinae of prodorsum present; (2a) dorsal field of prodorsum not bifurcate at distal end, lateral fields longer; (3a) rostral setae relatively short (ro 33), $in-in/ro-ro=4$; (4a) three pairs of lyrifissures ia , im and ips present; (5a) $h < h-h$; (6a) setae ps_4 situated at the level between setae ad_1 and ad_2 ; (7a) setae ad_1



FIGURES 1–8. *Notopthiracarus motumuka* sp. nov.: 1, lateral view of body (legs removed); 2, prodorsum, dorsal view; 3, left side of ventral plate; 4, subcapitulum, palpi removed; 5, palp, antiaxial view; 6, chelicera, antiaxial view; 7, femur I; 8, tibia IV. Scale bars: 1=100 μ m; 2–4, 6–8=50 μ m; 5=25 μ m.

similar as ad_2 , long, thick and hooked distally; (8a) setae d of femora I not bifurcate distally; in *N. repostus*, (1b) lateral carinae of prodorsum absent; (2b) dorsal field of prodorsum bifurcate at distal end, lateral fields shorter; (3b) rostral setae fairly long (ro 66), $in-in/ro-ro \approx 1.39$; (4b) two pairs of lyrifissures ia and im present; (5b) $h > h-h$; (6b) setae ps_4 situated at the level between setae ad_2 and ad_3 ; (7b) setae ad_1 thinner and much shorter than ad_2 , slightly hooked distally; (8b) setae d of femora I bifurcate distally.

***Notophthiracarus tamaki* sp. nov.**

(Figs. 9–19)

Material examined: Holotype: adult (NZAC, in alcohol, 83/166), New Zealand: AK, Tamaki Estuary Tohuna Torea, from litter, 13 Oct. 1983, leg. D. Russell. Paratypes: two adults (NZAC, in alcohol, 83/166), same data as holotype; two adults (NIGA, in alcohol, 83/166), same data as holotype.

Etymology. Named after type locality. It is here used as a noun in apposition.

Description. Measurements. Holotype: Prodorsum: length 330, width 210, height 124, setae: ss 50, ro 13, in 20, le 13, ex 23; notogaster: length 750, width 375, height 435; setae: c_1 13, d_1 15, e_1 23, h_1 18, ps_1 15; ventral region: ad_1 12, an_1 11, g_1 10; genito-aggenital plate 150×195 , ano-adanal plate 105×200 . Paratypes: Prodorsum: length 232–380, width 178–230, height 104–170; notogaster: length 510–858, width 305–440, height 370–550.

Integument. Colour brown. Surface of notogaster with distinct polygonal sculpture, other regions covered with very small and rounded foveoles with some distances between them.

Prodorsum (Figs. 9, 11–12). Median crista developed; posterior furrows present; lateral carinae absent; sigillar fields distinct, dorsal field narrow and bifurcate anteriorly, longer than lateral fields; sensilli (ss) short and fusiform, covered with small spines; other prodorsal setae (in , le , ro , ex) minute and smooth; comparative length: $ss > ex > in > le = ro$; mutual distance of setae: $in-in/ro-ro \approx 2.4$.

Notogaster (Fig. 9–10). Anterior cowl distinct, covered posterior part of prodorsum, not reaching interlamellar setae; one dorsal, longitudinal carinae present, starting from anterior margin and ending at the insertion levels between setae d_1 and e_1 ; it widens anteriorly and gradually narrows to an thick line at insertion level of setae d_1 , then slightly broad posteriorly; 15 pairs of minute setae ($c_1 < 1/10 c_1 - d_1$) present, similar in shape with prodorsal setae; setae c_1 close to anterior border, setae c_{2-3} much more remote than c_1 ; vestigial setae f_1 positioned anterior to setae h_1 ; two pairs of lyrifissures ia and im present.

Gnathosoma (Figs. 15–17). Subcapitulum normal (Fig. 15); setae h , m , and a simple and smooth; setae h more than two times longer than distance between them; adoral setae typical of family; palp (Fig. 16) 4-segmented, with femur and genu fused; palpal setation: 0-2-2-7(1); supracoxal seta simple and smooth; chelicera (Fig. 17) typical of family.

Ano-genital region (Figs. 9, 13–14). Genital setae (g) with formula: 5: 4; ano-adanal plates each with five pairs of minute and fine setae (an and ad).

Legs (Figs. 18–19). Setal counts for leg segments (without tarsi): I: 1-4-2(2)-4(1); II: 1-3-2(1)-3(1), III: 2-2-1(1)-2(1), IV: 2-1-1-2(1); chaetotaxy of legs complete; setae d on femora I inserted at level anterior to setae l'' ; setae a'' on tarsi I and setae ft'' on tarsi II curved distally; setae a'' on tarsi II curved distally; setae s and pv' on tarsi IV present; setae s on tarsi I and II present.

Remark. This new species is most similar to *Notophthiracarus tripartitus* Niedbala, 1989 in sharing the following features: notogaster with anterior cowl and dorsal carinae; median crista of prodorsum present; posterior furrows present; lateral carinae absent; dorsal field of prodorsum bifurcate distally; prodorsal, notogastral, anal and adanal setae short and fine; two pairs of lyrifissures

present; setae *h* much longer than distance between them; formula of genital setae: 5: 4; chaetotaxy of legs complete. However, the new species can be easily distinguished from the latter species by the following five characters (a versus b): in *N. tamaki* **sp. nov.**, (1a) surface of notogaster with distinct polygonal sculpture; (2a) sensilli short and fusiform; (3a) anterior cowl not concave in dorsal view, and small, not reaching interlamellar setae; (4a) notogaster with one short dorsal carina, not same in width longitudinally; (5a) vestigial setae f_1 positioned anterior to setae h_1 ; in *N. tripartitus*, (1b) surface of notogaster foveolate; (2b) sensilli long and lanceolate; (3b) anterior cowl concave in dorsal view, and large, reaching far beyond interlamellar setae; (4b) notogaster with two long dorsal carinae, nearly same in width longitudinally; (5b) vestigial setae f_1 positioned posterior to setae h_1 .

***Notophthiracarus rimi* sp. nov.**

(Figs. 20–29)

Material examined: Holotype: adult (NZAC, in alcohol, 72/227), New Zealand: Red I. Mercury Is. CL., from litter, 24 Nov. 1972, leg. G. W. Ramsay. Paratypes: four adults (NZAC, in alcohol, 72/227), same data as holotype; three adults (NIGA, in alcohol, 72/227), same data as holotype.

Etymology. The specific name (*rimi*) is abbreviation for Red Island, Mercury Islands. It is used here as a noun in apposition.

Description. Measurements. Holotype: Prodorsum: length 255, width 175, height 95, setae: *ss* 35, *ro* 48, *in* 75, *le* 10, *ex* 15; notogaster: length 504, width 320, height 310; setae: c_1 80, d_1 70, e_1 75, h_1 72, ps_1 70, ps_4 46; ventral region: ad_1 50, ad_2 70, ad_3 15, an_1 45, an_2 45; genito-aggenital plate 102×150, ano-adanal plate 95×178. Paratypes: Prodorsum: length 205–225, width 150–155, height 75–80; notogaster: length 370–375, width 255–257, height 245–255.

Integument. Colour yellowish. Surface of body covered with very small and rounded foveoles, especially dense on notogaster.


Prodorsum (Figs. 20–22). Median crista, lateral carinae and posterior furrows absent; sigillar fields distinct, dorsal field narrow, longer than lateral fields; sensilli (*ss*) with narrow stalk, and rounded and rough head; interlamellar setae (*in*) long, erect and stout, sparsely covered with small spines in distal half; rostral setae (*ro*) rough and semi-erect, much thinner than interlamellar setae; lamellar (*le*) and exobothridial (*ex*) setae short and fine; comparative length: $in > ro > ss > ex > le$; in/le 7.5; mutual distance of setae: $in-in/ro-ro \approx 3.73$.

Notogaster (Fig. 20). 15 pairs of setae ($c_1/c_1-d_1=0.68$) present, similar in shape with interlamellar setae; setae e_2 , h_2 and h_3 shortest; setae c_1 and c_3 near anterior border, setae c_2 much further; vestigial setae f_1 positioned anterior to setae h_1 ; two pairs of lyrifissures *ia* and *im* present.

Gnathosoma (Figs. 25–27). Subcapitulum normal (Fig. 25); setae *h*, *m*, and *a* simple and smooth; setae *h* shorter than distance between them; adoral setae typical of family; palp (Fig. 26) 4-segmented, with femur and genu fused; palpal setation: 0-2-2-7(1); supracoxal seta simple and smooth; chelicera (Fig. 27) typical of family.

Ano-genital region (Figs. 20, 23–24). Genital setae (*g*) with formula: 5: 4; ano-adanal plates each with five pairs of setae (*an* and *ad*), setae ad_1 and ad_2 long, thick, rough and slightly hooked, setae ad_3 minute and fine, setae an_1 and an_2 thinner and straight, setae ad_3 short and fine, similar in shape as genital setae; comparative length: $ad_2 > ad_1 > an_1 = an_2 > ad_3$.

Legs (Figs. 28–29). Setal counts for leg segments (without tarsi): I: 1-4-2(2)-4(1); II: 1-3-2(1)-3(1), III: 2-2-1(1)-2(1), IV: 2-1-1-2(1); chaetotaxy of legs complete; setae *d* on femora I inserted in the middle of article and slightly posterior to the level of setae *l''*; setae *a''* on tarsi I and setae *ft''* on tarsi II curved distally; setae *a''* on tarsi II curved distally; setae *s* and *pv'* on tarsi IV present; setae *s* on tarsi I and II present.



FIGURES 20–29. *Notopthiracarus rimi* sp. nov.: 20, lateral view of body (legs removed); 21, prodorsum, dorsal view; 22, sensillus, dorsal view; 23, right side of genito-aggenital plate; 24, right side of ano-adanal plate; 25, subcapitulum, palpi removed; 26, palp, antiaxial view; 27, chelicera, antiaxial view; 28, trochanter and femur I; 29, tibia IV. Scale bars: 20–21, 23–24=100µm; 25–29=50µm; 22=25µm.

Remark. This new species is very close to *Notophthiracarus claviger* Niedbala, 1993 in having the following features: lateral carinae absent, similar shape of prodorsal and notogastral setae, $h < h-h$, setae ad_3 short and fine, and chaetotaxy of legs complete. However, the new species can be easily distinguished from the latter species by the following six characters (a versus b): in *N. rimi* **sp. nov.**, (1a) notogaster elongated oval in shape; (2a) sigillar fields distinct, dorsal field narrower; (3a) setae h_1 inserted below level of e_2 , setae ps_1 situated much below level of h_2 , and setae ps_2 situated slightly below level of ps_3 ; (4a) formula of genital setae: 5: 4; (5a) setae ad_2 situated slightly posterior to the insertion level of an_1 ; (6a) setae d on femora I inserted slightly posterior to the level of setae l'' ; in *N. claviger*, (1b) notogaster rounded in shape; (2b) sigillar fields poorly visible, dorsal field broader; (3b) setae h_1 inserted slightly above level of e_2 , setae ps_1 situated slightly above level of h_2 , and setae ps_2 situated slightly above level of ps_3 ; (4b) formula of genital setae: 6: 3; (5b) setae ad_2 situated slightly posterior to the insertion level of an_2 ; (6b) setae d on femora I inserted much posterior to the level of setae l'' .

Acknowledgements

We thank all the colleagues who collected specimens used in this paper. The first author is very grateful to Prof. Jun Chen (Institute of Zoology, Chinese Academy of Sciences, Beijing, China) for supplying references on oribatid mites. His work was supported by the Funds for The Excellent Youth Scholars of "NEIGAE, CAS" (DLSYQ2012004), the Knowledge Innovation Programs of the Chinese Academy of Sciences (KSCX2-EW-Z-8), the Major Program of National Natural Science Foundation of China—Fauna Sinica (31093430), the key research program of the Chinese Academy of Sciences (Grant No. KZZD-EW-TZ-16), the China Scholarship Council, and the National Natural Science-Foundation of China (Grant No. 31101617). The second author's research on defining New Zealand mites was supported by Core funding for Crown Research Institutes from the Ministry of Business, Innovation and Employment's Science and Innovation Group.

References

- Liu, D. & Zhang, Z.-Q. (2013) The genus *Notophthiracarus* of New Zealand (Acari: Oribatida: Phthiracariidae): three new species and a key to 24 described species. *Zootaxa*, 3682 (2), 392–400. <http://dx.doi.org/10.11646/zootaxa.3682.2.11>
- Niedbala, W. (1989) Phthiracaroida (Acari, Oribatida) nouveaux du royaume australien. *Annales Zoologici*, 43(2), 19–50.
- Niedbala, W. (1992) *Phthiracaroida (Acari, Oribatida). Systematic Studies*. PWN-Polish Scientific Publishers, Warszawa, 612 pp.
- Niedbala, W. (1993) New species of Euptyctima (Acari, Oribatida) from New Zealand. *New Zealand Journal of Zoology*, 20, 137–159. <http://dx.doi.org/10.1080/03014223.1993.10422856>
- Niedbala, W. (1994) Supplement to the classification of Phthiracaroida, with redescriptions and descriptions of some species (Acari, Oribatida, Euptyctima). *Genus*, 5 (1–2), 1–152.
- Niedbala, W. (2000) The ptyctimous mites fauna of the Oriental and Australian Regions and their centres of origin (Acari: Oribatida). *Genus*, supplement, 1–493.
- Niedbala, W. (2002) Ptyctimous mites (Acari, Oribatida) of the Nearctic Region. *Monographs of the Upper Silesian Museum, Bytom*, 4, 1–261.
- Niedbala, W. (2012) An updated study of ptyctimous mite fauna (Acari: Oribatida) of the Australasian Region with a description of thirteen new species. *Acariologia*, 52, 183–228. <http://dx.doi.org/10.1051/acarologia/20122049>
- Suñas, L.S. (2014) Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo (Excepción fósiles). *Graellsia*, 60(número extraordinario), 3–305 (2004) (Actualizado en junio de 2006, en abril de 2007, en mayo de 2008, en abril de 2009, en julio de 2010, en febrero de 2011, en abril de 2012, en mayo de 2013 y en febrero de 2014). [Internet]. [cited 2014 Apr. 29]; Available from: http://escalera.bio.ucm.es/usuarios/bba/cont/docs/RO_1.pdf

Accepted by Qing-Hai Fan: 30 Apr. 2014; published 27 Jun. 2014