

An Analysis of Multiple Stressors on Resident Benthic Communities in a California Agricultural Stream

Authors: Hall, Lenwood W, Anderson, Ronald D, Killen, William D, and Alden, Raymond W

Source: Air, Soil and Water Research, 11(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/1178622118777761>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

An Analysis of Multiple Stressors on Resident Benthic Communities in a California Agricultural Stream

Lenwood W Hall Jr¹, Ronald D Anderson¹, William D Killen¹ and Raymond W Alden III²

¹Agricultural Experiment Station, Wye Research and Education Center, University of Maryland, Queenstown, MD, USA. ²Office of the Provost, Touro University, Henderson, NV, USA.

Air, Soil and Water Research
Volume 11: 1–18
© The Author(s) 2018
Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
DOI: 10.1177/1178622118777761



ABSTRACT: This 3-year study (2015–2017) was designed to characterize benthic communities (macroinvertebrates) and physical habitat in an agriculturally dominated waterbody in the Central Coast area of California (Santa Maria River). Benthic communities as represented by various metrics that represent richness, composition, tolerance/intolerance and trophic measures were used as response variables for the various stressors described below. Concurrent water quality evaluations, physical sediment parameters (grain size and total organic carbon [TOC]), pyrethroids, bulk metals—including simultaneously extracted metals (SEM) and acid volatile sulfides (AVS) ratios—and nutrients were measured. The relationship of various benthic metrics to physical habitat metrics, pyrethroids, metals, nutrients and sediment characteristics was evaluated for the 3-year data set. Total physical habitat scores in this watershed were considered to be poor. Samples collected for various sediment chemistry measurements were from depositional areas (fine grain areas primarily silt and clay) where hydrophobic chemicals such as pyrethroids could be found if sources exist. Dominant benthic taxa collected were generally considered to be tolerant to moderately tolerant of environmental stressors and rated as impaired based on a benthic index. Potentially toxic sediment concentrations of arsenic, cadmium and nickel were reported at various sites based on a comparison with existing threshold effect levels. Pyrethroid concentrations interpreted by using a highly protective toxics units approach with a laboratory sensitive taxon (*Hyalella*) suggested potential toxicity at various sites. Nutrient concentrations could not be interpreted within the context of potential impairment because the State of California has not developed nutrient criteria. The results of the stepwise linear regression models comparing benthic metrics with all environmental variables showed that TOC was the most important variable shaping the benthic communities. In contrast, pyrethroids, metals and physical habitat were not shown to be significant factors shaping benthic communities. The summary multivariate canonical correlation analysis indicated that less stressed, more diverse benthic communities tended to be associated more with TOC-rich finer sediments and lower concentrations of phosphorous-based nutrients, and more stressed, less diverse benthic communities tended to be associated with less organically rich, somewhat less fine sediments and higher phosphorous concentrations.

KEYWORDS: Benthic communities, agricultural stream, pyrethroids, metals, nutrients, physical habitat, sediment characteristics

RECEIVED: November 14, 2017. **ACCEPTED:** April 19, 2018.

TYPE: Original Research

FUNDING: The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was funded by the Pyrethroid Working Group

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Lenwood W Hall Jr, Agricultural Experiment Station, Wye Research and Education Center, University of Maryland, P. O. Box 169, Queenstown, MD 21658, USA.
Email: lwhall@umd.edu

Introduction

In the early 1990s, it was estimated that approximately 30% to 50% of the earth's land surface was impacted by non-point source pollutants.¹ Of the various activities associated with non-point source pollution, agricultural activities have been identified as one of the major contributors to impairment to surface waters of the United States.² The United States has more than 330 million acres of row crop agricultural land that produces an abundant supply of food and other products.³ Multiple stressors that can impact resident biological communities as a result of agricultural activities are associated with soil erosion, feeding operations, grazing, plowing, animal wastes, application of pesticides, irrigation water, and fertilizers.⁴ Therefore, decision-makers need to know the impacts of stressors associated with agricultural activities on resident biological communities in water bodies that are in close proximity to agriculture.

The common approaches used to assess impairments in agricultural water bodies due to multiple stressors are chemical monitoring, toxicity testing and biological assessments (bioassessment). Bioassessment, formally defined as a quantitative

survey of physical habitat and biological communities of a water body, is a well-established approach for determining the ecological condition of stream and river systems.^{5–8} Bioassessments provide a useful approach for integrating effects from physical, chemical, and biological stressors on aquatic organisms. The underpinnings of bioassessments are that the structure and function of an aquatic biological community can provide critical information about the quality of the surface water and sediment. Bioassessments are extremely valuable for determining the status of aquatic biological communities across large spatial scales and land use types (agricultural and urban). Information on the status of resident biological communities is particularly useful for determining impaired water bodies, developing Total Maximum Daily Loads (TMDLs), and measuring success of voluntary or regulatory actions. Bioassessments serve monitoring needs through three primary functions: (1) screening or initial assessment of conditions, (2) characterization of impairment and diagnosis, and (3) trend monitoring to evaluate improvements



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).




Figure 1. Santa Maria River watershed sample site locations.

from mitigation practices or further degradation. In addition, bioassessments also provide a direct means of measuring compliance with the goal of biotic integrity stipulated under the Clean Water Act because assemblages of aquatic organisms (ie, macroinvertebrates) comprise taxa that are differentially responsive to different environmental stressors.

Synthetic pyrethroids are a class of insecticides that have been used for over 35 years to control pests in over 120 economically important agricultural crops in the United States and globally.⁹ Ecological risk to sensitive aquatic biota (arthropods) from pyrethroid exposure has been predicted using highly protective approaches such as risk quotients (highest measured concentration/lowest toxicity value) and estimated environmental concentrations using models.⁹ However, more realistic approaches using currently available toxicity and monitoring data suggest that aquatic life may not be impacted.⁹ One water body in California's Central Coast that has been implicated as a high risk area for pyrethroids based on toxicity testing with a sensitive laboratory taxon (*Hyaella*) is the Santa Maria River watershed.¹⁰ The California Central Coast Water Board adopted the Santa Maria TMDL for toxicity and pesticides on January 30, 2014. This TMDL includes various pyrethroids. Eight of the top 12 pesticides used for agricultural crops in the Santa Maria River watershed in 2012 were pyrethroids. Since pyrethroids are highly hydrophobic, concentrations of these insecticides in sediments are a potential ecological risk issue.

Since 2006, bioassessment multiple stressor case studies have been conducted in four primarily urban wadeable California streams using benthic macroinvertebrates with concurrent measurements of habitat metrics, metals and pyrethroids to determine which stressors are most important in influencing the condition of benthic communities.¹¹ The summary results from these four case studies showed that physical habitat metrics were the most important factors influencing benthic community condition while metal concentrations were the second most important factor influencing benthic communities in these streams. Pyrethroids were the least important

constituents influencing benthic community conditions in these urban streams and for two of these streams pyrethroids were not a significant stressor to benthic communities when considered in a multiple stressor analysis. The bioassessment multiple stressor approach is ecologically relevant and provides a direct measure of stream health in concert with diagnostic analysis for identifying stressors, within a multiple stressor context, responsible for impairment. However, this approach has not been used for California water bodies located in agricultural areas where other investigators have documented pesticide occurrence and toxicity based on single species laboratory toxicity tests.¹² Therefore, this bioassessment multiple stressor approach was used in an intensive agricultural area of the Santa Maria River watershed in the Central Coast area of California to determine which stressors were most important in shaping benthic communities in this waterbody.

This 3-year study was designed to address the following specific objectives: (1) characterize benthic communities and physical habitat in Santa Maria River watershed during the spring of 2015, 2016, and 2017; (2) measure basic water quality parameters and nutrients in the water column and eight specific pyrethroids, total organic carbon (TOC), grain size, eight bulk metals and simultaneously extracted metals (SEM), and acid volatile sulfides (AVS) in sediment at various sites in concert with the bioassessments; and (3) use univariate and stepwise multiple regressions along with canonical correlation analysis to determine the relationship between various benthic metrics (ie, species richness, abundance) and physical habitat metrics (ie, embeddedness and channel flow status), pyrethroids, metals, nutrients and sediment characteristics from data sets based on sampling annually from 2015 to 2017.

Materials and Methods

Site selection

The Santa Maria River watershed is approximately 1.2 million acres composed of the large hydrologic areas containing the Cuyama Valley and the Sisque and Guadalupe watersheds.¹³ The Guadalupe hydrologic area is located in the lower watershed and is transected by the Santa Maria River which flows east to west from the confluence of the Cuyama and Sisquoc Rivers to the Pacific Ocean. The lower Santa Maria River watershed contains year-round intensively cultivated lands that supports a \$3.5 billion/year agricultural industry producing much of the nation's lettuce, artichokes, crucifer crops and strawberries.¹⁴

Our 12 study sites were located in the lower Santa Maria watershed (Figure 1). These 12 sites were sampled during the first week of March in 2015, 2016, and 2017. Water levels were generally higher in 2016 and 2017 than 2015 (a severe drought year). The lower watershed was adequately represented spatially as follows. Two sites were sampled in the mainstem Santa Maria River. Four sites were sampled in Solomon/Orcutt Creek. One site was sampled in both a channelized ditch at

Table 1. Sample site names, coordinates and three-year mean water parameters measured in Santa Maria River from 2015 to 2017.

SITE	LATITUDE	LONGITUDE	WATER TEMPERATURE (°C)	CONDUCTIVITY (μS)	PH	DISSOLVED OXYGEN (MG/L)	SALINITY (PPT)	TURBIDITY (NTU)
SM 1	34.96109	-120.64735	17.60	1987.67	8.10	10.03	1.20	121.65
SM 2	34.96013	-120.64289	18.23	1720.67	7.77	9.08	1.03	162.27
SM 3	34.95698	-120.60766	18.43	2342.00	8.08	7.91	1.40	178.30
SM 4	34.94854	-120.61559	14.80	1206.33	8.05	9.17	0.77	145.43
SM 5	34.94114	-120.57584	19.50	1837.00	8.01	10.40	1.03	94.90
SM 6	34.93232	-120.55531	17.43	2092.00	8.36	10.75	1.23	100.60
SM 7	34.9604	-120.48508	17.17	1138.67	8.64	8.87	0.70	36.33
SM 8	34.95383	-120.47218	17.23	707.00	8.65	6.41	0.43	40.60
SM 9	35.02736	-120.60011	16.37	1308.00	7.71	8.87	0.80	18.33
SM 10	35.02274	-120.58887	16.60	1660.33	7.86	9.17	1.00	15.73
SM 11	35.01407	-120.58594	16.97	1813.00	7.77	10.10	1.07	51.40
SM 12	34.9949	-120.57779	19.13	1706.67	8.21	8.56	1.00	166.53

Manzanita Berry Farm and at Main Street. Four sites were sampled in Oso Flaco Creek.

Water and sediment measurements

The following water parameters were measured at each stream site using procedures described in Kazyak¹⁵: temperature, pH, salinity, specific conductivity, dissolved oxygen, and turbidity (Table 1). All instruments were calibrated before and after sampling.

Grain size¹⁶ and TOC¹⁷ were measured on sediment samples collected from each site. Depositional areas—fine grain areas most likely to contain hydrophobic pesticides such as pyrethroids—were specifically sampled at each site when possible and three to five sediment samples from depositional areas were composited for the final sample. A stainless steel spoon (similar to a scoop) was used to collect the top 2–3 cm of sediment from each site. Approximately 1 L of sediment was collected from each site for grain size and TOC determinations as well as pyrethroids and metals. All sampling equipment was cleaned between sites using nitric acid, ethanol and distilled water. Sediment samples were stored in a cooler on ice in the field and later transferred to a refrigerator before shipment to Alpha Analytical Inc. for grain size and TOC analysis.

Physical habitat assessments

Physical habitat was evaluated at each site concurrently with benthic collections, water quality evaluations, nutrient analysis, sediment parameters, pyrethroids, and metals. The physical habitat evaluation methods followed protocols described in Harrington and Born.¹⁸ The physical habitat metrics used for

this study were based on nationally standardized protocols described in Barbour et al.¹⁹ A total of 10 continuous metrics scored on a 0 to 20 scale were evaluated (Table 2). Other non-continuous metrics including percent canopy, percent gradient, and substrate composition were also measured as reported in Hall et al.^{20–22}

Bulk metals and SEM/AVS analysis

The following bulk metals with existing threshold effects levels (TELS), conservative protective benchmarks, as described by Buchman²³ were measured on composited sediment samples for each site as previously described using EPA method 6020 m: arsenic (As), cadmium (Cd), chromium (Cr), copper (Cu), lead (Pb), nickel (Ni), and zinc (Zn). Mercury (Hg) was also measured on all sediment samples using EPA method 245.7m. The reporting limit (RL) for these metals except Zn ranged from 0.02 to 4.5 μg/g dry weight. The RL for zinc ranged from 14.4 to 21.0 μg/g dry weight.

SEM analysis was conducted for Ni, Cu, Zn, Cd, Pb, and Hg using EPA method 200.8 m. The reporting limits (μmol/dry g) for these SEMs were as follows: Ni (0.02–0.04), Cu (0.001–0.002), Zn (0.02–0.04), Cd (0.001–0.002), Pb (0.01–0.02), and Hg (0.0001). AVS were evaluated on sediment samples from each site using procedures described by Plumb.²⁴ SEM/AVS ratios were then developed for each site to provide insight on the bioavailability of these metals in sediment. The principle of SEM/AVS is based on the observation that there are some components in sediment that bind certain metals such that they are no longer available and therefore not toxic to benthic organisms.^{25,26} Sulfides in sediments have the ability to bind with divalent metals such as cadmium, copper, lead, mercury, nickel

Table 2. Scoring of 3-year mean physical habitat metrics for sites sampled in Santa Maria River from 2015 to 2017.

SITE	EPI SUBS	EMBEDD	VELDEP DIVERS	SEDIM DEP	CHAN FLOW STATUS	CHAN ALT	FREQ BEND RIFF	L BANK STAB	R BANK STAB	L BANK VEGET PROT	R BANK VEGET PROT	L BANK RIPAR ZONE	R BANK RIPAR ZONE	TOTAL
SM 1	3.67	0	3.67	0.33	17.33	16.00	8.67	1.67	4.00	2.00	5.33	6.00	8.33	77
SM 2	7.00	0	3.33	1.67	16.67	16.00	6.00	4.33	4.33	6.00	6.67	8.00	8.00	88
SM 3	0.33	0	1.67	0	13.33	0.33	1.33	0.33	1.00	0.67	1.00	0.33	0.33	21
SM 4	5.00	2.67	6.33	4.33	13.67	11.67	8.33	4.33	4.00	6.00	6.00	6.67	1.67	81
SM 5	1.00	0	4.67	0	12.00	1.00	1.00	0.67	0.67	0.33	1.00	0	0	22
SM 6	0.33	0	4.67	0	9.67	0.33	0	0	0	0.33	0.67	0	0	16
SM 7	2.67	0	0.33	3.00	15.33	0.33	0	5.33	7.33	5.00	4.67	0	0	44
SM 8	1.00	0	0.33	1.67	13.33	0.67	0.33	0.33	2.33	0.33	2.67	0	0	23
SM 9	3.00	0	4.00	0	17.33	2.33	0.67	0	8.33	0	7.67	0	6.00	49
SM 10	0.67	0	2.00	0.33	12.33	0.33	0.33	0	3.00	0	2.33	0	0	21
SM 11	5.33	0	1.67	0.33	12.33	0.67	2.67	0.33	2.00	1.00	3.00	0	0	29
SM 12	0.33	0	1.33	0.67	12.67	1.67	1.33	0	4.67	0	5.00	0	1.33	29

and zinc and may render these metals unavailable to the extent sulfides are available. Sediments from the study sites were therefore analyzed for the amount of SEM and for the amount of freely available divalent metals as SEM. Assuming that sulfides would bind with metals on a 1:1 molar basis, dividing SEM by the amount of AVS would suggest that these metals are available when the SEM/AVS ratio is greater than 1.

Nutrients analysis

The following nutrient analysis was conducted on water column samples at each site: ammonium, nitrite, nitrate, dissolved nitrogen, total nitrogen, soluble reactive phosphate, and total phosphorus.^{27–29}

Pyrethroid analysis

The pyrethroids bifenthrin, cypermethrin, cyfluthrin, deltamethrin, esfenvalerate, fenpropathrin, lambda-cyhalothrin and permethrin residues were extracted from sediment by shaking with methanol/water mixture and hexane for 1 hour. The sample was centrifuged and an aliquot of the upper hexane layer evaporated to dryness and re-dissolved in a small volume of hexane. The hexane sample was then subjected to a silica solid phase extraction (SPE) procedure prior to residue determination by gas chromatography with mass selective detection using negative ion chemical ionization (GC-MS/NICI). The limit of quantitation (LOQ) of the method was 0.14 to 0.29 ng/g dry weight for all pyrethroids.³⁰

Benthic macroinvertebrate sampling and identifications

Benthic macroinvertebrates were collected during the first week of March in 2015, 2016, and 2017 from three replicate samples at all 12 sample sites. The sampling procedures were conducted in accordance with methods described in Harrington and Born.¹⁸ Within each of these sample reaches, a riffle was located (if possible) for the collection of benthic macroinvertebrates. A tape measure was placed along the riffle and potential sampling transects were located at each meter interval of the tape. Using a random numbers table, three transects were randomly selected for sampling from among those available within the riffle. Benthic samples were taken using a standard D-net with 0.5 mm mesh starting with the most downstream portion of the riffle. A 30.5 cm x 61 cm section of the riffle immediately upstream of the net was disturbed to a depth of 10.2 to 15.2 cm to dislodge benthic macroinvertebrates for collection. Large rocks and woody debris were scrubbed and leaves were examined to dislodge organisms clinging to these substrates. Within each of the randomly chosen transects, three replicate samples were collected to reflect the structure and complexity of the habitat within the transect. If habitat complexity was lacking, samples were taken near the side margins and thalweg (deepest path) of

the transect and the procedures described above were followed. All samples were preserved in 95% ethanol.

Due to the physical nature of this water body, it was often difficult to locate a substantial number of riffles to sample. Therefore, all sites were sampled using the non-riffle method.¹⁸ This involved sampling the best available 30.5 cm x 61 cm sections of habitat throughout the reach using the same procedures described above. Best available habitat included general structure such as root wads, tires or other debris. Nine 30.5 cm x 61 cm sections were randomly selected for sampling (ie, stratified random sampling). Groups of three 30.5 cm x 61 cm sections were composited for each replicate for a total of three replicates per site.

All benthic samples were identified to the species level if possible. For taxa such as oligochaetes and chironomids, family and genus level, respectively, were often the lowest level of identification possible. Benthic macroinvertebrate subsampling (resulting in a maximum of 300 individuals) and identifications were conducted by California's Department of Fish and Wildlife (CDFW) in Rancho Cordova, California. The benthic macroinvertebrate samples were subsampled and sorted by personnel at the CDFW Laboratory located at Chico State University. Level 3 identifications (species level identifications) followed protocols outlined in Harrington and Born.¹⁸ CDFW taxonomists conducted the taxonomic identifications. Slide preparations and mounting for species such as midges and oligochaetes followed protocols from the United States Geological Survey National Quality Control Laboratory described in Moulton et al.³¹

Statistical analysis

A total of 899 statistical tests were conducted for the combined 2015, 2016, and 2017 data sets.²² In preparation for statistical analyses, the data for the 10 key benthic metrics were averaged across the three transects sampled for each site in the Santa Maria watershed. These data were merged with data sets of sediment concentrations of pyrethroids, sediment concentrations of metals, simultaneously extractable metals to AVS ratios, sediment characteristics (% TOC, % sand, and % fine sediments), nutrients in the water, and habitat metrics for each site. The sediment concentration data for pyrethroids were converted to toxicity units (TUs) by standardizing them to 1% TOC and dividing by *Hyaella* LC50 values that were also standardized to 1% TOC.³² Metal concentrations in sediment were also standardized to their relative toxicities by dividing the dry weight concentrations of each metal by their respective TEL values.²³

The statistical approach used was similar to that used for previous bioassessment/multiple stressor studies in California's urban streams.^{11,20,21,33} The potential associations between the benthic metrics and pyrethroids, metals, sediment characteristics, nutrients, and habitat metrics were explored by a series of regression techniques. Prior to this analysis, all data were unit

deviate standardized to place all dependent and independent variables on the same relative scales, as well as to produce more normalized distributions. Univariate general linear model regressions³⁴ were conducted to determine whether there were indications of significant relationships ($\alpha = 0.01$) between benthic metrics and concentrations of pyrethroids (expressed as TUs), metals in sediments (expressed as metals to TELs), SEM to AVS ratios, nutrients, sediment characteristics, and habitat metrics. Then, a series of stepwise multiple regressions³⁴ were conducted to determine potential relationships between the benthic metrics and combinations of environmental variables. Stepwise regressions were conducted for the benthic metrics versus the toxicants (pyrethroids and metals); the nutrients, sediment characteristics, and habitat metrics; and all variables combined into the same model. A second series of stepwise regressions were conducted for the benthic metrics versus principal components (PCs) of the environmental data that were produced by principal components analyses (PCA) with an orthogonal rotation (*Proc Factor*, principal components method with a “varimax” rotation³⁴). The confirmation models that have been utilized in other studies¹¹ could not be utilized in this study because there were fewer samples than the total number of environmental variables. However, separate PCAs were conducted on the toxicant data set (pyrethroids and the metals), the habitat variables (habitat metrics and sediment characteristics), and the nutrients.

The major PCs for each of these analyses (ie, the PCs that explained at least 5% of the variance in the data and having eigenvectors greater than 1.0) and the variables that had the greatest loadings on them were identified. The stepwise regression analyses were conducted on the benthic metrics versus the combined sets of PCs from all of these PCAs. Since multicollinearity could exist between the three sets of PCs, the two confirmation models described previously could not be employed.

A supplementary series of multivariate analyses were conducted to provide additional insight into the relationships between sets of benthic metrics and sets of environmental variables. Stepwise regression analyses were conducted on PCs of benthic metrics versus the entire set of PCs of the toxicants, PCs of nutrients, and PCs of the habitat variables.

The final series of multivariate analyses involved canonical correlation analyses of the PCs of the benthic data versus the PCs of the environmental data.³⁴ The canonical correlation analyses were conducted on the PCs, because the number of samples collected was relatively small compared to the total number of benthic and environmental variables to be analyzed. Thus, there were insufficient degrees of freedom to permit canonical correlation analysis of the full suite of benthic and environmental variables.

Results and Discussion

Water measurements

The 3-year mean ranges of water measurements from the 12 Santa Maria sites in Table 1 were as follows: temperature (14.8°C-19.1°C), conductivity (707-2342 uS); pH (7.7-8.6),

dissolved oxygen (6.4-10.8 mg/L), salinity (0.8-1.4 ppt), and turbidity (15.7-178 NTU). With the exception of salinity, the various water parameters were variable among the various sites. The turbidity values (>15.7 NTU) may be potentially stressful as all values exceeded the U.S. EPA turbidity criteria of 2.34 NTU for Nutrient Ecoregion III.³⁵

Physical habitat metrics

The 3-year mean physical habitat metrics (maximum score of 20 per metric) and total score (maximum of 200) for the Santa Maria sites are presented in Table 2. The various metrics as well as the total score were highly variable among the 12 sites. For example, the mean total scores ranged from 16 to 88. The highest total habitat scores were reported at the downstream sites (SM1, SM2, and SM4; see Figure 1). The narrative scoring for total physical scores is as follows: optimal (150-200), suboptimal (100-150), marginal (50-100), and poor (0-49).¹⁸ The mean total physical habitat score across all sites was 42 (ranked as poor). Nine of the 12 sites also had mean total habitat scores less than 49 reflecting poor habitat. The habitat scores reported in the Santa Maria River watershed were the lowest habitat scores we have reported from 11 different water bodies sampled in California since 2000.²²

Observed habitat conditions in wadeable waterbodies such as the Santa Maria River watershed are usually the result of the complex interplay between landscape alterations (ie, agricultural activities) and hydrogeomorphological factors. The physical habitat assessments conducted during this study were used to determine the suitability of the physical environment for aquatic biota such as benthic macroinvertebrates. Impaired physical habitat (including sediment loading) has been identified as a major stressor to aquatic life in California streams.^{36,37} Altered physical habitat structure is also considered one of the major stressors of aquatic systems throughout the United States resulting in extinctions, local extirpations and population reductions of aquatic fauna.³⁸ Identifying degraded physical habitat in streams such as the Santa Maria watershed is particularly critical for bioassessments as failure to do so can sometimes hinder investigations on the effects of toxic chemicals, as measured in this study, or other water quality related stressors as discussed later in this paper. Rankin³⁸ has also reported that there is a small but still significant risk of reporting a water quality related impact when one does not exist (false positive) when habitat assessments are insufficient or absent. Therefore, evaluation of physical habitat in agricultural streams in California's Central Coast is particularly important for multiple stressor analysis due to the intensive agriculture in the Santa Maria watershed.

TOC and grain size

The 3-year mean percent TOC ranged from 1.23% to 2.21% for the 12 Santa Maria sites as presented in Table 3. Percent sand was highly variable among sites ranging from 7.3 to 34.4. All sites were predominately fine grain areas, with silt ranging

Table 3. Three-year mean % TOC and grain size values for Santa Maria River from 2015 to 2017.

SITE	TOC	SAND	GRAVEL	SILT	CLAY
SM 1	1.98	33.50	0	42.30	24.20
SM 2	2.21	16.20	0	50.57	33.23
SM 3	2.02	12.63	0	57.77	29.60
SM 4	1.87	33.47	0	47.70	18.83
SM 5	1.99	24.10	0	48.87	27.03
SM 6	2.13	17.23	0.07	42.93	39.77
SM 7	1.46	24.30	0	49.60	26.10
SM 8	1.23	34.37	0	50.40	15.23
SM 9	1.79	9.27	0	43.07	47.67
SM 10	1.72	21.17	0.03	48.00	30.80
SM 11	1.86	7.27	0	55.90	36.83
SM 12	1.90	9.73	0.13	34.50	55.63

from 34.5% to 57.8% and clay ranging from 15.2% to 55.6%. Depositional areas—where hydrophobic chemicals such as pyrethroids could accumulate—were specifically targeted for sampling to increase the probability of measuring these insecticides if they were present in the water body.

Sediment mapping studies designed to determine the percent depositional area in this waterbody have not been conducted, but sediment mapping studies have been conducted in another Central Coast stream complex (Salinas streams) that are somewhat similar to the Santa Maria watershed. The results from the sediment mapping studies in the Salinas streams showed that depositional areas were not dominant as only 24% of the bed sediment was determined to be depositional area.³⁹ Based on field crew observations, it is highly likely that depositional area (silt and clay areas) would not be dominant in the Santa Maria river watershed.

Bulk metals and SEM/AVS

The 3-year mean concentrations for 8 metals measured in the Santa Maria watershed sites are presented in Table 4. TEL exceedances were reported at 8 sites for arsenic and at all 12 sites for cadmium and nickel. Natural sources for these metals in local soil cannot be ruled out as a contributor to the potentially toxic concentrations reported. For example, the cadmium TEL exceedances at all sites are likely related to the presence of shale-based soil in the Santa Maria watershed that have historically high concentrations of cadmium.⁴⁰ Serpentine soils in the Santa Maria watershed area, a known source of nickel as reported by other investigators, may be a natural source for the high nickel concentrations reported in this watershed.^{41,42} Possible anthropogenic sources of cadmium in the Santa Maria

watershed are phosphorous-based fertilizers or micronutrient applications.⁴³ Anthropogenic sources of arsenic in this watershed may also be related to the use of phosphorous-based fertilizers and micronutrients and possibly diffuse sources such as atmospheric fallout.⁴³

The SEM/AVS data in Table 5 showed that 10 sites had ratios greater than 1 with at least one metal exceeding at TEL at all sites thus suggesting predicted potential metals toxicity to resident benthic communities at these sites. The upstream site (SM-12) in Osco Flaco Creek had the highest SEM/AVS ratio (15.9), while the two downstream sites in the mainstem of the Santa Maria River had the lowest SEM/AVS ratios (both ratios less than 1).

Nutrients analysis

Ranges of the 3-year nutrient concentrations in the Santa Maria River watershed were as follows: dissolved ammonium (0.07–10.4 mg/L), dissolved nitrate (4.2–54.6 mg/L), dissolved nitrite (0.08–1.23 mg/L), dissolved nitrogen (4.6–70.2 mg/L), total nitrogen (6.2–85.7 mg/L), soluble reactive phosphate (0.05–1.2 mg/L), and total phosphorus (0.09–2.5 mg/L) (Table 6). Highest total nitrogen concentrations were reported at upstream site SM11 in Osco Flaco Creek (see Figure 1), while the highest total phosphorus was reported at upstream site SM7. Currently, the State of California is in the process of developing nutrient water quality criteria for the state so at this time there are no California nutrient criteria that can be compared with the nutrient measurements reported in the Santa Maria watershed.

However, the relationship of nutrients to macroinvertebrate communities has been addressed in various studies as described below. Ashton et al⁴⁴ reported that taxonomic and functional metrics of an index of biotic integrity (IBI) were significantly associated with ammonia-N and nitrite-N in Maryland streams. Wang et al⁴⁵ reported that 18 of 26 macroinvertebrate measures were significantly correlated with at least one nutrient measure in Wisconsin streams. These investigators specifically reported that EPT individuals and taxa and mean tolerance value metrics had the strongest correlations with most nutrient measures. Maul et al⁴⁶ reported that nutrients such as total phosphorus and ammonia were important variables for structuring benthic communities in northwest Mississippi streams. Other investigators also reported that nutrients tended to increase plant-associated macroinvertebrates at low levels with a flattening response at intermediate levels and a decline at high levels.⁴⁷ Therefore, based on the above studies, there is certainly background information to suggest that nutrients could have significant relationships to benthic communities in agricultural waterbodies such as the Santa Maria watershed.

Pyrethroids

Ranges of 3-year mean concentration of 8 pyrethroids normalized to 1% TOC presented in Table 7 were as follows: bifenthrin (0.23–31.9 ng/g), cyfluthrin (0.06–3.5 ng/g), cypermethrin

Table 4. Three-year mean bulk metals concentrations in sediment ($\mu\text{g/g dw}$) for Santa Maria River sites sampled from 2015 to 2017.

SITE	AS	CD	CR	CU	PB	HG	NI	ZN
SM 1	6.00	1.03	21.77	19.97	8.57	0.04	26.53	64.40
SM 2	6.62	0.97	25.47	21.57	9.12	0.05	25.50	74.33
SM 3	6.64	0.94	21.77	21.97	10.40	0.03	22.20	74.90
SM 4	4.24	0.97	24.50	18.50	6.69	0.04	38.90	53.57
SM 5	5.29	1.60	23.27	19.23	7.90	0.05	34.13	69.40
SM 6	4.85	1.77	31.07	18.57	6.49	0.06	44.73	87.27
SM 7	6.73	0.87	22.57	24.67	11.63	0.07	22.47	116.33
SM 8	5.17	0.83	16.80	17.60	10.13	0.03	18.47	84.90
SM 9	7.21	1.24	25.17	25.33	9.71	0.06	24.03	84.53
SM 10	7.35	1.07	20.43	20.57	9.87	0.05	22.43	68.57
SM 11	6.25	1.11	22.47	32.57	8.83	0.06	21.20	83.83
SM 12	8.68	0.90	32.13	33.27	13.27	0.10	34.37	97.17
TEL	5.9	0.596	37.3	35.7	35	0.174	18	123

Metals concentrations exceeding threshold effects levels (TELs) are in bold.

Table 5. Three-year mean concentrations of acid volatile sulfide (AVS), simultaneously extracted metals (SEM), and the SEM/AVS ratios in sediment for sites sampled in the Santa Maria River from 2015 to 2017. Bold SEM/AV ratios >1 suggest metals are bioavailable and may be toxic. AVS values below the detection limit were assigned a value of $\frac{1}{2}$ the detection limit in the SEM/AVS calculation.

STATION	(CONCENTRATIONS IN $\mu\text{MOLE/G DRY WEIGHT}$)						TOTAL SEM	SEM/AVS
	AVS	NI	CU	ZN	CD	PB		
SM 1	9.22	0.15	0.14	0.33	0.01	0.01	0.64	0.62
SM 2	1.07	0.16	0.16	0.38	0.00	0.03	0.74	0.79
SM 3	1.28	0.13	0.12	0.29	0.01	0.02	0.57	4.86
SM 4	0.26	0.29	0.14	0.40	0.01	0.02	0.86	8.69
SM 5	5.43	0.25	0.14	0.49	0.01	0.02	0.91	3.55
SM 6	0.89	0.34	0.16	0.79	0.01	0.02	1.32	5.79
SM 7	21.82	0.15	0.21	0.76	0.01	0.04	1.16	4.34
SM 8	25.81	0.14	0.14	0.80	0.01	0.04	1.12	3.75
SM 9	3.00	0.14	0.19	0.71	0.01	0.03	1.07	2.25
SM 10	0.29	0.16	0.17	0.47	0.01	0.03	0.84	6.56
SM 11	1.76	0.14	0.32	0.57	0.01	0.02	1.06	5.04
SM 12	0.07	0.23	0.29	0.55	0.01	0.04	1.12	15.99

(0.13–1.3 ng/g), deltamethrin (0.05–0.12 ng/g), esfenvalerate (0.04–4.5 ng/g), fenprothrin (0.04–7.8 ng/g), lambda cyhalothrin (0.02–24.3 ng/g), and permethrin (0.27–22.9 ng/g). Highest concentrations of pyrethroids (ng/g at 1% TOC) based on maximum concentrations in descending order were reported for bifenthrin, lambda-cyhalothrin, permethrin, fenprothrin, esfenvalerate, cyfluthrin, cypermethrin, and

deltamethrin. Highest concentrations of three pyrethroids (bifenthrin, fenprothrin, and lambda-cyhalothrin) occurred at upstream site SM11 in Oso Flaco creek.

Toxic units (TU) calculations were determined for each pyrethroid by dividing the 1% TOC normalized concentration by the sediment *Hyalella* LC50 concentration (a species highly sensitive to pyrethroids) that was also 1% TOC normalized

Table 6. Three-year mean nutrient concentrations from Santa Maria river sites sampled from 2015 to 2017.

SITE	DISSOLVED AMMONIUM (MG N/L)	DISSOLVED NITRATE (MG N/L)	DISSOLVED NITRITE (MG N/L)	DISSOLVED NITROGEN (MG N/L)	TOTAL NITROGEN (MG N/L)	SOLUBLE REACTIVE PHOSPHATE (MG P/L)	TOTAL PHOSPHORUS (MG P/L)
SM 1	0.75	16.93	0.32	17.98	20.13	0.29	0.64
SM 2	0.62	19.30	0.35	20.23	22.73	0.35	0.64
SM 3	0.37	29.93	1.04	31.30	32.90	0.23	0.64
SM 4	0.23	17.14	0.19	17.54	19.07	0.41	0.79
SM 5	0.33	33.69	0.35	34.40	37.07	0.41	0.69
SM 6	0.08	30.91	0.82	31.81	34.47	0.42	0.90
SM 7	0.38	10.22	0.27	10.84	12.83	1.19	2.53
SM 8	0.09	4.17	0.29	4.55	6.18	0.20	0.40
SM 9	0.27	32.27	0.26	32.77	33.30	0.09	0.19
SM 10	0.07	31.43	0.08	31.60	32.77	0.05	0.09
SM 11	10.37	51.80	0.48	70.15	85.70	0.22	0.46
SM 12	5.41	54.57	1.23	61.23	72.90	0.64	1.61

(Table 8).³² TU concentrations exceeding 1.0 were predicted to be potentially toxic. The sum of pyrethroid TUs using *Hyalella* indicated that 9 of the 12 sites were predicted to be toxic due to pyrethroids. The highest sum of pyrethroid TU (11.7) was reported at site SM11 in Oso Flaco creek (see Figure 1).

The 3-year mean sum of pyrethroid TUs for all 12 Santa Maria site means was 2.7. This value is approximately 50% lower than the 3-year mean of another group of California Central Coast streams (three streams in the Salinas River watershed) where the mean of 13 Salinas stream mean site means of 6.1 was calculated.⁴⁸ Therefore, it appears that pyrethroid sediment concentrations are somewhat lower in the Santa Maria watershed when compared with data from other California Central Coast streams.

Benthic communities

The total number of different benthic taxa collected in the Santa Maria watershed in 2015, 2016 and 2017 was 75, 90, and 46, respectively. While the total number of individual benthic taxa collected by year was 8872 for 2015, 7880 for 2016, and 3607 for 2017. For 2017, both the number of different benthic taxa and the number of individuals was lower than for the previous 2 years. The reason for this difference may be related to the much greater flow in 2017 as reported in Hall et al²² compared to the previous 2 years that also resulted in reduced instream vegetation (habitat for benthic invertebrates).

The five most dominant benthic taxa collected during the 3-year study comprising 51% of the total taxa and the reported % of the total were as follows: *Tubificidae* unidentified immature (oligochaetes)—13.4%; *Cricotopus* (chironomid)—10.7%; *Physa*

(snails)—6.8%; *Cricotopus bicinctus* group (chironomid)—6.5%; and *Enchytraeidae* (oligochaetes)—6.4% (Table 9). These five taxa are generally considered tolerant to moderately tolerant of environmental stressors.¹⁸ The 3-year mean values for the various benthic metrics were highly variable among the 12 Santa Maria sites. For example, taxa richness ranged from 12 to 23, percent tolerant taxa ranged from 41 to 71, percent collectors-gatherers ranged from 31 to 72, percent predators ranged from 3.3 to 18, and total abundance ranged from 232 to 6506 (Table 10).

The state of California has recently developed a state-wide index that translates complex data about benthic macroinvertebrates into a measure of stream health.⁴⁹ This index is called the California Stream Condition Index (CSCI). The CSCI combines a multimetric index that measures ecological structure and function and an observed to expected (O/E) index that measures taxonomic completeness. Categories of biological condition based on 0-1 scoring are as follows: ≤ 0.62 = very likely altered; 0.63-0.79 = likely altered; 0.80-0.91 = possibility altered; and ≥ 0.92 likely intact. The 3-year mean value for the Santa Maria sites sampled was only 0.33 with values ranging from 0.19 to 0.53—very likely altered category—so the benthic communities in this watershed would be considered impaired.

Relationship of benthic metrics to all stressors

Interpretation of the relationships of benthic metrics to environmental stressors requires an understanding of how different benthic metrics are expected to respond to impairment.^{18,50} In the statistical analyses presented below, the definition of the benthic metrics and expected response of the various benthic

Table 7. Three-year mean pyrethroid pesticide concentrations @ 1% TOC in sediment (ng/g dw) for sites sampled in the Santa Maria River from 2015 to 2017.

SITE	% TOC	BIFENTHRIN	CYFLUTHRIN	CYPERMETHRIN	DELTAMETHRIN	ESFENVALERATE	FENPROPATHRIN	LAMBDA-CYHALTHRIN	PERMETHRIN
SM 1	1.98	1.82	0.36	0.61	0.07	1.96	0.56	1.80	1.67
SM 2	2.21	0.83	0.21	0.41	0.05	1.33	0.30	0.67	0.63
SM 3	2.02	0.77	0.56	0.61	0.05	2.17	0.04	1.17	1.07
SM 4	1.87	3.26	0.34	0.33	0.05	2.33	0.97	5.49	1.27
SM 5	1.99	9.94	3.47	1.25	0.05	1.78	5.91	1.18	4.61
SM 6	2.13	0.23	0.61	0.80	0.06	4.50	0.04	0.98	8.35
SM 7	1.46	3.18	0.97	1.06	0.08	1.99	1.27	3.83	22.94
SM 8	1.23	2.26	0.42	0.73	0.12	1.58	0.47	0.99	7.36
SM 9	1.94	11.42	0.16	0.22	0.06	1.27	3.31	3.78	2.11
SM 10	1.57	14.60	0.07	0.13	0.07	0.41	5.97	0.20	0.43
SM 11	1.86	31.92	0.24	0.29	0.07	0.44	7.81	24.25	1.28
SM 12	1.90	13.99	0.06	0.19	0.06	0.04	3.37	0.34	0.27

TOC: total organic carbon.

Table 8. Three-year mean toxic unit (TU) calculations based on *Hyalella* for pyrethroid concentrations (1% TOC normalized) for sites sampled in the Santa Maria River 2015 to 2017. The mean sum of TUs by site is also included. Values exceeding 1.0 TU are in bold.

Site	BIFENTHRIN	CYFLUTHRIN	CYPERMETHRIN	DELTAMETHRIN	ESFENVALERATE	LAMBDA-CYHALOTHRIN	PERMETHRIN	SUM
	TU	TU	TU	TU	TU	TU	TU	TU
SM 1	0.35	0.03	0.16	0.01	0.13	0.40	0.02	1.10
SM 2	0.16	0.02	0.11	0.01	0.09	0.15	0.00	0.54
SM 3	0.14	0.16	0.16	0.01	0.14	0.26	0.01	0.78
SM 4	0.63	0.03	0.09	0.01	0.15	1.22	0.01	2.14
SM 5	1.90	0.32	0.33	0.01	0.12	0.27	0.04	2.99
SM 6	0.04	0.06	0.21	0.01	0.29	0.22	0.08	0.91
SM 7	0.61	0.09	0.28	0.01	0.13	0.88	0.21	2.21
SM 8	0.43	0.04	0.19	0.02	0.10	0.22	0.07	1.07
SM 9	2.20	0.01	0.06	0.01	0.08	0.84	0.02	3.22
SM 10	2.81	0.01	0.04	0.01	0.03	0.05	0.00	2.93
SM 11	6.14	0.02	0.08	0.01	0.03	5.40	0.01	11.70
SM 12	2.69	0.01	0.05	0.01	0.00	0.08	0.01	2.83

Table 9. Five dominant taxa collected in the Santa Maria River from 2015 to 2017. The entire benthic taxa list is available in Hall et al.^{20–22}

LOWER TAXA	HIGHER TAXA	N	TOTAL %	CUMULATIVE %
<i>Tubificidae unid.imm.</i>	<i>Tubificid</i>	2721	13.365	13.365
<i>Cricotopus</i>	<i>Chironomidae</i>	2182	10.718	26.730
<i>Physa</i>	<i>Physidae</i>	1384	6.798	37.448
<i>Cricotopus bicinctus grp</i>	<i>Chironomidae</i>	1318	6.474	44.246
<i>Enchytraeidae</i>	<i>Tubificida</i>	1295	6.361	50.720

metrics to impairment is as follows: taxonomic richness—total number of individual taxa (decrease); % dominant taxa—percent composition of the single most abundant taxa (increase); Shannon Diversity—an index that accounts for both abundance and evenness of species present (decrease); tolerance value—value between 0 and 10 weighted for abundance of individuals designated as pollution tolerant (higher values) and intolerant (lower values) (increase); % tolerant taxa—% of organisms in sample that are highly tolerant to impairment as indicated by a tolerance value of 8, 9, or 10 (increase); % collectors/filterers—% of macrobenthos that collect or filter fine particulate matter (increase); % collectors/gatherers—% of macrobenthos that collect and gather fine particulate matter (increase); % grazers—% of macrobenthos that graze on periphyton (variable); % predators—% predator individuals (decrease); and abundance—total number of individuals (decrease).

The results of the stepwise analyses of benthic metrics versus all environmental variables from the combined 2015, 2016, and 2017 data set are shown in Table 11. Sediment TOC was the most frequently observed significant environmental variable that, in combination with other variables, displayed relationships with benthic metrics: displaying direct relationships with *taxonomic richness*, % *collectors/gatherers*, % *grazers*, and % *predators*; and displaying inverse relationships with *tolerance value*, and % *collectors/filterers*. Some of the benthic metrics that displayed direct relationships with TOC also displayed inverse relationships with various metals: *taxonomic richness* was inversely related to copper; % *collectors/gatherers* was inversely related to chromium; and % *predators* was inversely related to nickel. The % *predators* metric was also inversely related to Total TUs. *Shannon's index*, a diversity metric, was inversely related to lambda cyhalothrin, total phosphorous, and directly related to fines, while % *dominant taxon*, the index that is indicative of lack of diversity, displayed the opposite relationships with these environmental variables. The benthic metric % *tolerant taxa*, generally indicative of stressed communities when high, was inversely related to the habitat metric *bend/riffle frequency*, dissolved nitrogen, and directly related to ammonium. Most of the relationships between benthic metrics and the various environmental variables were individually quite weak (most R^2 values <0.20), but some combinations of the environmental variables explained more of the variations of some of the benthic metrics. Thus, with the exception of TOC, no

single environmental variable or group of variables appeared to emerge from these analyses as being the dominant one in influencing benthic communities.

The results of the stepwise regression analyses of benthic metrics versus PCs of toxicants, nutrients, and habitat variables are presented in Table 11. While there were fewer significant relationships between benthic metrics and PCs than with combinations of individual environmental variables, several benthic metrics were significantly related to HabPC3, the PC positively loaded by TOC and fine sediments: *tolerance value* and % *collectors/filterers* were inversely related to HabPC3, and % *collectors/gatherers* was directly related to this PC. The metrics % *collectors/gatherers* and % *collectors/filterers* were also inversely related to ToxPC4, the PC positively loaded by lambda cyhalothrin, copper, and, to a lesser extent, by bifenthrin. The diversity metric *Shannon's index* was inversely related to NutPC2, the PC that was positively loaded by total phosphorous and dissolved phosphorous.

Most of the relationships shown to be significant in the stepwise regressions involving combinations of individual environmental variables were not supported by the results from the stepwise regressions of the PCs of the groups of environmental variables. Even those significant relationships that were observed between a few of the benthic metrics and environmental PCs did not consistently make sense ecologically for typical benthic communities. *Tolerance value*, % *collectors/filterers*, and % *collectors/gatherers*, three benthic metrics generally believed to be indicative of stressed communities did not display consistent relationships to HabPC3 (the PC indicative of fine, organic rich sediments), since the former two metrics displayed inverse relationships with it, while the latter metric displayed a direct relationship to it. Moreover, % *collectors/filterers*, and % *collectors/gatherers* both displayed inverse relationships to the ToxPC4. If the toxicants positively loaded by this PC (lambda cyhalothrin, copper and bifenthrin) had represented a stress to the benthic communities, one would have expected these benthic metrics to display a direct relationship.

In considering these results, a caveat should be noted. The limited number of samples for the combined 3-year data set prevented the typical confirmation analyses that has been used and previously as described in the statistical analysis section of this paper. Thus, multicollinearity may exist not only between the individual environmental variables, but also between the

Table 10. Three-year mean benthic metrics by site for the 12 Santa Maria River sites sampled from 2015 to 2017.

SITE	SM 1	SM 2	SM 3	SM 4	SM 5	SM 6	SM 7	SM 8	SM 9	SM 10	SM 11	SM 12
Taxonomic richness	19	22	12	14	20	16	15	20	23	22	20	13
% Dominant taxon	35.00	26.07	30.53	62.20	33.20	29.27	54.17	65.13	31.27	47.00	30.27	35.93
Number Ephemeroptera taxa	1	0	0	0	0	0	0	0	0	0	0	0
Number Plecoptera taxa	0	0	0	0	0	0	0	0	0	0	0	0
Number Trichoptera taxa	1	0	0	0	0	0	0	0	0	0	0	0
EPT taxa	1	0	0	0	0	0	0	0	0	0	0	0
EPT Index (%)	3.33	0	0	0	0	0	0	0	0	0	0	0
Sensitive EPT Index (%)	1.00	0	0	0	0	0	0	0	0	0	0	0
Shannon diversity	2.09	2.20	1.83	1.06	1.89	1.98	1.41	1.15	2.05	1.84	1.97	1.72
Tolerance value	7.96	8.55	7.82	7.92	8.86	7.50	8.11	7.61	8.97	7.68	7.97	6.89
% Intolerant taxa (0-2)	2.33	4.67	0.00	0.00	5.00	0.00	4.00	0.00	2.67	0.00	2.67	0.00
% Tolerant taxa (8-10)	41.67	57.67	59.67	53.67	53.00	55.33	70.67	63.67	54.00	56.00	59.33	40.67
% Baetidae	1.00	0	0	0	0	0	0	0	0	0	0	0
% Chironomidae	54.67	40.00	43.00	63.33	15.00	34.33	47.00	67.67	44.00	42.00	42.00	45.67
% Hydropsychidae	1.00	0	0	0	0	0	0	0	0	0	0	0
% Collectors gatherers	59.67	57.33	47.33	56.67	44.00	47.33	53.33	64.33	61.00	30.67	71.67	60.00
% Collector-filterers	26.33	32.00	29.33	32.67	35.67	28.67	32.00	28.67	29.00	52.33	20.00	9.00
% Scrapers	4.33	7.33	20.67	10.00	16.67	19.00	11.00	2.00	7.00	6.00	6.33	4.67
% Predators	13.00	16.00	11.33	8.67	7.00	7.33	17.00	18.00	13.00	13.00	5.67	3.33
% Shredders	4.67	0	0	0	0	2.00	0	0	0	0	0	0
Total abundance (#/sample)	778	524	232	6506	2489	1124	1451	1006	832	1673	3748	546

Table 11. Results of stepwise multiple linear regression models for the Santa Maria watershed in the combined 2015, 2016, and 2017 data set: (a) benthic metrics versus toxicity units for pyrethroids, habitat metrics, nutrients, sediment characteristics, and metals to TEL ratios; and (b) benthic metrics versus principal components of the environmental data; only variables that were significant at $\alpha = 0.01$ were included in the models (NS = not significant). The direction of the relationship for each significant variable is indicated (+ = direct; - = inverse), as is the contributed R^2 value.

BENTHIC METRICS	(A) SIGNIFICANT ENVIRONMENTAL VARIABLES (R^2)	(B) SIGNIFICANT ENVIRONMENTAL PRINCIPAL COMPONENTS ^a
Taxonomic richness	+TOC (0.14), -Copper (0.13)	NS
% Dominant taxon	+Lambda-cyhal. (0.16), +Total Phosphorous (0.13), --Fines (0.13)	NS
Shannon diversity	-Lambda cyhalothrin (0.17), -Total Phosphorous (0.16), +Fines (0.15)	-NutPC2 (0.18)
Tolerance value	-TOC (0.29)	-HabPC3 (0.18)
% Tolerant taxa	- Frequency of Bends (0.20), -Dissolved Nitrogen (0.19), + Ammonium (0.17)	NS
% Collectors/gatherers	-Chromium (0.22), +TOC (0.16)	+HabPC3 (0.21), -ToxPC4 (0.12)
% Collectors/filterers	-TOC (0.29)	-HabPC3 (0.30), -ToxPC4(0.14)
% Grazers	+TOC (0.21)	NS
% Predators	-Nickel (0.21), -Total TUs (0.14), +TOC (0.15)	NS
Abundance	NS	NS

TOC: total organic carbon.

^aThe toxicants with the largest loadings for the toxicant principal components (proportion of variance explained by PCs are in parentheses) are: ToxPC1 (0.30) = + Lead, +Arsenic, +Zinc, and +Mercury; ToxPC2 (0.17) = +Nickel, +Cadmium, and +Chromium; ToxPC3 (0.13) = +Permethrin and +Cypermethrin; ToxPC4 (0.11) = +Lambda-cyhal., +Copper, and +Bifenthrin. The habitat principal components with the largest loadings are: HabPC1 (0.46) = +Channel alteration, +Bend/Riffle frequency, +Riparian zone, +Total score, +Width, and +Velocity depth diversity; HabPC2 (0.12) = +Bank stability, +Channel flow status, +Vegetative protection of stream banks, +Depth, and + Epifaunal substrate; HabPC3 (0.12) = +TOC and +Fines. The nutrient principal components with the largest loadings are: NutPC1 (0.51) = +Total Nitrogen, +Dissolved Nitrogen, +Nitrates, and +Ammonium; NutPC2 (0.29) = +Total phosphorus and +Dissolved phosphorus; NutPC3 (0.13) = +Nitrites.

PCs calculated for each group of variables separately. Thus, the significant, albeit rather weak, patterns that were observed may have been confounded by this multicollinearity (i.e. other correlated environmental variables, PCs or combinations may have been responsible for the relationships). The definitive confirmation analyses could not be conducted to determine whether significant patterns between benthic metrics and habitat variables persisted when the potential effects of toxicants were taken into account and vice versa. Thus, the results of the stepwise statistical analyses should not be over-interpreted.

Multivariate analysis of the PCs of benthic metrics versus PCs of environmental variables

The results of the stepwise regressions of PCs for the benthic metrics versus the PCs for all of the environmental variables for the combined 2015, 2016, and 2017 data sets are presented in Table 12. BenPC1 (the PC that was positively loaded by the benthic metrics % collectors/gatherers, taxonomic richness, % predators, and abundance, and negatively loaded by % collectors/filterers) was directly related to HabPC3 (the PC that was positively loaded by the sediment characteristics TOC and Fines) and was inversely related to ToxPC2 (the PC that was positively loaded by the metals nickel, cadmium, and chromium). Thus, the stepwise regressions for the PCs tended to confirm some of the patterns observed for the benthic metrics associated with BenPC1 in the analyses of individual benthic metrics with combinations of environmental variables shown in Table 11.

BenPC2 (the PC that was positively loaded by the diversity metric *Shannon's index* and was negatively loaded by % *Dominant taxon*) was inversely related to NutPC2 (the nutrient PC that was positively loaded by total phosphorous and dissolved phosphorous) and to ToxPC4 (the toxicant PC that was positively loaded by lambda cyhalothrin, copper, and, to a lesser extent, by bifenthrin). These relationships are similar to those observed for the relationships between the *Shannon's index* and % *dominant taxon*, and the environmental variables lambda cyhalothrin and phosphorus (see Table 11).

For the combined 2015, 2016, and 2017 data set, a canonical correlation was conducted on the benthic metric PCs versus the PCs for the environmental variables (ie, PCs for toxicants, nutrients, and habitat metrics). The results from the canonical correlation analysis are shown in Figure 2. There was a significant and moderately strong ($R^2 = 0.65$) relationship between the Canonical Variate for benthic PCs (CVBen; plotted on the y-axis) and the Canonical Variate I for PCs for environmental data (CVEnv; plotted on the x-axis). The CVBen for benthic metrics was positively correlated to BenPC1 and BenPC2, and displayed a somewhat weaker negative correlation to BenPC3. The benthic PCs that were positively loaded by benthic metrics associated with more diverse and healthier benthic communities tended to increase going up the y-axis, while those loaded on metrics associated with less diverse, more stressed communities tended to increase going down the y-axis. The CVEnv was directly correlated to HabPC3, the PC that was positively loaded by the sediment characteristics TOC and

Table 12. Results of stepwise regression analyses of principal components for benthic metrics (BenPC^a) versus the principal components for environmental data (see footnote for Table 11 for loadings of the ToxPCs, HabPCs, and NutPCs) for the Santa Maria watershed in the combined 2015, 2016, and 2017 data set. Only variables that were significant at $\alpha = 0.01$ were included in the models (NS = not significant). The direction of the relationship for each significant variable is indicated (+ = direct; - = inverse), as is the contributed R^2 values.

BENTHIC METRIC PCS	PROB.	R^2	SIGNIFICANT ENVIRONMENTAL PCS (R^2)
BenPC1	<0.001	0.33	+HabPC3 (0.17), -ToxPC2 (0.16)
BenPC2	<0.001	0.33	-ToxPC4 (0.17), -NutPC2 (0.16)
BenPC3	NS		
BenPC4	NS		

^aThe benthic metrics with the largest loadings for the benthic principal components (proportion of variance explained by PCs are in parentheses) are: BenPC1 (0.42) = +% collectors/gatherers, -% collectors/filterers, +taxonomic richness, +% predators, and +abundance; BenPC2 (0.19) = +Shannon, -% dominant taxon; BenPC3 (0.14) = +tolerance value and +tolerant taxa; BenPC4 (0.09) = +% Grazers.

Fines, and displayed a somewhat weaker negative correlation to NutPC2, the PC that was negatively loaded by total phosphorous and dissolved phosphorous. Samples higher on the x-axis tended to be from sites characterized by more organic rich, fine sediments and lower phosphorous concentrations.

In looking at the trend line relationship in Figure 2, data points found in the upper right hand quadrant of the figure tended to be from samples that were characterized by somewhat healthier, more diverse benthic communities, finer and more organic rich sediments, and lower phosphorous concentrations at these sites. Conversely, data points found in the lower left quadrant tended to be from more stress impaired, less diverse benthic communities with less organic rich sediments and greater phosphorous concentrations.

The direct relationship we have reported between TOC and diversity measures dominated by tolerant benthic taxa (oligochaetes) in the Santa Maria watershed has been addressed in other studies.^{51,52} Grumiaux et al.⁵¹ reported that pollution resistance benthic taxa abundance increased with an increase in organic content in rivers and canals in northern France. Since our sites in the Santa Maria watershed are dominated by tolerant benthic taxa, such as oligochaetes, the direct relationship we have reported with TOC and diversity measures for benthic assemblages dominated by tolerant taxa is supported by these authors. In another study conducted in the San Francisco estuary, Thompson and Lowe⁵² reported that tolerant taxa such as oligochaetes increased with elevated concentrations of TOC. The results from the Thompson and Lowe⁵² study also support our findings showing that TOC is an important factor shaping benthic communities dominated by tolerant taxa such as oligochaetes.

Overall, several observations may be made from the results of the canonical correlation analysis comparing the overall patterns of combinations of benthic metrics to combinations of environmental variables. First of all, the ToxPCs related to toxicants did not emerge as appearing to be significant to the BenPCs in the 3-year Santa Maria River data. The nitrogen-based nutrients also did not appear to be significant in the analyses of the 3-year data set. Rather, higher concentrations of




Figure 2. Results of the canonical correlation analysis of PCs for benthic metrics versus PCs for environmental data (i.e. PCs separately calculated for habitat metrics, nutrients, and toxicants in sediments) for the Santa Maria River combined 2015, 2015, and 2017 data set: canonical variate 1 for BenPCs versus the canonical variate 1 for HabPCs, NutPCs, and ToxPCs (see footnotes in Tables 11 and 12 for loadings of variables for PCs). The PCs for benthic biological metrics (y-axis) and the PCs for the environmental (x-axis) that were most highly correlated with the canonical variates are shown ($r \geq 0.5$); moderate correlations ($r \sim 0.5$) are shown in parentheses, along with the direction of their relationships (shown by arrows). The regression trend line and 95% confidence limits are indicated by solid and dashed lines, respectively.

phosphorous-based nutrients (NutPC2) appears to be somewhat important in characterizing the environmental conditions associated with more stressed, less diverse benthic communities (BenPC2 and BenPC3). The habitat metrics that characterize physical stream quality conditions did not appear to be significant in relationship to the benthic communities in the “big picture” canonical correlation analysis. This observation is quite different from the results of previous studies in California urban streams¹¹ in which some of the habitat metrics represented key environmental factors that were shown to be associated with the relative health of benthic communities as indicated by the benthic metrics.

A few final caveats should be made regarding the multivariate analyses. While the stepwise analyses involving the PCs for various groups of environmental PCs (toxicants, nutrients, habitat metrics, and sediment characteristics) did address issues associated with multicollinearity between variables within the groups, there may still be multicollinearity between the PCs of the different groups of environmental variables. Thus, one PC could have been selected as being significant during the stepwise process, others that were correlated with it would have been discarded, even though they may be ecologically significant. The canonical correlation analysis tends to overcome this issue, since important environmental PCs identified as being correlated to the CVEnv by the model could be mutually related to the CVBen and the BenPCs to which it was correlated. Therefore, the relatively modest number of samples for even the 3-year data set prevents taking the ideal statistical approach of using PCs for the entire set of environmental variables (in which case, the PCs would not be correlated to each other) or even the use of the entire individual environmental variables in the canonical analyses.¹¹ Furthermore, the relatively small data set may affect the power of the multivariate models to detect more subtle relationships.

Results overview of the statistical analysis

There were relatively few significant relationships between benthic metrics and toxicants in the sediments in the stepwise analyses of the data from the Santa Maria watershed. Most of the statistically significant relationships that were observed between benthic metrics and pyrethroids were largely influenced by an outlier sample (Site SM11) observed in 2017 that had very high concentrations of bifenthrin and to a somewhat lower lambda-cyhalothrin TU, although still greater than 1. However, these significant relationships disappeared with these two pyrethroids in the stepwise analyses for combined 2015, 2016, and 2017 data set analyses, suggesting that the impact of the outlier data point was no longer a factor in the analyses of a somewhat larger data set.

There were also few relationships between the benthic metrics and metals. For the analyses of the combined 2015, 2016, and 2017 data set, the benthic metric % *predators* was inversely related to nickel. Nickel was quite high in the sediments with the TEL value being exceeded at all sites (Table 4). While the relationships between this benthic metric and nickel made sense ecologically, the strength of this relationship only explained a low to moderate proportion of the variance for this metric ($R^2 = 0.21$). However, none of the benthic metrics were significantly related to SEM to AVS ratios for any of the metals, including nickel (ie, the benthic metrics were not significantly related to the forms of the metals generally believed to be most biologically available), so it is difficult to say that nickel in sediments represent a major factor influencing the benthic communities of Santa Maria watershed.

There were more apparent relationships between benthic metrics and nutrients. The benthic metric % *tolerant taxa* displayed a direct relationship with ammonium in the combined 2015–2017 data set. This relationship persisted in the stepwise analyses of all environmental data, as well as those involving PC loaded by nitrogen nutrients. In contrast, the phosphorous-based nutrients did not display significant relationships in the stepwise analyses involving only the nutrients, but they were significant in relationship to diversity-related metrics in combination with other environmental variables in the stepwise analyses for all variables and in the multivariate analyses: *Shannon's index* was inversely related to total phosphorous, while % *dominant taxon* was directly related to it. These relationships persisted in the multivariate analyses involving benthic and environmental PCs.

In contrast to the results of our previous bioassessment multiple stressor studies, the benthic metrics did not have many significant relationships with traditional habitat metrics in the Santa Maria data set.¹¹ *Tolerance taxa* was inversely related to *Bend/Riffle frequency* in the stepwise analyses of the combined data set, but it did not display a significant role in the big picture multivariate analyses (ie, the regressions involving the PC of the habitat metrics or the canonical correlation analysis).

TOC in sediment displayed a significant relationship with benthic metrics in the stepwise regression analyses and the multivariate analyses. The metrics *tolerance value* and % *collectors/filterers* were inversely related to TOC in sediments, while % *grazers* was directly related to it. These benthic metrics displayed similar relationships to HabPC3, the PC that was positively loaded by TOC in the sediment and fine sediments.

The summary multivariate canonical correlation analysis indicated that less stressed, more diverse benthic communities tended to be associated more with TOC-rich finer sediments and lower concentrations of phosphorous-based nutrients, and more stressed, less diverse benthic communities tended to be associated with less organically rich, somewhat less fine sediments and higher phosphorous concentrations. Neither toxicants nor habitat metrics were shown to be important factors in this analysis. The extremely poor habitat metric scores observed consistently at all sites throughout the Santa Maria watershed may be responsible for this group of variables not being a major factor that would shape spatiotemporal patterns of the benthic communities throughout the waterbody.

Conclusion

Resident benthic communities in the Santa Maria River watershed were reported to be impaired based on a benthic index developed by the State of California and the dominant presence of tolerant taxa. A suite of possible constituents that could be responsible for the benthic community impairment, such as pyrethroids, metals, nutrients, sediment characteristics, and altered physical habitat conditions, were analyzed concurrently in order to identify which stressor or stressors were most

influential. Predictive approaches based on laboratory-derived toxicity data for toxicants (pyrethroids and metals) suggested that these toxicants could be toxic to resident benthic communities. However, a thorough statistical analysis of all stressors based on field data that yields an observed response did not support this prediction. The consistently poor habitat conditions at all sites in the Santa Maria waterbody was not identified in the statistical analysis to be a major contributor to the impaired benthic communities possibly due to a lack of variation in habitat conditions among the sites. This finding is in sharp contrast to other studies conducted in urban streams in California using a similar sampling design where habitat conditions were determined to be a key stressor.

The key constituents identified as major contributors in shaping benthic communities were TOC and phosphorus-based nutrients. In summary, the less stressed, more diverse benthic communities highly populated by tolerant taxa such as oligochaetes tended to be associated more with TOC-rich finer sediments and lower concentrations of phosphorous-based nutrients while the more stressed, less diverse benthic communities tended to be associated with less organically rich, somewhat less fine sediments and higher phosphorous concentrations. Field approaches conducted over multiple years in a waterbody designed to determine the relationships of multiple stressors to resident benthic communities provide the most ecologically relevant assessment of which stressors are most important in shaping benthic communities. Relying solely on single species laboratory toxicity data to predict the impact of specific toxicants on resident benthic communities may provide misleading conclusions if confirmation from a bioassessment approach addressing possible multiple stressor impacts is not determined.

Author Contributions

WK and RA conducted the field sampling. RA conducted the statistical analysis.

REFERENCES

- Pimental D. *World Soil Erosion and Conservation*. Cambridge, UK: Cambridge University Press; 1993.
- Humenik FJ, Smolen MD, Dressing SA. Pollution from non-point sources. *Environ Sci Tech*. 1987;21:737–742.
- Nonpoint Source: Agriculture U. S. Environmental Protection Agency Website. <https://www.epa.gov/nps/non-point-source-agriculture>. Up-dated January 19, 2017. Accessed October 25, 2017.
- Loague KD, Corwin L, Ellsworth TR. The challenge of predicting non-point source pollution. *Environ Sci Tech*. 1998;32:130–133.
- Karr JR, Chu EW. *Restoring Life in Running Waters—Better Biological Monitoring*. Covelo, CA: Island Press; 1999.
- Yoder CO, Rankin ET. Biological criteria program development and implementation in Ohio. In: Davis WD, Simon TP, eds. *Biological Assessment and Criteria: Tools of Water Resource Planning and Decision Making*. Boca Raton, FL: Lewis Publishers; 1995:109–144.
- Barbour MT, Gerritsen J, Griffith GE, et al. A framework for biological criteria for Florida streams using benthic macroinvertebrates. *J N Am Benthol Soci*. 1996;15:185–211.
- Wright JF, Sutcliffe DW, Furse MT. *Assessing the Biological Quality of Freshwater: RIVPACS And Other Techniques*. Ableside Cumbria, UK: Freshwater Biological Association; 2000.
- Giddings JM, Hendley P, Holmes C, et al. *Ecological Risk Assessment of Agricultural Uses of Nine Synthetic Pyrethroids*. Valdosta, GA: Pyrethroid Working Group; 2015. Report. PWG-ERA-18.
- Central Coast Regional Water Quality Control Board. *Adopting Total Maximum Daily Loads for Toxicity and Pesticides in the Santa Maria River Watershed*. San Luis Obispo, CA: California Regional Water Quality Control Board; 2014.
- Hall LW Jr, Anderson RD, Killen WD, Alden RW III. A summary of case studies designed to determine the influence of multiple stressors on benthic communities in urban California streams. In: Jones RL, Shamin M, Jackson SH, eds. *Describing the Behavior and Effects of Pesticides in Urban and Agricultural Settings*. Vol 1168. Washington, DC: American Chemical Society; 2014:135–152.
- Anderson BS, Phillips BM, Hunt JW, et al. Evidence of pesticide impacts in the Santa Maria River watershed, California, USA. *Environ Toxicol Chem*. 2006;25:11160–11170.
- Stillwater Sciences. *Santa Maria River Instream Flow Study: Flow Recommendations for the Steelhead Passage*. Santa Barbara, CA: Stillwater Sciences; 2012. Report prepared for California Ocean Protection Council.
- Smalling KL, Orlando JL. *Occurrence of Pesticides in Surface Water and Sediments from Three Central California Watersheds 2008–09*. Reston, VA: US Geological Survey; 2011. Data Series 600.
- Kazyak PF. *Maryland Biological Stream Survey Sampling Manual*. Annapolis, MD: Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division; 1987.
- American Society for Testing and Materials (ASTM). *Standard Test Method for Particle Size Analysis of Soils (Method D422-63)*. Philadelphia, PA: ASTM; 1998.
- United States Environmental Protection Agency (US EPA). *SW-846, Method 9060A: Total Organic Carbon*. Washington, DC: US EPA Office of Water; 2004. Report.
- Harrington J, Born M. *Measuring the Health of California Streams and Rivers—A Methods Manual for Water Resource Professionals, Citizen Monitors and Natural Resource Students*. Sacramento, CA: Sustainable Land Stewardship International Institute; 2000. Report.
- Barbour MT, Gerritsen J, Snyder BD, Stribling S. *Rapid Bioassessment Protocols for Use in Wadeable Streams and Rivers*. Washington, DC: EPA; 1999. EPA-841-B-99-002.
- Hall LW Jr, Killen WD, Anderson RD, Alden RW III. *An Assessment of Benthic Communities with Concurrent Physical Habitat, Pyrethroids, Metals and Nutrient Analysis in the Santa Maria Watershed in 2015*. Queenstown, MD: University of Maryland, Wye Research and Education Center; 2015. Progress Report.
- Hall LW Jr, Killen WD, Anderson RD, Alden RW III. *An Assessment of Benthic Communities with Concurrent Physical Habitat, Pyrethroids, Metals and Nutrient Analysis in the Santa Maria Watershed in 2016*. Queenstown, MD: University of Maryland, Wye Research and Education Center; 2016. Progress Report.
- Hall LW Jr, Killen WD, Anderson RD, Alden RW III. *An Assessment of Benthic Communities with Concurrent Physical Habitat, Pyrethroids, Metals and Nutrient Analysis in the Santa Maria Watershed in 2017 and 2015–2017*. Queenstown, MD: University of Maryland, Wye Research and Education Center; 2017. Final Report.
- Buchman MF. *NOAA Screening Reference Tables*. Seattle, WA: Coastal Protection and Restoration Division, National Oceanic and Atmospheric Administration; 1999. NOAA HAZMAT Report 99–1.
- Plumb RH. *Procedures for Handling and Chemical Analysis of Sediment and Water Samples*. Vicksburg, MS: US Army Corps of Engineers, US Army Engineer Waterways Experimental Station; 1981. Report.
- Dituro DM, Mahony JD, Hansen DJ, et al. Toxicity of cadmium in sediment: the role of acid volatile sulfide. *Environ Toxicol Chem*. 1990;9:1488–1502.
- Dituro DM, Mahony JD, Hansen DJ, et al. Acid volatile sulfide predicts the acute toxicity of cadmium and nickel in sediments. *Environ Sci Tech*. 1992;26:96–101.
- O'Dell JW. *Determination of Ammonia Nitrogen by Semi-automated Colorimetry (Method 350.1)*. Cincinnati, OH: Environmental Monitoring Systems Laboratory, Office of Research and Development. U. S. Environmental Protection Agency; 1993. Report.
- O'Dell JW. *Determination of Nitrate-nitrite by Automated Colorimetry (Method 353.2)*. Cincinnati, OH: Environmental Monitoring Systems Laboratory, Office of Research and Development. U. S. Environmental Protection Agency; 1993. Report.
- O'Dell JW. *Determination of Phosphorus by Semi-automated Colorimetry (Method 365.1)*. Cincinnati, OH: Environmental Monitoring Systems Laboratory, Office of Research and Development. U. S. Environmental Protection Agency; 1993.
- Reed RL. *Laboratory Validation: Validation of the Residue Analytical Method: Residue Analytical for the Determination of Residues of Bifenthrin, Cypermethrin, Cyfluthrin, Deltamethrin, Efenvalerate, Fenprothrin, Lambda-cyhalothrin and Permethrin in Sediment*. Sacramento, CA: Morse Laboratories, Inc.; 2006. Final Report. Protocol No. MLI-06-02. Morse Labs Project No. ML06-1286-PWG.

31. Moulton SR II, Carter JL, Grotheer SA, Cuffney TF, Short TM. *Methods of Analysis by the U. S. Geological Survey National Water Quality Laboratory—Processing, Taxonomy and Quality Control of Benthic Macroinvertebrate Samples*. Sacramento, CA: US Geological Survey; 2000. Report 2000–212.
32. Amweg EL, Weston DP, You J, Lydy MJ. Pyrethroid insecticides and sediment toxicity in urban creeks for California and Tennessee. *Environ Sci Tech*. 2006;40:1700–1706.
33. Hall LW Jr, Killen WD, Anderson RD, Alden RW III. The relationship of benthic community metrics to pyrethroids, metals, and sediment characteristics in Cache Slough, California. *J Environ Sci Health A Tox Hazard Subst Environ Eng*. 2016;51:154–163. doi:10.1080/10934529.2015.1087744.
34. SAS Institute Inc. *Statistical Analysis Program*. Version 9.1. Cary, NC: SAS Institute Inc.; 2010.
35. United States Environmental Protection Agency (US EPA). *Fact Sheet EPA Ecological Nutrient Criteria*. Washington, DC: US EPA; 2002. EPA-822-F-02-008.
36. Griffith MB, Husby P, Kaufmann PR. Analysis of macroinvertebrate assemblages in relation to environmental gradients among lotic habitats of California's Central Valley. *Environ Monitor Assess*. 2003;82:281–309.
37. Hall LW Jr, Killen WD, Alden RW III. Relationship of farm level pesticide use and physical habitat on benthic community status in a California agricultural stream. *Hum Ecol Risk Assess*. 2007;13:843–869.
38. Rankin ET. Habitat indices in water resource quality assessments. In: Davis WS, Simon TP eds. *Biological Assessment and Criteria: Tools for Water Resource Planning and Decision Making*. Boca Raton, FL: Lewis Publishers; 1995:181–208.
39. Hall LW Jr, Anderson RD, Killen WD. Mapping of depositional and non-depositional areas in Salinas, California streams with concurrent pyrethroid and benthic macroinvertebrate assessments. *J Environ Sci Health A Tox Hazard Subst Environ Eng*. 2013;48:1–13.
40. Holmgren GGS, Meyers MW, Chaney RL, Daniels RB. Cadmium, lead, zinc, copper and nickel in agricultural soils of the United States of America. *J Environ Qual*. 1993;22:335–348.
41. Woodring WP, Bramlette MN. *Geology and Paleontology of the Santa Maria District California*. Washington, DC: United States Government Printing Office; 1950. Geological Survey Professional Paper 222.
42. Kruckeberg AR. Plant species in relation to serpentine soils. *Ecology*. 1954;35:267–274.
43. Chen W, Krage N, Wu L. Arsenic, cadmium, and lead in California cropland soils: role of phosphate and micronutrient fertilizers. *J Environ Qual*. 2008;37:689–695.
44. Ashton MJ, Morgan RP II, Stranko S. Relations between macroinvertebrate, nutrients, and water quality criteria in Wadeable streams of Maryland, USA. *Environ Monit Assess*. 2014;186:1167–1182.
45. Wang L, Robertson DM, Garrison PJ. Linkages between nutrients and assemblages of macroinvertebrates and fish in Wadeable streams: implications to nutrient criteria development. *Environ Manag*. 2007;39:194–212.
46. Maul JD, Farris JL, Milam CD, Cooper CM, Testa S III. The influence of stream habitat and water quality on macroinvertebrate communities in degraded streams of northwest Mississippi. *Hydrobiologia*. 2004;518:79–94.
47. Miracle MR, Moss B, Vincente E, et al. Response of macroinvertebrates to experimental nutrient and fish additions in European localities at different latitudes. *Limnetica*. 2006;25:585–612.
48. Hall LW Jr, Killen WD, Anderson RD, Alden RW III. A three year assessment of the influence of physical habitat, pyrethroids and metals on benthic communities in two urban California streams. *J Ecosys Ecograph*. 2013;3:133. doi:10.4172/21577625.1000133.
49. Rehn AC, Mazor RD, Ode PR. *The California Stream Condition Index (CSCI): A New Statewide Biological Monitoring Tool for Assessing the Health of Freshwater Streams*. Sacramento, CA: State Water Resources Control Board; 2015. SWAMP Technical Memorandum TM-2015-0002.
50. Stribling JB, White JS, Jessup BK. *Development of a Benthic Index of Biotic Integrity for Maryland Streams*. Owings Mills, MD: Tetra Tech, Inc.; 1998. Report of Maryland Department of Natural Resources.
51. Grumiaux F, Lepretre A, Dhainaux-Gourtois N. Effect of sediment quality on benthic macroinvertebrate communities in streams in the north of France. *Hydrobiologia*. 1998;385:33–46.
52. Thompson B, Lowe S. Assessment of macrobenthos response to sediment contamination in the San Francisco estuary, California, USA. *Environ Toxicol Chem*. 2004;23:2178–2187.