

Household Water Treatment Practice and Associated Factors in Rural Households of Sodo Zuria District, Southern Ethiopia: Community-Based Cross-Sectional Study

Authors: Admasie, Amha, Abera, Keefelegn, and Feleke, Fentaw Wassie

Source: Environmental Health Insights, 16(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/11786302221095036>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Household Water Treatment Practice and Associated Factors in Rural Households of Sodo Zuria District, Southern Ethiopia: Community-Based Cross-Sectional Study

Environmental Health Insights
Volume 16: 1–7
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/11786302221095036



Amha Admasie¹ , Kefelegn Abera² and Fentaw Wassie Feleke³ 

¹School of Public Health, Bahir Dar University, Bahir Dar, Ethiopia. ²Teaching and Referral Hospital, Wolaita Sodo University, Bahir Dar, Ethiopia. ³College of Health Science, Woldia University, Wodia, Ethiopia.

ABSTRACT

INTRODUCTION: In Ethiopia, access to safe drinking water is very low, and even safe water at the point of distribution is subjected to frequent and substantial contamination during collection, transport, and storage. The purpose of this study was to assess the level of household water treatment practices and associated factors in rural households of the Sodo Zuria district, southern Ethiopia.

METHODS: A community-based cross-sectional study was conducted in 836 households using a multistage sampling technique. A structured and pre-tested questionnaire was used. Binary and multivariate logistic regression analysis was used.

RESULTS: The household water treatment practice was 44.1%. Households having a higher estimated monthly income, AOR = 1.5 (1.23, 3.47), older age greater than 45 years, AOR = 1.69 (1.08, 2.64), fetching water twice a day, AOR = 2.8 (1.21, 9.17), weekly washing of the water storage container, AOR = 0.3 (0.11, 0.83), and using the dipping technique to draw water from the collection jar, AOR = 1.67 (1.14, 2.42) were significant factors in the practice of household water treatment in the study.

CONCLUSIONS: The household water treatment practice was low. Higher estimated monthly income, older household heads, fetching water twice per day, washing the water storage container weekly, and dipping techniques to draw water from water storage containers were significant factors of household water treatment practices. Thus, proper hygiene of water storage, and engaging the community in income-generating activities were recommended.

KEYWORDS: Ethiopia, household water treatment practices, rural households

RECEIVED: November 23, 2021. **ACCEPTED:** March 28, 2022.

TYPE: Original Research

FUNDING: The author(s) received no financial support for the research, authorship, and/or publication of this article.

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Fentaw Wassie Feleke, College of Health Science, Woldia University, P.O.BOX: 400, Wodia, Ethiopia. Email: fentawwassie@gmail.com

Introduction

Globally, the common household water treatment technologies are chlorination, filtration, solar disinfection and boiling,^{1,2} and boiling is the most frequently used method.^{2,3} Household water treatment and safe storage is the term used to describe the process of treating and safely storing water in the home.⁴ In general, it is a way of lowering disease-causing pathogens⁵ and improving the quality of drinking water at the household level.⁶ Worldwide, more than 1 billion people lack access to drinking water, which is equal to 17% of the global population.⁴

It is estimated that globally every day, the diarrheal disease kills 2195 children, outnumbering AIDS, malaria, and measles combined which makes it the second leading cause of death in children under 5 years of age.⁷ A significant proportion of diarrheal disease can be prevented through safe drinking and adequate water sanitation and hygiene (WASH).^{8,9}

Copious studies from low-income and middle-income populations throughout the world have reported that indicators of household-level WASH are significant predictors of linear growth of children after adjusting for a wide array of other

covariates including birth weight, gestational age at birth, maternal height, maternal education, and household wealth.¹⁰ However, a wide range of studies indicated that HWTS can improve drinking water quality prior to consumption and it has been found as a cost-effective method that can reduce the risk of diarrhea significantly.^{11–14} Even though the benefit of household water treatment practice to decrease the risk of enteric infection is well understood in different studies, there are different hurdles that prevent consistent practice; which include psychosocial, contextual, and technology-related factors.^{15,16}

Although tens of millions of people rely on improved community water supplies, water supply authorities may not deliver microbiologically qualified water, so water needs to be treated at the household level.^{17,18} In low-income countries, numerous household water treatments (HWT) have been confirmed to significantly improve drinking water quality in field trials.¹⁹ It was reported that there was more than 25% reduction in diarrheal disease among children <2 years of age using flocculent disinfectant compared to untreated water handling methods.²⁰ Waterborne diseases have a harmful effect on public health



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

when drinking water is of poor quality²¹ and due to a problem in the universal provision of treated and piped water, and HWT technology is lacking.^{6,22,23} Common sources of drinking water in low-income countries are shallow wells, ponds, streams, rivers, and lakes, which are extremely contaminated with fecal pathogens needs HWT.²⁴

Evidence from the Demographic and Health Survey of sub-Saharan Africa also showed that only 18% of households treat their drinking water appropriately.²⁵ In Ethiopia, various studies found that efforts to minimize water-borne diseases by treating drinking water at the source were insufficient to reduce water-borne diseases in the country unless drinking water was treated and sanitary handling is practiced at home.²⁶

In Ethiopia, access to safe drinking water is very low, and even safe water at the point of distribution is subjected to frequent and substantial contamination during collection, transport, and storage. Meanwhile, conventional water treatment plants are scarce, and even existing plants are vulnerable to frequent interruptions and technical malfunctions.²² Even improved water sources often fail to provide safe drinking water due to an infusion of fecal contamination in the course of the distribution system and mode of use.²⁷⁻²⁹ Treatment of point of use water at the household level, such as boiling and chlorination, is the most cost-effective intervention to prevent diarrheal disease. The effect of chlorine-treated household water treatment on the reduction of diarrhea episodes is variable, ranging from no protective effect to 85% reduction.⁴ However, household water treatment is widespread throughout the world but in Ethiopia a maximum of 6% and below it of the households practiced HWT.^{30,31}

The United Nations program set the SDG 6.1 universal and equitable access to safe and affordable drinking water for all by the end of 2030.³² Despite increasing efforts to promote household water treatment in Ethiopia through routine health extension program, the reasons for the low practice are not yet clear. In addition to the limited research done on this regard, there is no conclusive and consistent evidence on factors that determine the practice of household water treatment among rural households. It will also enrich literatures available on household water treatment practice and may trigger other researcher to conduct related studies in various parts of the globe. Therefore, the main objective of this study was to evaluate the current status of household water treatment practices and associated factors in rural households in Sodo Zuria District, South Ethiopia.

Methods

Study setting: The study was carried out in rural households in the Sodo Zuria district, southern Ethiopia, from August 1st to September 30, 2018. The Sodo Zuria district is located in the Wolaita zone 329 kilometers south of Addis Ababa, the capital of Ethiopia. The district has 36 Kebeles (sub-districts/lowest administrative unit in Ethiopian context) and 35231 households. The district had also about a total of 172632 populations.

The common sources of household water supply in the district are unimproved water sources such as a river, well, and spring water.³³

Study design and population: A community-based cross-sectional study design was used. All households in the district were the source population and all randomly selected households from the selected kebeles were the study population. Respondents whose age is 18 years and above (preferably female), and who have been living for at least 6 months in the district were included in the study. Whereas, respondents who were seriously ill and could not communicate to give information were excluded from the study.

Sample size: A single population proportion formula was used considering the proportion of HWT practices from a study conducted in north-west Ethiopia which is 44.8%.⁵ With an assumption of 95% confidence level, 5% margin of error, design effect of 2 and an anticipated nonresponse of 10% were considered. Based on the above assumptions, using the single proportion formula, the total sample size was found to be 836.

Sampling procedure: A multistage sampling procedure was used by first selecting 8 Kebeles from 36 Kebeles using the simple random sampling method. Based on the proportional allocation, households were distributed in each Kebele. The households of study participants were selected using systematic sampling techniques after calculating the sampling interval from the 8 kebeles number of household divided by the number of study sample size households which was $6688/836=8$. Then the data were collected every eighth household.

Variables

Outcome variable: The outcome variable for this study was household water treatment practice (Yes/No) which is dictated as "Yes" if at least one of the following options is practiced at the household; such as boil, add bleach/chlorine, strain through a cloth, use water filter, solar disinfection, let it stand and settle, and it is measured by self-report.^{4,34}

Explanatory variables: The explanatory variables included socio-demographic (Age, sex, educational status, and religion, marital status, occupational status, family size, head of a household, and monthly income), water supply, storage, and hygiene factors.

Data collection tool and procedure: Data were collected using a pretested and structured questionnaire in a face-to-face interview of household heads. The questionnaire was adapted based on the available literature. The questionnaire was originally prepared in English and then translated into Amharic. Finally, it was translated back to English to check for consistency. Training was given to 5 data collectors and 1 supervisor for 2 days. A pre-test was conducted on 5% of the sample size in Kebeles which were not included in the actual data collection. Following the pre-test result, the tool and procedures were modified for ease of convenience and to ensure data quality. Data collection procedures were genuinely supervised to check

the consistency, clarity, and completeness of the questionnaires filled daily during data collection.

Operational definitions

Household water treatment practices: Households who used at least one of the following household water treatment methods before drinking: boiling, adding bleach/chlorine, letting the water pass through a clean cloth (filtering), solar disinfection (SODIS), and settling in the last 2 weeks at least once before drinking.⁴

Data management and analysis procedure: First, the questionnaires were checked for completeness, then coded and entered into the computer using the Epidata3.1 statistical software, then cleaned and exported to SPSS 20 statistical software for analysis. Data were presented in descriptive statistics, tables, and charts. Binary logistic regression analysis was used to screen the eligible independent variables effect on the outcome variable at P -values $<.25$. The necessary assumption of logistic regression was checked using Hosmer and Lemeshow goodness-of-fit-test statistics to assess the fitness of the model. Collinearity was checked using a cutoff point based on the variance inflation factor (VIF) <10 or the tolerance test >0.1 . To control for confounding variables in the bivariate analysis, a multivariate analysis was run. In the final model, the level of significance of the variables was determined at P -value $<.05$.

Results

Socio-demographic characteristics of the respondents

Eight hundred and thirty-three respondents had participated in the study with a response rate of 99.6%. The mean age of the respondents was 38.22 and (SD \pm 8.2) years. Among the total of the respondents, 625 (75%) were women and 702 (84.3%) were married. Regarding educational status, 403 (44.8%) of them had no formal education and the majority 626 (75.2%) of the subjects were farmers in occupation (Table 1).

Water handling practices. About 344 (41%) participants had reported that they stored water in 2 containers and 378 (45.4%) of the household containers had a capacity of <20 L. The majority (774 (90.9%)) of the respondents used covered water containers for water storage. Two-third, 116 (65%) of the respondents reported that the government water supply system was interrupted. Regarding water sources, more than half of the household, 524 (62.9%) had an unimproved source (Table 2).

Household water treatment practice. About 367 (44.1%) of the respondents had used different household water treatment methods to treat water before drinking. Concerning the mode of water withdrawal from the storage container, approximately 449 (53.9%) had drawn water by dipping (during data collection) from the storage container and the majority 757 (90.9%) of them had washed their water storage container. The water

Table 1. Socio-demographic characteristics of respondents, Sodo Zuria district, South, Ethiopia, 2018 (n=833).

VARIABLES	RESPONSES	FREQUENCY	PERCENT
Sex	Female	625	75
	Male	208	25
Age	18-35 y	323	38.8
	36-45 y	351	42.1
	>45 y	159	19.1
Educational status	No formal education	403	48.4
	Literate	430	51.6
Occupation	Farmer	626	75.2
	Housewife	73	8.8
	Merchant	80	9.6
	Daily laborer	53	6.4
Marital status	Single	64	7.7
	Married	702	84.3
	Divorced/ widowed	67	8.0
Family size	≤ 5	583	70.0
	>5	240	30.0
Monthly income (ETB)	≤ 500	327	39.2
	501-999	382	45.9
	≥ 1000	124	14.9

treatment practice reported by the study respondents 245 (29.5%) had used boiling as their water treatment option prior to drinking. About 84 (10.1%) and 38 (4.5%) had practiced adding chlorine/bleach and settling, respectively. More than half of the 466 households (54.9%) had never used any method to have safe drinking water (Table 3).

Factors associated with household water treatment practice. During bivariate logistic regression analysis, sex, age, education, occupation, monthly income, frequency of fetching water per day, ways of drawing water, and frequency of washing the water storage container were screened as candidate independent variables for multivariable logistic regression analysis with HWT practice with P -value $<.25$. During multivariate analysis, monthly income, age, the frequency of fetching water, the frequency of washing water containers, and mode of drawing water maintain their significant association with HWT practice at P -value $<.05$.

Households with estimated income greater than 1000.00 ETB are 1.5 times more likely to practice HWT than those

Table 2. Water handling practices in Sodo Zuria district Southern, Ethiopia, 2018 (n=833).

VARIABLES	RESPONSES	FREQUENCY	PERCENT
Number of containers used to store water	1	224	26.6
	2	344	41.3
	3 and more	265	32.1
A capacity of the container in liters	≤20L	259	31.1
	>20L	574	69.9
Do your water container have a cover	Yes	757	90.9
	No	76	9.1
A material used to wash the container (n=806)	Water with soap	377	46.8
	Water only	429	53.2
A container used to fetch water	Jerry can	581	69.9
	Pot	212	25.5
	Plastic container	38	4.6
A frequency of fetching water per day	1	576	69.1
	2 or more	257	30.9
Nature of government water supply (n=176)	Continuous	60	34.1
	Intermittent	116	65.9
Water sources	Unimproved	524	62.9
	Improved	309	37.1

with income <500 ETB, AOR=1.5 (95% CI=1.23-3.47). Those who had an age greater than 45 years were 1.69 times more likely to practice HWT than 18 to 35 years, AOR=1.69 (95% CI=1.08, 2.64). Households who collected water twice a day were 2.8 times more likely to practice HWT than households who drank more than twice, AOR=2.8 (95% CI=1.21-9.17). The weekly washing of the container has 30% higher odds of HWT practice than those who washed their water storage container daily, AOR=0.3 (95% CI=0.11, 0.83). Using a dip method to draw water from the collection jar was 1.67 times more likely to practice HWT than who used pouring, AOR=1.67 (95% CI=1.14, 2.42) (Table 4).

Discussion

According to the current study, 44.1% of study participants practiced HWT at least once during 2 weeks before the data collection. Even though the current finding showed that 44.7% and 50.9% of drinking water sources in the region were contaminated with *Escherichia coli* and enterococci respectively³⁵ the level of household water treatment practice was found to be 44.1%. This finding is consistent with the study done Northwest Ethiopia (44.8%),⁵ Nigeria (45%)³⁶ whereas it was lower than

Table 3. Household water treatment practices in rural households of Sodo Zuria district, Southern, Ethiopia, 2018 (n=833).

VARIABLES	RESPONSE	FREQUENCY	PERCENT
Methods of water drawing from storage container	Dipping	604	72.5
	Pouring	229	27.5
Do you wash water storage container?	Yes	757	90.9
	No	76	9.1
A frequency of washing the container (n=757)	Daily	385	50.9
	2-3d interval	244	32.2
	Weekly	128	16.9
Do you store water for more than 3d?	Yes	450	54
	No	383	46
Type of water storage container	Jerry can	536	64.3
	Pot	246	29.5
	Plastic container	40	4.8
	Iron container	11	1.4
Do you treat water prior to drinking?	Yes	367	43.9
	No	466	56.1
Time taken to fetch water from water sources	<30 min	669	80.4
	≥30 min	164	19.6
HWT methods	Boiling	245	29.5
	Chlorine bleach	84	10.1
	Settling	38	4.5
	Never used	466	55.9

studies revealed from Zambia (72.6%), Peru 70.9%,³⁷ and a study in Western Pacific region (66.8%).²⁷ But it was higher than the findings from Ethiopian demographic health survey 2016 which is 7%,³⁴ Southern Ethiopia Gibe district revealed 34.3%,²⁶ northern Ethiopia Degadamot district 14%,³⁸ and in developing countries 33.3%.³⁹ It was also higher than studies revealed from outside Ethiopia such as in Biye community, Kaduna State of Nigeria 32.4%, in Eastern Mediterranean region 13.6%,²⁷ and in African region 18.2%.³⁷ The possible explanations for this difference might be related with households' perception difference across different contexts. The discrepancy from these reports might be explained that the report included all regional states with different socio-demographic, socioeconomic background, related to sample size, study design, and study period variations.

Households with higher income were 1.5 times more likely to practice HWT than those with low income. This finding is in line with the finding of Maria Elena Figueroa

Table 4. Factors associated with household water treatment practices in rural households of Sodo Zuria district, southern, Ethiopia, 2018 (n=833).

VARIABLES		HWT PRACTICE		COR (95% CI)	AOR (95% CI)
		YES	NO		
Sex	Male	107	101	1.49 (1.09, 2.04)	1.16 (0.8, 1.68)
	Female	260	365	Reference value	Reference value
Age in years	18-35	129	194	Reference value	Reference value
	36-45	160	191	1.26 (0.93, 1.71)	1.44 (0.9, 2.03)
	>45	78	81	1.45 (0.99, 2.12)	1.69 (1.08, 2.64)
Educational	Literate	197	233	1.16 (0.88, 1.52)	1.35 (0.98, 1.87)
	Illiterate	170	233	Reference value	Reference value
Occupation	Merchant	112	118	Reference value	Reference value
	Farmer	87	86	1.07 (0.72, 1.58)	1.4 (0.89, 2.18)
	House wife	107	141	0.8 (.56, 1.15)	1.07 (0.71, 1.62)
	Daily labor	61	121	0.53 (0.35, 0.79)	0.76 (0.49, 1.19)
Estimated income	≤500 ETB	128	199	Reference value	Reference value
	501-999 ETB	168	214	1.22 (0.90, 1.65)	1.2 (0.87, 1.64)
	≥1000 ETB	53	71	2.08 (1.37, 3.17)	1.5 (1.23, 3.47)
Frequency of fetching water per day	Once	275	301	4.72 (1.94, 11.49)	3.33 (0.01, 7.84)
	Twice	134	85	3.28 (1.31, 8.12)	2.8 (1.21, 9.17)
	Above 2	6	31	Reference value	Reference value
Water storage (>3d)	Yes	182	267	1.36 (1.04, 1.80)	1.3 (0.96, 1.77)
	No	185	199	Reference value	Reference value
Frequency of washing water storage containers	Daily	204	197	Reference value	Reference value
	2-3d	113	160	0.68 (0.50, 0.93)	0.65 (0.46, 0.91)
	Weekly	39	93	0.41 (0.27, 0.62)	0.3 (0.11, 0.83)
Methods of water drawing	Dipping	293	311	1.97 (1.45, 2.76)	1.67 (1.14, 2.42)
	Pouring	74	155	Reference value	Reference value

and D. Lawrence Kincaid.^{40,41} This study was confirmed with the evidence revealed from Mexico groups the lowest family income households compared together the highest income households distribution shows that water consumption is favorable to a larger extent to the households with the highest income.⁴² This was also consistent a study revealed in China⁴³ and Nigeria.³⁶

Respondents with higher age were 1.69 times more likely to practice HWT than household heads of lower age. A similar finding was reported by Maria Elena Figueroa.⁴⁰ The possible explanation for this finding might also be that older age participants may have understood through exposure the need for household water treatment to reduce diarrheal disease, and their educational level³⁸ may also encourage practice.

This study revealed that participants who draw their water from storage vessel by dipping were 1.67 times more likely to practice household water treatment than those who draw their water by pouring. This finding is in line with the study done in Northern and southern Ethiopia.^{5,26} The possible explanation might be due to the fact that those who withdraw water from the storage container by dipping thought that dipping the container for drawing water increases the risk of contamination, and they might get information from health professionals on the possible ways of water contamination. Since hands can enter the jar, causing water contamination and to avoid those contaminants, respondents may employ either of the methods for HWT. Apparently, those who pour water and store it in narrow-neck containers may be more protective against contamination.³⁷

Households who fetched water twice per day were 2.8 times more likely to practice HWT than households who fetched water more than twice per day. This is in agreement with a study from north-west Ethiopia^{5,26} and Nigeria.³⁶ The possible explanation might be due to the fact that those who were fetching water more frequently might have higher tendency to store their water, which in turn empowers them to treat their water, by storing.

Respondents who washed their container weekly have 30% lowered odds of HWT practice than those who washed their container daily. The reason may be that the person who washed daily may think that for a daily washed water storage container, contaminants are less likely in the water contamination. But the CDC report revealed that only washing the water storage container is not guaranteed for the absence of disease-causing organisms in water.⁴⁴ Generally household water treatment practice has a range of multilevel influences. Beyond the model of providing on-going safe water education by health extension workers, potential initiatives could be improved by community mobilization activities that include community leaders, women's groups, etc., in promoting water treatment at community engagements. This is confirmed by a qualitative analysis of factors influencing household water treatment practices among consumers of self-supplied water in rural Ethiopia.⁴⁵

Limitation of the Study

The current study used a large sample size compared to previous studies. As a limitation, since the information for this study was collected mainly through interviews and self-reports, there is a possibility that some of the responses might suffer from social desirability bias and to avoid this we add an observation method as a method of data collection besides the interview. Moreover, since the study employed cross sectional study design, it may be difficult to establish temporal relationship between the outcome and response variables. In addition, it could have been better if the water bacteriological analysis were considered in the method.

Conclusion

The HWT practice was low. Having a higher estimated monthly income, being of older age, fetching water twice per day, washing water storage container weekly, and using dipping method to draw water from the container were determinant factors of HWT practice. Therefore, dipping method drawing water, water storage hygienic practice, and strengthening income generating activities are highly recommended. Further scientific studies through qualitative methods were also recommended.

Acknowledgements

Our deepest appreciation goes to Wolaita Sodo University; Sodo Zuria district health office; all data collectors, supervisors, and study participants who participated in the study.

Author Contributions

KA designed the study protocol. KA and AA involved in protocol development. KA involved in data collection and analysis. KA and AA involved in data analysis. AA prepared the manuscript and edited by FWF. AA supervised and monitored the protocol. All authors read and approved the final manuscript.



Availability of Data and Materials

The data sets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethical Considerations

Ethical approval and clearance was obtained from the Institutional Ethics Review Committee (IRC) of Wolaita Sodo University, School of Health Sciences and Medicine, School of Public Health. Permission was also obtained from the concerned bodies of Sodo Zuria district offices. Written and verbal consent was obtained from the study participants. Privacy and confidentiality were maintained. Participants were told that they have the right to participate and/or withdraw any time they feel uncomfortable by the interview.

ORCID iDs

Amha Admasie  <https://orcid.org/0000-0002-9826-1270>
Fentaw Wassie Feleke  <https://orcid.org/0000-0002-4482-877X>

REFERENCES

- Enger KS, Nelson KL, Rose JB, Eisenberg JN. The joint effects of efficacy and compliance: a study of household water treatment effectiveness against childhood diarrhea. *Water Res.* 2013;47:1181-1190.
- Cotruvo JA, Sobsey MD. Point-of-use water treatment for home and travel. *Water Health.* 2009;2:103.
- Cohen A, Tao Y, Luo Q, et al. Microbiological evaluation of household drinking water treatment in rural China shows benefits of electric kettles: a cross-sectional study. *PLoS One.* 2015;10:e0138451.
- World Health Organization. *A Toolkit for Monitoring and Evaluating Household Water Treatment and Safe Storage Programmes.* WHO; 2012.
- Belay H, Dagnew Z, Abebe N. Small scale water treatment practice and associated factors at Burie Zuria Woreda rural households, Northwest Ethiopia, 2015: cross sectional study. *BMC Public Health.* 2016;16:887-888.
- Mengistie B, Berhane Y, Worku A. Household water chlorination reduces incidence of diarrhea among under-five children in rural Ethiopia: a cluster randomized controlled trial. *PLoS One.* 2013;8:e77887.
- Prevention C. *Diarrhea: Common Illness Global Killer.* US Department of Health and Human Services; 2013.
- Soboksa NE. Associations between improved water supply and sanitation usage and childhood diarrhea in Ethiopia: an analysis of the 2016 demographic and health survey. *Environ Health Insights.* 2021;15:11786302211002552.
- Ko SH, Sakai H. Water sanitation, hygiene and the prevalence of diarrhea in the rural areas of the delta region of Myanmar. *J Water Health.* 2022;20:149-156.
- Pickering AJ, Null C, Winch PJ, et al. The WASH benefits and SHINE trials: interpretation of WASH intervention effects on linear growth and diarrhoea. *Lancet Glob Health.* 2019;7:e1139-e1146.
- Verhoughstraete M, Reynolds KA, Pearce-Walker J, Gerba C. Cost-benefit analysis of point-of-use devices for health risks reduction from pathogens in drinking water. *J Water Health.* 2020;18:968-982.
- Mokoena MM, Mudau LS, Mokgobu MI, Mukhola MS. The use of sodium hypochlorite at point-of-use to remove microcystins from water containers. *Toxins.* 2021;13:207.

13. Hill CL, McCain K, Nyathi ME, et al. Impact of low-cost point-of-use water treatment technologies on enteric infections and growth among children in Limpopo, South Africa. *Am J Trop Med Hyg*. 2020;103:1405-1415.
14. Alemu BM, Berhane Y, Yalew AW. Household water chlorination reduces incidence of diarrhea among under five in rural Ethiopia: a cluster randomized controlled trial. In: *PAA 2017 Annual Meeting*; April 27-29, 2017; Chicago, IL.
15. Daniel D, Sirait M, Pande S. A hierarchical Bayesian belief network model of household water treatment behaviour in a suburban area: a case study of Palu—Indonesia. *PLoS One*. 2020;15:e0241904.
16. Lilje J, Kessely H, Mosler H-J. Factors determining water treatment behavior for the prevention of cholera in Chad. *Am J Trop Med Hyg*. 2015;93:57-65.
17. Clasen T. Household water treatment and safe storage to prevent diarrheal disease in developing countries. *Curr Environ Health Rep*. 2015;2:69-74.
18. Banerjee A, McFarland DA, Singh R, Quick R. Cost and financial sustainability of a household-based water treatment and storage intervention in Zambia. *J Water Health*. 2007;5:385-394.
19. Clasen T, Nadakatti S, Menon S. Microbiological performance of a water treatment unit designed for household use in developing countries. *Trop Med Int Health*. 2006;11:1399-1405.
20. Dubois AE, Crump JA, Keswick BH, et al. Determinants of use of household-level water chlorination products in rural Kenya, 2003-2005. *Int J Environ Res Public Health*. 2010;7:3842-3852.
21. Mwabi JK, Mamba BB, Momba MN. Removal of *Escherichia coli* and faecal coliforms from surface water and groundwater by household water treatment devices/systems: a sustainable solution for improving water quality in rural communities of the Southern African development community region. *Int J Environ Res Public Health*. 2012;9:139-170.
22. Dessie A, Alemayehu E, Mekonen S, Legesse W, Kloos H, Ambelu A. Solar disinfection: an approach for low-cost household water treatment technology in southwestern Ethiopia. *J Environ Health Sci Eng*. 2014;12:25.
23. Jain S, Sahanoon OK, Blanton E, et al. Sodium dichloroisocyanurate tablets for routine treatment of household drinking water in periurban Ghana: a randomized controlled trial. *Am J Trop Med Hyg*. 2010;82:16-22.
24. Mohamed H, Clasen T, Njee RM, Malebo HM, Mbuligwe S, Brown J. Microbiological effectiveness of household water treatment technologies under field use conditions in rural Tanzania. *Trop Med Int Health*. 2016;21:33-40.
25. Geremew A, Damtew YT. Household water treatment using adequate methods in sub-Saharan countries: evidence from 2013–2016 demographic and health surveys. *J Water Sanit Hyg Dev*. 2020;10:66-75.
26. Tafesse B, Gobena T, Baraki N, Alemeshet Asefa Y, Adare Mengistu D. Household water treatment practice and associated factors in Gibe district southern Ethiopia: a community based cross-sectional study. *Environ Health Insights*. 2021;15:11786302211060150.
27. Clasen TF. Household water treatment and the millennium development goals: keeping the focus on health. *Environ Sci Technol*. 2010;44:7357-7360.
28. Freeman MC, Trinies V, Boisson S, Mak G, Clasen T. Promoting household water treatment through women's self help groups in rural India: assessing impact on drinking water quality and equity. *PLoS One*. 2012;7:e44068.
29. Murray A, Pierre-Louis J, Joseph F, Sylvain G, Patrick M, Lantagne D. Need for certification of household water treatment products: examples from Haiti. *Trop Med Int Health*. 2015;20:462-470.
30. Geremew A, Mengistie B, Mellor J, Lantagne DS, Alemayehu E, Sahilu G. Appropriate household water treatment methods in Ethiopia: household use and associated factors based on 2005, 2011, and 2016 EDHS data. *Environ Health Prev Med*. 2018;23:46-11.
31. Azage M, Motbainor A, Gedamu G. Access to improved water and household water treatment practice in rural communities of Amhara Region, Ethiopia. *PAMJ-One Health*. 2021;6:4.
32. United Nations Development Programme. *Sustainable Development Goals. United Nations Sustainable Development*. United Nations Development Programme; 2015.
33. Sodo Zuria District Water and Health Office Departments. *Annual Report of Sodo Zuria Water, Mine, and Energy Office and Health Department, 2018*. Sodo Zuria District Water and Health Office Departments; 2018.
34. ICF, CSA. *Ethiopia Demographic and Health Survey 2016*. CSA and ICF; 2016. Accessed March 5, 2020. <https://dhsprogram.com/pubs/pdf/FR328/FR328.pdf/>
35. Alemayehu TA, Weldetinsae A, Dinssa DA, et al. Sanitary condition and its microbiological quality of improved water sources in the Southern Region of Ethiopia. *Environ Monit Assess*. 2020;192:319.
36. Abubakar IR. Understanding the socioeconomic and environmental indicators of household water treatment in Nigeria. *Utilities Policy*. 2021;70:101209.
37. Rosa G, Kelly P, Clasen T. Consistency of use and effectiveness of household water treatment practices among urban and rural populations claiming to treat their drinking water at home: a case study in Zambia. *Am J Trop Med Hyg*. 2016;94:445-455.
38. Tsegaye D, Aniley Y, Negese B, Mengesha Z. Assessment of knowledge and practice of household water treatment and associated factors in rural kebeles of Degadamat woreda, north-West Ethiopia, 2020. *J Bacteriol Parasitol*. 2021;12:403.
39. Daniel D. Factors influencing household water treatment adoption in rural areas in developing countries; 2021. <https://doi.org/10.4233/uiid:6f6e7a1b-65ac-4876-9531-24988a563e36>
40. Elena Figueroa M, Kincaid DL. *Social, Cultural and Behavioral Correlates of Household Water Treatment and Storage*. Center Publication HCI 2010-1: Health Communication Insights, Johns Hopkins Bloomberg School of Public Health, Center for Communication Programs; 2010.
41. Stoler J, Miller JD, Brewis A, et al. Household water insecurity will complicate the ongoing COVID-19 response: evidence from 29 sites in 23 low- and middle-income countries. *Int J Hyg Environ Health*. 2021;234:113715.
42. Morales-Novelo J, Rodríguez-Tapia L, Revollo-Fernández D. Inequality in access to drinking water and subsidies between low and high income households in Mexico City. *Water*. 2018;10:1023.
43. Liao X, Chai L, Liang Y. Income impacts on household consumption's grey water footprint in China. *Sci Total Environ*. 2021;755:142584.
44. CDC/UNAIDS. *Household Water Treatment Options in Developing Countries: Boiling*. CDC/UNAIDS; 2009.
45. Tamene A. A qualitative analysis of factors influencing household water treatment practices among consumers of self-supplied water in rural Ethiopia. *Risk Manag Healthc Policy*. 2021;14:1129-1139.