

Whose Habitat Is It Anyway? Role of Natural and Anthropogenic Habitats in Conservation of Charismatic Species

Authors: Kumar, Mavatur Ananda, Vijayakrishnan, Sreedhar, and Singh, Mewa

Source: Tropical Conservation Science, 11(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/1940082918788451>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Whose Habitat Is It Anyway? Role of Natural and Anthropogenic Habitats in Conservation of Charismatic Species

Tropical Conservation Science
 Volume 11: 1–5
 © The Author(s) 2018
 Reprints and permissions:
sagepub.com/journalsPermissions.nav
 DOI: 10.1177/1940082918788451
journals.sagepub.com/home/trc



Mavatur Ananda Kumar¹, Sreedhar Vijayakrishnan^{1,2}, and Mewa Singh³

Abstract

Developmental activities have been one of the major drivers of conversion of natural forest areas into mosaics of forest fragments, agriculture, and plantations, threatening the existence of wildlife species in such altered landscapes. Most conservation research and actions are protected area centric and seldom addresses the importance of landscape matrices around these protected areas in providing habitats to a wide range of species. In this article, we bring out the crucial role of natural and anthropogenic habitats for the existence of three charismatic species, namely, Asian elephants, leopard, and lion-tailed macaques. The larger public perception of where the animals should be and where the animals actually are is also discussed. We emphasize that, while habitat generalists often adapt behaviorally and ecologically to modified landscapes, habitat specialists, such as the lion-tailed macaques could find survival harder, with increasing anthropogenic pressures and loss of their habitats.

Keywords

anthropogenic habitats, generalists, habitat use, mammals, protected area, specialists

Globally, burgeoning human developmental activities have threatened natural forest areas, resulting in destruction and fragmentation of habitats and in turn restricting wild fauna within islands of forests or pushing them into surrounding human-use areas. Much of the focus on wildlife conservation has been on the *fortresses of conservation*, protected areas (PAs), which focuses on setting aside wildlife sanctuaries and national parks. However, today, wide range of taxa extensively range outside these conventional protected boundaries, into human-use landscapes, and there has been little attention paid to the ecology of such wildlife that uses these modified habitats dominated by commercial and subsistence agriculture or plantations (see Figure 1). It is only in the recent past that conservation of such species outside PAs has gained momentum and is established as an important area of scientific investigation (Graham, Douglas-Hamilton, Adams, & Lee, 2009; Mudappa, Kumar, & Raman, 2014).

Continued persistence of wildlife in human-use areas warrants ecological adaptation in terms of use of diverse habitats and behavioral modifications in species as a response to pressures associated with modified

landscapes. Such spillover animal populations, particularly wide-ranging species such as elephants, leopards, primates, and so on, outside forest areas come into frequent contact with people, leading to what is largely referred to as human–wildlife conflict.

India, being one of the largest populated countries with 1.3 billion people, still remains as high biodiversity area, where several taxa thrive in human-dominated areas that surround natural habitats (Sridhar, Raman, & Mudappa, 2008; Chang, Karanth, & Robbins, 2018). In the Western Ghats of India, 25% of forest cover

¹Nature Conservation Foundation, Mysore, Karnataka, India

²National Institute of Advanced Studies, Indian Institute of Science Campus, Bangalore, Karnataka, India

³Biopsychology Laboratory and Institution of Excellence, University of Mysore, Mysore, Karnataka, India

Received 18 June 2018; Accepted 21 June 2018

Corresponding Author:

Mavatur Ananda Kumar, Rainforest Restoration Research Station, Nature Conservation Foundation, 8/364, Cooperative Colony, Valparai 642127, Tamil Nadu, India.
 Email: anand@ncf-india.org

loss was estimated between 1973 and 1995, resulting in degradation of forests and increased conversion to agriculture and plantations (Jha, Dutt, & Bawa, 2000). In southern India alone, nearly 10% of available forest area has been reduced due to monoculture plantations such as tea, coffee, *Eucalyptus*, and teak (Santiapillai & Jackson, 1990). Nevertheless, forest remnants among plantations still support wide range of endangered and endemic charismatic species in certain key and important high biodiversity areas in the Western Ghats (Sridhar et al., 2008; Mudappa et al., 2014). Their existence in such altered landscapes is influenced by many factors such as resource availability and distribution, land-use patterns, cultural and religious practices, beliefs of the local communities, and the kind of interactions the species has with people. Semiurban and urban peripheries, off late, provide a unique ecosystem to the species, changing their food habits, distribution, and their physiological processes (Sinha & Vijaykrishnan, 2017).

This article is an attempt to highlight the importance of natural and modified landscapes for the sustenance of three species, namely, the Asian elephant (*Elephas*

maximus), leopard (*Panthera pardus*), and the lion-tailed macaque (LTM; *Macaca silenus*). The Asian elephant, a highly endangered species, uses diverse habitats encompassing forested and nonforested production landscapes across its range. India holds the largest Asian elephant population spreading across 13 Indian states. A recent study by Madhusudan et al. (2015) indicates that more than 60% of Asian elephant population distribution lies outside PAs in the southern Indian state of Karnataka alone. Asian elephants are known to use commercial plantations such as tea, coffee, rubber, cardamom, and oil palm plantations, which are often juxtaposed with elephant landscapes. These habitats are extremely important for movement of elephants; and often, they act as refuges, which provide food and shelter in modified landscapes (Kumar, Mudappa, & Shankar Raman, 2010; Srinivasaiah, Anand, Vaidyanathan, & Sinha, 2012). On the other hand, elephants in hard-edge interfaces of forest and farm landscapes tend to depend on crops, leading to intense conflict between people and elephants.

Similarly, leopards (Athreya, Odden, Linnell, Krishnaswamy, & Karanth, 2013; Navya, Athreya,




Figure 1. Schematic representation of the interrelationship between natural and anthropogenic habitats.

Mudappa, & Raman, 2014; Sidhu, Raman, & Mudappa, 2015; Kshetry, Vaidyanathan, & Athreya, 2017) are known to adapt to human-use landscapes, dominated by tea and coffee plantations, sugarcane fields in villages, and are found even in urbanized spaces, coming into frequent contact with people (see Figure 2). However, these frequent interactions need not necessarily imply incidences of conflict, as indicated by Kshetry et al. (2017), where the authors show that while leopards use ground cover in tea habitats extensively, this is not significantly correlated with conflict occurrence. Abundant wild prey presence in certain modified landscapes such as tea-coffee-forest-fragmented matrix of Valparai plateau in the Anamalai Hills, leopard diet primarily consists of wild prey such as Indian muntjac, Indian spotted chevrotain, sambar deer, Indian porcupine, and so on, as against domestic prey (Sidhu et al., 2015). On the other hand, in large expanse of agricultural habitats with high human density and low wild prey availability, leopards are largely dependent on domestic animals such as dogs, goats, cattle, cats, and so on, which constitute more than 87% of their diet (Athreya et al., 2013). Nevertheless, persistence of large carnivore populations in altered habitats depends on effective conflict avoidance rather than management measures and its enforcement. This also ensures that large carnivore conservation is possible in modified landscapes, thereby enhancing coexistence in such landscapes (Athreya et al., 2013).

On the other hand, species such as LTM, an endangered habitat specialist, listed as Schedule I (highest order of protection in the country) animal under the Indian Wildlife Protection Act, is found only in the narrow strips of rainforests along the Western Ghats of India. Due to exploitation of rainforests to establish commercial plantations such as tea, coffee, rubber, logging, and degradation, many LTM groups are found in fragmented rainforest habitats across its range (Singh, Kaumanns, Sushma, & Molur, 2009; Umaphathy, Hussain, &

Shivaji, 2011; Kumara et al., 2014) (see Figure 3). Their use and survival in rainforest fragments are influenced by resource quality and certain habitat characteristics such as basal area, tree density, height of the tree, and so on (Umaphathy et al., 2011). Moreover, studies have also revealed that LTM populations in fragmented habitats have large group size, high male to female and female to immature ratios due to lack of dispersal spaces, depleted genetic diversity, and reduced home ranges as compared with groups that inhabit contiguous forests, making this species vulnerable in islands of forests surrounded by plantations (Ram et al., 2015; Singh, Kumara, Kumar, & Sharma, 2001; Umaphathy et al., 2011; Ram et al., 2015). Unlike other species of the same genus—the Bonnet macaque (*Macaca radiata*) and Rhesus macaque (*Macaca mulatta*)—the LTMs are less adaptable and are vulnerable to extensive changes in their habitats that often threaten their survival. Diversion of rainforests for coffee, rubber, and tea plantations surrounding LTM habitats alters their ranging and




Figure 3. Adult lion-tailed macaque in a rainforest fragment in the Anamalai Hills, southern India. Image credits: Ganesh Raghunathan.



Figure 2. Leopard in a fuel station amidst plantation-habitat mosaic in the Anamalai Hills, southern India. Image credits: Ganesh Raghunathan.



Figure 4. Elephants in a tea plantation in the Anamalai Hills, southern India. Image credits: Sreedhar Vijayakrishnan.

availability of food resources while influencing their behavioral responses (Singh et al., 2001; Erinjery, Kavana, & Singh, 2015), leading to negative interactions with people in some areas.

Often, these human–wildlife interactions are negative in nature leading to loss of life on both sides besides property and livestock damages. This has forced state forest departments to adopt extreme reactive measures such as capture and removal, or translocation, as a way to resolve or mitigate human–wildlife conflict. Such reactive measures neither resolve conflicts nor benefit wildlife conservation in modified landscapes (Athreya, Odden, Linnell, & Karanth, 2011; Fernando, Leimgruber, Prasad, & Pastorini, 2012). In 2012, the Karnataka Elephant Task Force constituted by Karnataka State High Court recommended importance of zonation of habitats or habitations for elephants and people by creating elephant conservation zone, coexistence zone, and elephant removal zone as a way to promote elephant conservation and resolve conflicts. However, such clear zone demarcation is based on extent of forest cover availability, intensity of conflicts, and intensity of land use, supporting livelihood of people in natural and modified elephant landscapes, and often, not based on understanding of elephant movement patterns and considering their ranging behavior outside PAs. From an anthropogenic perspective, such zonation may be helpful in terms of management, often in the form of short-term solutions for human–wildlife interactions. But for wide-ranging species such as elephants or leopards, presence of food resources and cover in altered landscapes may act as suitable habitats.

For example, in the Anamalai Hills, which holds second largest Asian elephant population in India, the 220 km² of Valparai plateau dominated by tea, coffee, and *Eucalyptus* plantations, interspersed with rainforest fragments, supports around 120 elephants amid 70,000 people. Elephants primarily use these plantations to move across the remnant fragments and the surrounding PAs (see Figure 4). Rainforest fragments, *Eucalyptus* plantations with secondary vegetation, and coffee habitats with abundant grass make the Valparai region an important resource habitat and act as refuges for elephants, while habitats such as tea facilitate movement for elephants at night. While studies show that the proportion of time the elephants spend in human-use areas is fairly high, their physiological parameters, measured in terms of fecal glucocorticoid metabolite concentrations, seem to show signs of adaptation to changing environments (Vijaykrishnan, Kumar, Umaphathy, Kumar, & Sinha, 2018).

Our research in coffee-paddy-dominated areas of Hassan region in south India shows that monoculture refuges of *Acacia* and *Eucalyptus*, coffee plantations, and remnant forest habitats provide critical habitats for elephants. Capture and removal of elephants as a

measure to mitigate human–elephant conflict in this region have not helped resolve conflicts but resulted in recolonization of the area by elephant herds, and conflicts continue to exist. Similarly, coffee-dominated Coorg region in the state of Karnataka has been intensively used by elephants throughout the year as estates are large in size with native canopy cover and lush green grass availability due to presence of agriculture, ponds or tanks, and presence of *Acacia* and *Eucalyptus* plantations enabling growth of secondary vegetation, thereby acting as feeding and sheltering habitats for elephants (Bal, Nath, Nanaya, Kushalappa, & Garcia, 2011). The presence of crops such as paddy and jackfruit amid coffee habitats attracts elephants, leading to high incidence of conflicts (Bal et al., 2011).

While generalists such as elephants and leopards adapt fairly well to changing environments, the persistence of the LTM, a habitat specialist species, would depend on retention or establishment of canopy connectivity that would connect forest fragments to facilitate movement of males and improve genetic diversity in fragmented populations (Kumar, Singh, Kumara, Sharma, & Bertsch, 2001; Umaphathy et al., 2011; Ram et al., 2015). The larger question of inbreeding is yet to be explored in large mammals in fragmented landscapes but facilitating free passage by establishing physical and functional connectivity using tree cover areas of monoculture habitats to nearest forest habitats would enhance their survival in production landscapes.

This article intends to bring out the point that wide-ranging species such as elephants and leopards, which are habitat generalists, can adapt and thrive fairly well in landscape matrices of anthropogenic habitats and in changing environments. While in the case of habitat sensitive and specialist species such as LTMs, further degradation or disappearance of rainforest habitats could potentially lead to local extinction of species.

A major challenge for wildlife conservation outside PAs is that most people are unaware of substantial biodiversity that exists in their own lands. The necessity for increased stewardship of PAs and communicating the beauty and ecological importance of wildlife in production landscapes to combat the thinking of restricting wild animals to forests alone is important. As biologists, we need to bring science and conservation efforts to the general public by getting people involved in conservation by understanding the role of nonforest areas for wildlife conservation. This transparent and integrative approach could help develop an appreciation of the criticality of natural and nonnatural habitats in biodiversity conservation. Through combined efforts among researchers, conservation agencies, policy makers, and citizens, it is not a distant possibility to bring meaningful changes in the idea or perception of what constitutes a habitat and how they are viewed, managed, and appreciated.

Acknowledgments

The authors wish to thank Tamil Nadu and Karnataka state forest departments, plantation companies, and local communities for their continued support and Nature Conservation Foundation, National Institute of Advanced Studies, and University of Mysore for institutional support. The authors also thank Ganesh Raghunathan for letting them use his images for the paper. Whitley Fund for Nature, Elephant Family, Rufford Foundation, Van Tienhoven Foundation, Nilekani Philanthropies and Oracle India have supported the long-term research.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

- Athreya, V., Odden, M., Linnell, J. D., & Karanth, K. U. (2011). Translocation as a tool for mitigating conflict with leopards in human-dominated landscapes of India. *Conservation Biology*, 25(1): 133–141.
- Athreya, V., Odden, M., Linnell, J. D., Krishnaswamy, J., & Karanth, U. (2013). Big cats in our backyards: Persistence of large carnivores in a human dominated landscape in India. *PLoS One*, 8(3): 2–9.
- Bal, P., Nath, C. D., Nanaya, K. M., Kushalappa, C. G., & Garcia, C. (2011). Elephants also like coffee: Trends and drivers of human–elephant conflicts in coffee agroforestry landscapes of Kodagu, Western Ghats, India. *Environmental Management*, 47(5): 789–801.
- Chang, C. H., Karanth, K. K., & Robbins, P. (2018). Birds and beans: Comparing avian richness and endemism in arabica and robusta agroforests in India's Western Ghats. *Scientific Reports*, 8, 3143.
- Erinjery, J. J., Kavana, T. S., & Singh, M. (2015). Food resources, distribution and seasonal variations in ranging in lion-tailed macaques, *Macaca silenus* in the Western Ghats, India. *Primates*, 56(1): 45–54.
- Fernando, P., Leimgruber, P., Prasad, T., & Pastorini, J. (2012). Problem–elephant translocation: Translocating the problem and the elephant? *PLoS One*, 7(12): 1–9.
- Graham, M. D., Douglas-Hamilton, I., Adams, W. M., & Lee, P. C. (2009). The movement of African elephants in a human-dominated land-use mosaic. *Animal Conservation*, 12(5): 445–455.
- Jha, C. S., Dutt, C. B. S., & Bawa, K. S. (2000). Deforestation and land use changes in Western Ghats, India. *Current Science*, 79, 239–238.
- Karnataka Elephant Task Force. (2012). *Report of the Karnataka Elephant Task Force, Submitted to the High Court of Karnataka, Bangalore, India*. Retrieved from <http://www.indiaenvironmentportal.org.in/files/file/Report%20of%20the%20Karnataka%20Elephant%20Task%20Force.pdf>.
- Kshetry, A., Vaidyanathan, S., & Athreya, V. (2017). Leopard in a tea-cup: A study of leopard habitat-use and human-leopard interactions in north-eastern India. *PLoS One*, 12(5): e0177013.
- Kumar, M. A., Mudappa, D., & Shankar Raman, T. R. (2010). Asian elephant *Elephas maximus* habitat use and ranging in fragmented rainforest and plantations in the Anamalai Hills, India. *Tropical Conservation Science*, 33(322): 143–158.
- Kumar, M. A., Singh, M., Kumara, H. N., Sharma, A. K., & Bertsch, C. (2001). Male migration in lion-tailed macaques. *Primate Report*, 59, 5–17.
- Kumara, H. N., Sasi, R., Suganthasakthivel, R., Singh, M., Sushma, H. S., Ramachandran, K. K., & Kaumanns, W. (2014). Distribution, demography, and conservation of lion-tailed macaques (*Macaca silenus*) in the Anamalai Hills Landscape, Western Ghats, India. *International Journal of Primatology*, 35(5): 976–989.
- Madhusudan, M. D., Sharma, N., Raghunath, R., Baskaran, N., Bipin, C. M., Gubbi, S., . . . Pillay, R. (2015). Distribution, relative abundance, and conservation status of Asian elephants in Karnataka, southern India. *Biological Conservation*, 187, 34–40.
- Mudappa, D., Kumar, M. A., & Raman, T. R. S. (2014). Wildlife conservation in landscapes fragmented by plantation crops in India. In: M. Rangarajan, M. D. Madhusudan, & G. Shahabuddin (Eds.). *Nature Without Borders*, (pp. 178–214). Hyderabad, India: Orient Blackswan Pvt Ltd.
- Navya, R., Athreya, V., Mudappa, D., & Raman, T. S. (2014). Assessing leopard occurrence in the plantation landscape of Valparai, Anamalai Hills. *Current Science*, 107(9): 1381–1385.
- Ram, M. S., Marne, M., Gaur, A., Kumara, H. N., Singh, M., Kumar, A., & Umopathy, G. (2015). Pre-historic and recent vicariance events shape genetic structure and diversity in endangered lion-tailed macaque in the Western Ghats: Implications for conservation. *PLoS One*, 10(11): e0142597.
- Santiapillai, C., & Jackson, P. (1990). *The Asian elephant: An action plan for its conservation*. Gland, Switzerland: IUCN.
- Sidhu, S., Raman, T. S., & Mudappa, D. (2015). Prey abundance and leopard diet in a plantation and rainforest landscape, Anamalai Hills, Western Ghats. *Current Science*, 109(2): 323.
- Singh, M., Kaumanns, W., Sushma, H. S., & Molur, S. (2009). The lion-tailed macaque *Macaca silenus* (Primates: Cercopithecidae): Conservation history and status of a flagship species of the tropical rainforests of the Western Ghats, India. *Journal of Threatened Taxa*, 1(3): 151–157.
- Singh, M., Kumara, H. N., Kumar, M. A., & Sharma, A. K. (2001). Behavioural responses of lion-tailed macaques (*Macaca silenus*) to a changing habitat in a tropical rain forest fragment in the Western Ghats, India. *Folia Primatologica*, 72(5): 278–291. Retrieved from doi.org/10.1002/9781119179313.wbprim0458.
- Sinha, A., & Vijayakrishnan, S. (2017). Primates in urban settings. *The International Encyclopedia of Primatology*. Retrieved from <https://doi.org/10.1002/9781119179313.wbprim0458>.
- Sridhar, H., Raman, T. S., & Mudappa, D. (2008). Mammal persistence and abundance in tropical rainforest remnants in the southern Western Ghats, India. *Current Science*, 94(6): 748–757.
- Srinivasaiah, N. M., Anand, V. D., Vaidyanathan, S., & Sinha, A. (2012). Usual populations, unusual individuals: Insights into the behavior and management of Asian elephants in fragmented landscapes. *PLoS One*, 7(8): e42571.
- Umopathy, G., Hussain, S., & Shivaji, S. (2011). Impact of habitat fragmentation on the demography of lion-tailed macaque (*Macaca silenus*) populations in the rainforests of Anamalai Hills, Western Ghats, India. *International Journal of Primatology*, 32(4): 889–900.
- Vijayakrishnan, S., Kumar, M. A., Umopathy, G., Kumar, V., & Sinha, A. (2018). Physiological stress responses in wild Asian elephants *Elephas maximus* in a human-dominated landscape in the Western Ghats, southern India. *General and Comparative Endocrinology*. Retrieved from <https://doi.org/10.1016/j.ygcen.2018.05.009>.