

Monetary Diet Cost is Associated with not only Favorable but also Unfavorable Aspects of Diet in Pregnant Japanese Women: The Osaka Maternal and Child Health Study

Authors: Murakami, Kentaro, Miyake, Yoshihiro, Sasaki, Satoshi, Tanaka, Keiko, Ohya, Yukihiro, et al.

Source: Environmental Health Insights, 3(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/EHI.S2508>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

OPEN ACCESS

Full open access to this and thousands of other papers at <http://www.la-press.com>.

Monetary Diet Cost is Associated with not only Favorable but also Unfavorable Aspects of Diet in Pregnant Japanese Women: The Osaka Maternal and Child Health Study

Kentaro Murakami¹, Yoshihiro Miyake², Satoshi Sasaki¹, Keiko Tanaka², Yukihiro Ohya³, Yoshio Hirota⁴, and Osaka Maternal and Child Health Study Group⁵

¹Department of Social and Preventive Epidemiology, School of Public Health, the University of Tokyo, Tokyo, Japan.

²Department of Public Health, Faculty of Medicine, Fukuoka University, Fukuoka, Japan. ³Division of Allergy, Department of Medical Specialties, National Center for Child Health and Development, Tokyo, Japan. ⁴Department of Public Health, Osaka City University School of Medicine, Osaka, Japan. ⁵Other members of the Osaka Maternal and Child Health Study Group are listed in the Acknowledgments. Email: kenmrkm@m.u-tokyo.ac.jp

Abstract: While several observational studies in European countries have shown that higher monetary diet cost is associated with healthier diets, information on the relationship of cost to diet quality in other countries is sparse, including Japan. This cross-sectional study examined the association between monetary diet cost and dietary intake in a group of pregnant Japanese women. Subjects were 596 pregnant Japanese housewives. Dietary intake was estimated using a validated, self-administered, comprehensive diet history questionnaire. Monetary diet cost was calculated using retail food prices. Values of monetary diet cost and nutrient and food intake were energy-adjusted using the density method. Monetary diet cost was associated positively with the intake of protein, total fat, saturated fatty acids, dietary fiber, cholesterol, sodium, potassium, calcium, magnesium, iron, vitamins A, D, E, C, and folate, and inversely with that of carbohydrate. For foods, cost was associated positively with the intake of potatoes, pulses and nuts, fish and shellfish, meat, dairy products, vegetables, and fruits, and inversely with that of rice and bread. No association was seen for noodles, confectioneries and sugars, fats and oils, or eggs. Cost was also associated inversely with dietary energy density. In conclusion, monetary diet cost was associated with not only favorable aspects of diet, including a higher intake of dietary fiber, key vitamins and minerals, fruits, and vegetables and lower dietary energy density, but also unfavorable aspects, including a higher intake of fat and sodium and lower intake of carbohydrate and rice, in a group of pregnant Japanese women.

Keywords: diet cost, nutrient intake, food intake, pregnant women, Japan

Environmental Health Insights 2009:3 27–35

This article is available from <http://www.la-press.com>.

© the authors, licensee Libertas Academica Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://www.creativecommons.org/licenses/by/2.0>) which permits unrestricted use, distribution and reproduction provided the original work is properly cited.



Introduction

While food choice is influenced by a large number of factors,¹ the price of food is an important determinant.^{2,3} Generally, energy-dense and nutrient-dilute foods such as cereals, fats and oils, and sugar and confectioneries provide dietary energy at lowest cost.⁴⁻⁹ Conversely, the cost per calorie of energy-dilute and nutrient-dense foods, including fish and shellfish, vegetables, and fruits is much higher.⁴⁻⁹ If healthier foods cost more then so too will healthier diets, and several observational studies conducted in European countries have in fact shown that healthful diets are more expensive than less healthful diets,⁴⁻⁹ albeit that the currently available procedure for estimating monetary diet cost is based on the assumption that all foods are purchased and then prepared and consumed at home, and hence cannot consider the monetary cost of eating out.¹⁰ Given the important role of nutrition in promoting health,^{11,12} the positive association between monetary diet cost and diet quality may explain, at least in part, the socioeconomic gradient in health widely observed in Western countries, with higher socioeconomic position associated with better health.¹³⁻¹⁵

However, information on the relationship of monetary diet cost to diet quality in other countries is sparse, including Japan. Considering the unclear or even inverse association between socioeconomic position and health status observed in several Japanese populations,¹⁶⁻¹⁸ the association between monetary diet cost and diet quality may differ between European countries and Japan. In a study of Japanese female dietetic students, for example, monetary diet cost was not necessarily associated with healthier dietary intake patterns.¹⁹ This finding may have been confounded by a lack of information on whether the participants bought their food themselves, however, given that the majority (89%) lived with their family.¹⁹ Because the relationship between monetary diet cost and diet quality is an important public health issue, further investigation in Japanese people is clearly needed. Additionally, the association between monetary diet cost and dietary intake has not been investigated among women during pregnancy, a nutritionally important lifestyle stage, even in Western countries.

The present cross-sectional study examined the association between monetary diet cost and dietary

intake in a group of pregnant Japanese women using baseline data from the Osaka Maternal and Child Health Study (OMCHS). We selected only housewives who did not habitually eat out and who lived in nuclear families in non-agricultural societies for accurate estimation of monetary diet cost. We hypothesized that monetary diet cost is associated with both favorable and unfavorable aspects of diet.

Methods

Subjects

The subjects in the present analysis were participants in the baseline survey of the OMCHS, an ongoing prospective cohort study, the main purpose of which was to investigate preventive and risk factors for maternal and child health problems. Details of the OMCHS have been published elsewhere.^{20,21} Briefly, all pregnant women in Neyagawa City, Osaka Prefecture (metropolis in Japan with a total population of approximately 8.8 million), were recruited between November 2001 and March 2003. Of 3639 eligible women, 627 (17.2%) took part in the study. An additional 375 pregnant women living in other neighborhood areas were enrolled in the study between December 2001 and November 2003, giving a total of 1002 pregnant women completing the baseline survey. Written informed consent was obtained from each woman. The protocol of the OMCHS was approved by the ethics committees of Osaka City University School of Medicine, Japan.

For the present analysis, we excluded women living in an expanded family ($n = 127$), those working outside the home ($n = 288$), and those who dined out 16 times or more during the preceding month (i.e. four times or more per week) ($n = 118$). Because some women fell into more than one exclusion category, the final analysis sample comprised 596 women.

Measurements

Baseline assessment of the OMCHS included a set of two self-administered questionnaires. The participants mailed the answered questionnaires to the data management center. Research technicians completed missing or illogical data by telephone interview.

One of the self-administered questionnaires was a validated, self-administered, comprehensive diet history questionnaire (DHQ) that assesses dietary habits during the preceding month. Details of the



DHQ's structure, calculation of dietary intake, and validity for commonly studied nutritional factors have been published elsewhere.^{22–25} Briefly, the DHQ is a 16-page structured questionnaire which asks about the consumption frequency and portion size of selected foods commonly consumed in Japan as well as general dietary behavior and usual cooking methods.²² Estimates of daily intake for foods (150 items in total), nutrients, and energy were calculated using an ad hoc computer algorithm for the DHQ,^{22,25} based on the Standard Tables of Food Composition in Japan.²⁶

Monetary diet cost (Japanese yen/day) was calculated by multiplying the amount of each food estimated from the DHQ (g/day) by the estimated price of the food (Japanese yen/g) and summing the products (1 Japanese yen = 0.0077 pound sterling = 0.0085 euros = 0.0110 US dollars in February 2009). Detailed descriptions of the cost calculation method as well as the monetary cost of each food have been published elsewhere.^{19,27} Briefly, the price of each food was determined based on the National Retail Price Survey 2004.²⁸ For foods whose prices were not published in the survey (13 items), prices were taken from the websites of a nationally distributed supermarket (Seiyu, Tokyo, Japan) or fast-food restaurant (McDonalds, Tokyo, Japan and Mister Donut, Tokyo, Japan) chains. Alcoholic beverages (six items), non-caloric beverages (four items), and water (three items) were excluded from calculation.⁹ Costs of combined foods such as pizza were calculated using the prices of frozen equivalents.⁸ Costs were estimated based on the assumption that all foods were purchased and then prepared and consumed at home.¹⁰

To minimize the influence of dietary misreporting, an ongoing controversy in studies that collect dietary information using self-report instruments,²⁹ values of monetary diet cost as well as intakes of nutrients and foods were energy-adjusted using the density method (i.e. monetary diet cost per 1000 kcal of energy, percentage of energy for energy-providing nutrients, and amount per 1000 kcal of energy for other nutrients and foods). Dietary energy density (kcal/g) was calculated based on foods only (128 items, excluding 22 beverages), as described elsewhere.³⁰ In a previous study of 92 Japanese women, Pearson correlation coefficient between the DHQ and 16-day weighed

dietary records were 0.57 for monetary diet cost, suggesting satisfactory validity of the DHQ in terms of monetary diet cost.²⁷

The other self-administered questionnaire elicited information on potential confounding factors, including age (28 years or less, 29–31 years, and 32 years or more), gestational age (14 weeks or less, 15–20 weeks, and 21 weeks or more), parity (nulliparous and multiparous), cigarette smoking (never, former, and current), changes in diet in the previous one month (none or few, slight, and substantial), physical activity (low; and medium or high), partner's education [low (12 years or less), medium (13–14 years), and high (15 years or more)], education [low (12 years or less), medium (13–14 years), and high (15 years or more)], and household income as calculated considering differences in household size and composition³¹ [divided into approximate tertiles: low (2,499,999 yen/year or less), medium (2,500,000–3,099,999 yen/year), and high (3,100,000 yen/year or more)].

Statistical analysis

All statistical analyses were performed using SAS statistical software version 9.1 (SAS Institute Inc., Cary, NC, USA). Using the PROC GLM procedure, linear regression models were constructed to examine the association of monetary diet cost with dietary intake of selected nutrients and foods. For analyses, subjects were categorized into quartiles according to monetary diet cost. Mean (and standard error) values of nutrient and food intake were calculated by quartile of monetary diet cost with adjustment for potential confounding factors, including age, gestational age, parity, cigarette smoking, changes in diet in the previous month, physical activity, partner's education, education, and household income. Tests for trends were performed by modeling the median value of each quartile category as a continuous variable. All reported *P*-values are two-tailed, and a *P*-value of <0.05 was considered statistically significant.

Results

Dietary characteristics of the 596 women are shown in Table 1. Mean monetary diet cost was 458.9 Japanese yen/1000 kcal, while mean percentages of intake of protein, total fat, and carbohydrate were 13.4%,

**Table 1.** Dietary characteristics of the subjects (n = 596).^a

Variable	
Monetary diet cost (Japanese yen/1000 kcal) ^b	458.9 ± 72.7
Nutrient intake	
Protein (% of energy)	13.4 ± 2.0
Total fat (% of energy)	30.0 ± 5.5
Saturated fatty acids (% of energy)	8.5 ± 2.0
Carbohydrate (% of energy)	55.6 ± 6.3
Dietary fiber (g/1000 kcal)	6.5 ± 1.8
Cholesterol (mg/1000 kcal)	164.5 ± 59.5
Sodium (mg/1000 kcal)	2039 ± 437
Potassium (mg/1000 kcal)	1151 ± 270
Calcium (mg/1000 kcal)	294.5 ± 97.8
Magnesium (mg/1000 kcal)	121.3 ± 26.2
Iron (mg/1000 kcal)	3.66 ± 0.84
Vitamin A (μg retinol equivalents/1000 kcal)	341.3 ± 317.8
Vitamin D (mg/1000 kcal)	3.34 ± 1.78
Vitamin E (mg alpha-tocopherol/1000 kcal)	4.25 ± 0.91
Vitamin C (mg/1000 kcal)	58.6 ± 28.5
Folate (μg/1000 kcal)	157.4 ± 49.2
Food intake (g/1000 kcal)	
Rice	133.9 ± 56.6
Bread	38.8 ± 22.5
Noodles	43.5 ± 32.4
Potatoes	15.2 ± 11.4
Confectioneries and sugars	33.8 ± 17.3
Fats and oils	13.6 ± 6.2
Pulses and nuts	26.2 ± 16.9
Fish and shellfish	26.1 ± 13.5
Meat	32.5 ± 15.2
Eggs	17.9 ± 13.0
Dairy products	103.8 ± 67.5
Vegetables	115.5 ± 67.4
Fruits	94.8 ± 92.7
Dietary energy density (kcal/g)	1.38 ± 0.20

^aValues are mean ± standard error.

^b1 Japanese yen = 0.0077 pound sterling = 0.0085 euros = 0.0110 U.S. dollars in February 2009.

30.0%, and 55.6% of energy, respectively. Selected characteristics by quartile of monetary diet cost are shown in Table 2. Women in the higher quartiles of cost tended to be never smokers and have changes in diet in the previous month, a partner with higher education, higher education, and higher household income.

Dietary characteristics by quartile of monetary diet cost are shown in Table 3. At the nutrient level, monetary diet cost was associated positively with the intake of protein, total fat, saturated fatty acids, dietary fiber, cholesterol, sodium, potassium, calcium, magnesium, iron, vitamins A, D, E, and C, and folate, and inversely with that of carbohydrate. At the food level, monetary diet cost was associated positively with the intake of potatoes, pulses and nuts, fish and shellfish, meat, dairy products, vegetables, and fruits, and inversely with that of rice and bread. No association was seen for noodles, confectioneries and sugars, fats and oils, or eggs. Further, monetary diet cost was inversely associated with dietary energy density.

Discussion

Consistent with a previous study of Japanese female dietetic students,¹⁹ we found that monetary diet cost was associated with not only favorable aspects of diet such as a higher intake of dietary fiber, key vitamins and minerals, and fruits and vegetables and lower dietary energy density, but also unfavorable aspects of diet such as a higher intake of fat and sodium and lower intake of carbohydrate and rice in a group of pregnant Japanese women. These findings somewhat conflict with those of previous Western studies, in which monetary diet cost has been consistently associated with healthier diets, including higher intake of key micronutrients, fruits, and vegetables, lower intake of fats and sweets, and lower dietary energy density.⁴⁻⁹

For Japanese people, rice is a staple food which is consumed at almost every meal, accompanied by a main and several side dishes consisting mainly of fish and shellfish, meat, egg, vegetables, and pulses. Rice is relatively inexpensive compared with the component foods of such main and side dishes in Japan.¹⁹ We previously speculated that persons with a limited budget for foods would mainly consume rice, with a relatively restricted amount or variety of

**Table 2.** Selected characteristics according to quartile category of monetary diet cost (n = 596).^a

Variable	Quartile 1 (n = 149)	Quartile 2 (n = 149)	Quartile 3 (n = 149)	Quartile 4 (n = 149)	P ^b
Monetary diet cost (Japanese yen/1000 kcal) ^c					
Median	379.6	434.1	475.3	536.6	
Range	223.6–412.8	412.9–453.5	453.6–501.4	501.5–829.2	
Age					0.18
28 years or less	72 (48.3)	54 (36.2)	51 (34.2)	57 (38.3)	
29–31 years	33 (22.2)	53 (35.6)	43 (28.9)	48 (32.2)	
32 years or more	44 (29.5)	42 (28.2)	55 (36.9)	44 (29.5)	
Gestational age					0.41
14 weeks or less	50 (33.6)	58 (38.9)	51 (34.2)	53 (35.6)	
15–20 weeks	45 (30.2)	45 (30.2)	57 (38.3)	52 (34.9)	
21 weeks or more	54 (36.2)	46 (30.9)	41 (27.5)	44 (29.5)	
Parity					0.029
Nulliparous	54 (36.2)	70 (47.0)	57 (38.3)	78 (52.4)	
Multiparous	95 (63.8)	79 (53.0)	92 (61.7)	71 (47.7)	
Cigarette smoking					0.021
Never	88 (59.1)	111 (74.5)	101 (67.8)	106 (71.1)	
Former	20 (13.4)	19 (12.8)	28 (18.8)	20 (13.4)	
Current	41 (27.5)	19 (12.8)	20 (13.4)	23 (15.4)	
Changes in diet in the previous one month					0.0007
None or few	56 (37.6)	50 (33.6)	35 (23.5)	35 (23.5)	
Slight	61 (40.9)	66 (44.3)	76 (51.0)	64 (43.0)	
Substantial	32 (21.5)	33 (22.2)	38 (25.5)	50 (33.6)	
Physical activity					0.97
Low	101 (67.8)	81 (54.4)	91 (61.1)	98 (65.8)	
Medium or high	48 (32.2)	68 (45.6)	58 (38.9)	51 (34.2)	
Partner's education					0.0020
Low (12 years or less)	69 (46.3)	68 (45.6)	58 (38.9)	50 (33.6)	
Medium (13–14 years)	30 (20.1)	25 (16.8)	20 (13.4)	25 (16.8)	
High (15 years or more)	50 (33.6)	56 (37.6)	71 (47.7)	74 (49.7)	
Education					<0.0001
Low (12 years or less)	64 (43.0)	61 (40.9)	46 (30.9)	38 (25.5)	
Medium (13–14 years)	58 (38.9)	60 (40.3)	61 (40.9)	63 (42.3)	
High (15 years or more)	27 (18.1)	28 (18.8)	42 (28.2)	48 (32.2)	
Household income					<0.0001
Low (2,499,999 yen/year or less)	76 (51.0)	55 (36.9)	45 (30.2)	45 (30.2)	
Medium (2,500,000–3,099,999 yen/year)	50 (33.6)	54 (36.2)	53 (35.6)	45 (30.2)	
High (3,100,000 yen/year or more)	23 (15.4)	40 (26.9)	51 (34.2)	59 (39.6)	

^aValues are number of subjects (%) unless otherwise indicated.^bMantel-Haenszel chi-square test.^c1 Japanese yen = 0.0077 pound sterling = 0.0085 euros = 0.0110 US dollars in February 2009.

Table 3. Dietary characteristics according to quartile category of monetary diet cost (n = 596).^a

Variable	Quartile 1 (n = 149)	Quartile 2 (n = 149)	Quartile 3 (n = 149)	Quartile 4 (n = 149)	P for trend ^b
Nutrient intake					
Protein (% of energy)	12.1 ± 0.1	13.2 ± 0.1	13.7 ± 0.1	14.6 ± 0.1	<0.0001
Total fat (% of energy)	27.8 ± 0.5	30.2 ± 0.4	30.8 ± 0.4	31.2 ± 0.4	<0.0001
Saturated fatty acids (% of energy)	7.5 ± 0.2	8.5 ± 0.2	8.9 ± 0.2	8.9 ± 0.2	<0.0001
Carbohydrate (% of energy)	58.5 ± 0.5	55.5 ± 0.5	54.6 ± 0.5	54.0 ± 0.5	<0.0001
Dietary fiber (g/1000 kcal)	5.6 ± 0.1	6.0 ± 0.1	6.5 ± 0.1	7.7 ± 0.1	<0.0001
Cholesterol (mg/1000 kcal)	149.1 ± 4.9	167.9 ± 4.9	173.7 ± 4.8	167.2 ± 4.9	0.012
Sodium (mg/1000 kcal)	1890 ± 36	1997 ± 36	2120 ± 36	2150 ± 36	<0.0001
Potassium (mg/1000 kcal)	937 ± 17	1087 ± 17	1191 ± 18	1391 ± 18	<0.0001
Calcium (mg/1000 kcal)	245.3 ± 7.2	286.0 ± 7.1	308.6 ± 7.1	338.2 ± 7.1	<0.0001
Magnesium (mg/1000 kcal)	106.8 ± 1.9	115.9 ± 1.9	123.4 ± 1.9	139.1 ± 1.9	<0.0001
Iron (mg/1000 kcal)	3.22 ± 0.06	3.50 ± 0.06	3.68 ± 0.06	4.21 ± 0.06	<0.0001
Vitamin A (μg retinol equivalents/1000 kcal)	230.9 ± 25.4	313.1 ± 25.0	343.3 ± 24.9	477.9 ± 25.0	<0.0001
Vitamin D (mg/1000 kcal)	2.59 ± 0.14	3.08 ± 0.14	3.52 ± 0.14	4.18 ± 0.14	<0.0001
Vitamin E (mg alpha-tocopherol/1000 kcal)	3.69 ± 0.07	4.09 ± 0.07	4.36 ± 0.07	4.85 ± 0.07	<0.0001
Vitamin C (mg/1000 kcal)	43.0 ± 2.1	50.4 ± 2.1	60.1 ± 2.0	80.8 ± 2.1	<0.0001
Folate (μg/1000 kcal)	133.0 ± 3.5	144.7 ± 3.5	157.4 ± 3.5	194.6 ± 3.5	<0.0001
Food intake (g/1000 kcal)					
Rice	169.1 ± 4.4	136.1 ± 4.3	123.7 ± 4.3	106.5 ± 4.3	<0.0001
Bread	44.7 ± 1.9	40.9 ± 1.9	36.3 ± 1.8	33.3 ± 1.9	<0.0001
Noodles	45.1 ± 2.7	48.1 ± 2.7	40.8 ± 2.7	39.9 ± 2.7	0.07
Potatoes	12.9 ± 0.9	14.5 ± 0.9	15.8 ± 0.9	17.8 ± 0.9	0.0001
Confectioneries and sugars	33.8 ± 1.5	33.7 ± 1.4	34.8 ± 1.4	32.6 ± 1.4	0.65
Fats and oils	13.8 ± 0.5	13.9 ± 0.5	13.3 ± 0.5	13.2 ± 0.5	0.30
Pulses and nuts	21.8 ± 1.4	23.0 ± 1.3	27.6 ± 1.3	32.3 ± 1.3	<0.0001
Fish and shellfish	18.5 ± 1.0	23.7 ± 1.0	29.3 ± 1.0	33.1 ± 1.0	<0.0001
Meat	24.8 ± 1.2	32.7 ± 1.2	33.1 ± 1.2	39.3 ± 1.2	<0.0001
Eggs	17.6 ± 1.1	19.3 ± 1.1	19.1 ± 1.1	15.7 ± 1.1	0.18
Dairy products	84.7 ± 5.5	109.3 ± 5.4	109.8 ± 5.4	111.3 ± 5.4	0.0016
Vegetables	73.8 ± 4.9	103.3 ± 4.8	119.6 ± 4.8	165.1 ± 4.8	<0.0001
Fruits	60.1 ± 7.4	78.0 ± 7.3	104.2 ± 7.2	136.9 ± 7.3	<0.0001
Dietary energy density (kcal/g)	1.48 ± 0.02	1.41 ± 0.02	1.36 ± 0.01	1.27 ± 0.02	<0.0001

^aValues are means ± standard error. Adjustment was made for age (28 years or less, 29–31 years, and 32 years or more), gestational age (14 weeks or less, 15–20 weeks, and 21 weeks or more), parity (nulliparous and multiparous), cigarette smoking (never, former, and current), changes in diet previous one month (none or few, slight, and substantial), physical activity (low; and medium or high), and partner's education [low (12 years or less), medium (13–14 years), and high (15 years or more)], education [low (12 years or less), medium (13–14 years), and high (15 years or more)], and household income [low (2,419,999 yen/year or less), medium (2,420,000–3,619,999 yen/year), and high (3,620,000 yen/year or more)].

^bTest for linear trend in general linear regression.



main and side dishes, while those with a larger food budget would have an increased amount or variety of main and side dishes and decreased consumption of rice.¹⁹ This hypothesis may be supported by the decreasing consumption of rice and increasing consumption of other foods such as meat, vegetables, fish and shellfish, and pulses³² seen with the increase in the Gross National Product of Japan³³ from 1955 to 2000.¹⁹

With regard to the unfavorable aspects of diet at the nutrient level, the higher intake of total fat, saturated fatty acids, and cholesterol and lower intake of carbohydrate seen with increasing monetary diet cost appeared to be due to lower rice consumption (and dietary intake patterns associated with lower rice intake) with increasing monetary diet cost; in apparent support of this, rice intake was in fact negatively associated with the intake of total fat (Pearson correlation coefficient: -0.52), saturated fatty acids (-0.48), and cholesterol (-0.19) and positively with carbohydrate intake (0.44). The higher intake of sodium seen with increasing monetary diet cost may be due to higher intake of vegetables, meat, fish and shellfish, and pulses and nuts, because in Japan these foods are usually accompanied by seasonings with a salty taste, such as salt, soy sauce, miso (fermented soybean paste), and dressings. Intake of these foods was in fact positively associated with sodium intake in the present study (Pearson correlation coefficients: 0.33 , 0.12 , 0.30 , and 0.35 , respectively).

Several limitations of the study warrant mention. First, the study subjects were selected pregnant Japanese housewives. Additionally, response rate for the 627 women living in Neyagawa City was only 17.2%, and could not be calculated for the remaining 375 women living in other areas because the exact number of eligible subjects among the sources from which they recruited was not available. Thus, the present results could not be extrapolated to other populations.

Although the validity of the DHQ appears reasonable with regard to commonly studied nutritional factors,^{22–25} it was not specifically designed to measure monetary diet cost. This limitation is common to other studies on this topic.^{4–9,19} Additionally, food prices were derived from the National Retail Price Survey and websites of nationally distributed supermarket

and fast food restaurant chains. As this procedure gives only an approximation of actual diet costs, the results of the present study should be interpreted with caution, although a similar methodology has been used in all previous observational studies.^{4–9,19} Nevertheless, the satisfactory validity of this DHQ regarding monetary diet cost against 16-day weighed dietary records as described above may provide some reassurance.²⁷

In conclusion, monetary diet cost was associated with not only favorable but unfavorable aspects of diet in a group of pregnant Japanese women. Because the relation between monetary diet cost and diet quality is an important public health topic, further research is needed to investigate whether the association found in this specific group of subjects would also be observed in other Japanese populations.

Acknowledgments

We thank the Neyagawa City Government, Hirakata City Government, Katano City Government, Shijonawate City Government, Kaizuka City Government, Takaishi City Government, Hannan City Government, Neyagawa City Medical Association, Hirakata City Medical Association, and Kadoma City Medical Association for their valuable support; and Tomoko Shibazaki, Nahoko Nishimura, and Naomi Takaoka for their assistance.

Other members of the Osaka Maternal and Child Health Study Group are as follows: Shoichi Miyamoto (Department of Public Health, Osaka City University Graduate School of Medicine); Ichiro Matsunaga, Toshiaki Yoshida, and Hajime Oda (Osaka Prefectural Institute of Public Health); Hideharu Kanzaki and Mitsuyoshi Kitada (Department of Obstetrics and Gynecology, Kansai Medical University); Yorihiro Horikoshi (Department of Obstetrics and Gynecology, Kansai Medical University Kori); Osamu Ishiko, Yuichiro Nakai, Junko Nishio, and Seiichi Yamamasu (Department of Obstetrics and Gynecology, Osaka City University Graduate School of Medicine); Jinsuke Yasuda (Department of Obstetrics and Gynecology, Matsushita Memorial Hospital); Seigo Kawai (Department of Obstetrics and Gynecology, Hoshigaoka Koseinenkin Hospital); Kazumi Yanagihara (Yanagihara Clinic); Koji Wakuda (Department of Obstetrics and Gynecology, Jujimoto Hospital); Tokio Kawashima (Kyoritsu Women's Clinic); Katsuhiko



Narimoto (Ishida Hospital Obstetrics, Gynecology); Yoshihiko Iwasa (Iwasa Women's Clinic); Katsuhiko Orino (Orino Lady's Clinic); Itsuo Tsunetoh (Tsunetoh Obstetrics and Gynecology); Junichi Yoshida (Yoshida Clinic); Junichi Iito, (Iito Obstetrics and Gynecology Clinic); Takuzi Kaneko (Kaneko Sanfujinka); Takao Kamiya (Kamiya Ladies Clinic); Hiroyuki Kuribayashi (Kuribayashi Clinic); Takeshi Taniguchi (Taniguchi Hospital); Hideo Takemura (Kosaka Women's Hospital); and Yasuhiko Morimoto (Aizenbashi Hospital).

Sources of Support

This study was supported by a Grant-in-Aid (13770206, 16790351) for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology and Health and Labour Sciences Research Grants, Research on Allergic Disease and Immunology from the Ministry of Health, Labour, and Welfare.

Disclosure

The authors report no conflicts of interest.

References

1. Shepherd R. Resistance to changes in diet. *Proc Nutr Soc.* 2002;61:267–72.
2. Lennernas M, Fjellstrom C, Becker W, Giachetti I, Schmitt A, Remaut de Winter A, et al. Influences on food choice perceived to be important by nationally-representative samples of adults in the European Union. *Eur J Clin Nutr.* 1997;51:S8–15.
3. Glanz K, Basil M, Maibach E, Goldberg J, Snyder D. Why Americans eat what they do: taste, nutrition, cost, convenience, and weight control concerns as influences on food consumption. *J Am Diet Assoc.* 1998;98:1118–26.
4. Cade J, Upmeier H, Calvert C, Greenwood D. Costs of a healthy diet: analysis from the UK Women's Cohort Study. *Public Health Nutr.* 1999;2:505–12.
5. Drewnowski A, Darmon N, Briand A. Replacing fats and sweets with vegetables and fruits—a question of cost. *Am J Public Health.* 2004;94:1555–9.
6. Drewnowski A, Monsivais P, Maillot M, Darmon N. Low-energy-density diets are associated with higher diet quality and higher diet costs in French adults. *J Am Diet Assoc.* 2007;107:1028–32.
7. Andrieu E, Darmon N, Drewnowski A. Low-cost diets: more energy, fewer nutrients. *Eur J Clin Nutr.* 2006;60:434–6.
8. Schroder H, Marrugat J, Covas MI. High monetary costs of dietary patterns associated with lower body mass index: a population-based study. *Int J Obes.* 2006;30:1574–9.
9. Maillot M, Darmon N, Vieux F, Drewnowski A. Low energy density and high nutritional quality are each associated with higher diet costs in French adults. *Am J Clin Nutr.* 2007;86:690–6.
10. Drewnowski A, Darmon N. The economics of obesity: dietary energy density and energy cost. *Am J Clin Nutr.* 2005;82:265S–73S.
11. Smith GD, Brunner E. Socio-economic differentials in health: the role of nutrition. *Proc Nutr Soc.* 1997;56:75–90.
12. Darmon N, Drewnowski A. Does social class predict diet quality? *Am J Clin Nutr.* 2008;87:1107–17.
13. Winkleby MA, Jatulis DE, Frank E, Fortmann SP. Socioeconomic status and health: how education, income, and occupation contribute to risk factors for cardiovascular disease. *Am J Public Health.* 1992;82:816–20.
14. Lantz PM, House JS, Lepkowski JM, Williams DR, Mero RP, Chen J. Socioeconomic factors, health behaviors, and mortality: results from a nationally representative prospective study of U.S. adults. *JAMA.* 1998;279:1703–8.
15. Lahelma E, Martikainen P, Laaksonen M, Aittomaaki A. Pathways between socioeconomic determinants of health. *JEpidemiol Community Health.* 2004;58:327–32.
16. Martikainen P, Ishizaki M, Marmot MG, Nakagawa H, Kagamimori S. Socioeconomic differences in behavioural and biological risk factors: a comparison of a Japanese and an English cohort of employed men. *Int J Epidemiol.* 2001;30:833–8.
17. Nishi N, Makino K, Fukuda H, Tataru K. Effects of socioeconomic indicators on coronary risk factors, self-rated health and psychological well-being among urban Japanese civil servants. *Soc Sci Med.* 2004;58:1159–70.
18. Hirokawa K, Tsutsumi A, Kayaba K. Impacts of educational level and employment status on mortality for Japanese women and men: the Jichi Medical School cohort study. *Eur J Epidemiol.* 2006;21:641–51.
19. Murakami K, Sasaki S, Okubo H, Takahashi Y, Hosoi Y, Itabashi M. Monetary costs of dietary energy reported by young Japanese women: association with food and nutrient intake and body mass index. *Public Health Nutr.* 2007;10:1430–9.
20. Miyake Y, Miyamoto S, Ohya Y, Sasaki S, Matsunaga I, Yoshida T, et al. Relationship between active and passive smoking and total serum IgE levels in Japanese women: baseline data from the Osaka Maternal and Child Health Study. *Int Arch Allergy Immunol.* 2004;135:221–8.
21. Miyake Y, Sasaki S, Ohya Y, Miyamoto S, Matsunaga I, Yoshida T, et al. Soy, isoflavones, and prevalence of allergic rhinitis in Japanese women: the Osaka Maternal and Child Health Study. *J Allergy Clin Immunol.* 2005;115:1176–83.
22. Sasaki S, Yanagibori R, Amano K. Self-administered diet history questionnaire developed for health education: a relative validation of the test-version by comparison with 3-day diet record in women. *J Epidemiol.* 1998;8:203–15.
23. Sasaki S, Yanagibori R, Amano K. Validity of a self-administered diet history questionnaire for assessment of sodium and potassium: comparison with single 24-hour urinary excretion. *Jpn Circ J.* 1998;62:431–5.
24. Sasaki S, Ushio F, Amano K, Morihara M, Todoriki T, Uehara Y, et al. Serum biomarker-based validation of a self-administered diet history questionnaire for Japanese subjects. *J Nutr Sci Vitaminol.* 2000;46:285–96.
25. Murakami K, Sasaki S, Takahashi Y, Okubo H, Hirota N, Notsu A, et al. Reproducibility and relative validity of dietary glycaemic index and load assessed with a self-administered diet-history questionnaire in Japanese adults. *Br J Nutr.* 2008;99:639–48.
26. Science and Technology Agency. Standard Tables of Food Composition in Japan, Fifth Revised and Enlarged Edition. Tokyo, Japan: Printing Bureau of the Ministry of Finance; 2005 (in Japanese).
27. Murakami K, Sasaki S, Takahashi Y, Okubo H, Hirota N, Notsu A, et al. Comparability of weighed dietary records and a self-administered diet history questionnaire for estimating monetary cost of dietary energy. *Environmental Health Insights.* 2008;1:35–43. http://www.la-press.com/article.php?article_id=1016.
28. Price Statistics Office, Statistics Bureau. National Retail Price Survey in Japan. Available at: <http://www.stat.go.jp/data/kouri/index.htm>. Accessed June 26, 2008 (in Japanese).
29. Murakami K, Sasaki S, Takahashi Y, Uenishi K, Yamasaki M, Hayabuchi H, et al. Misreporting of dietary energy, protein, potassium and sodium in relation to body mass index in young Japanese women. *Eur J Clin Nutr.* 2008;62:111–8.
30. Murakami K, Sasaki S, Takahashi Y, Uenishi K; for the Japan Dietetic Students' Study for Nutrition and Biomarkers Group. Dietary energy density is associated with body mass index and waist circumference, but not with other metabolic risk factors, in free-living young Japanese women. *Nutrition.* 2007;23:798–806.



31. Laaksonen M, Rahkonen O, Martikainen P, Lahti E. Socioeconomic position and self-rated health: the contribution of childhood socioeconomic circumstances, adult socioeconomic status, and material resources. *Am J Public Health*. 2005;95:1403–9.
32. Ministry of Health, Labour, and Welfare of Japan. The National Nutrition Survey in Japan, 1955–2000. Tokyo, Japan: Ministry of Health and Welfare, 1957–2002 (in Japanese).
33. Ministry of Health, Labour, and Welfare. Statistical tables: trend of medical expenditure, Gross National Product, and Net National Income in Japan. Available at: <http://www1.mhlw.go.jp/toukei/k-iryohi/hyo1.html>. Accessed June 26, 2008 (in Japanese).

Publish with Libertas Academica and every scientist working in your field can read your article

“I would like to say that this is the most author-friendly editing process I have experienced in over 150 publications. Thank you most sincerely.”

“The communication between your staff and me has been terrific. Whenever progress is made with the manuscript, I receive notice. Quite honestly, I’ve never had such complete communication with a journal.”

“LA is different, and hopefully represents a kind of scientific publication machinery that removes the hurdles from free flow of scientific thought.”

Your paper will be:

- Available to your entire community free of charge
- Fairly and quickly peer reviewed
- Yours! You retain copyright

<http://www.la-press.com>