

Cloning and Sequence Analysis of the Amylase Gene from the Rice Pest *Scirpophaga incertulas* Walker and its Inhibitor from Wheat (Variety MP Sehore)

Authors: Sharma, Poonam, Shankar, Pallavi R., Subramaniam, Geetha, Kumar, Amit, Tandon, Aanchal, et al.

Source: International Journal of Insect Science, 1(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/IJIS.S3384>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

OPEN ACCESS

Full open access to this and thousands of other papers at <http://www.la-press.com>.

Cloning and Sequence Analysis of the Amylase Gene from the Rice Pest *Scirpophaga incertulas* Walker and its Inhibitor from Wheat (Variety MP Sehore)

Poonam Sharma, Pallavi R. Shankar, Geetha Subramaniam, Amit Kumar, Aanchal Tandon, C.G. Suresh, Meenakshi V. Rele and Lalitha Sunil Kumar

Division of Biochemical Sciences, National Chemical Laboratory, Pune 411008, India. Email: ls.kumar@ncl.res.in

Abstract: *Scirpophaga incertulas* Walker (Lepidoptera: Pyralidae), commonly known as yellow stem borer, is a predominant monophagous pest of rice, which causes 5% to 30% loss of the rice crop. We report for the first time, the cloning and sequence analysis of the amylase gene of this pest. The cloned gene translates into a protein of 487 amino acids having a predicted molecular weight of 54,955 daltons and a theoretical pI of 5.9. The 3D structure of the amylase is predicted from its amino acid sequence by homology modeling using the structure of the amylase from *Tenebrio molitor* L (Coleoptera: Tenebrionidae). We also report the purification of a dimeric α -amylase inhibitor from a local variety of wheat MP Sehore that is specific for the amylase of this pest and does not inhibit human salivary amylase or porcine pancreatic amylase. The gene encoding this inhibitor has been cloned and its sequence has been analysed to find a possible explanation for this specificity.

Keywords: *S. incertulas* walker, α -amylase, cDNA cloning, homology modeling, wheat dimeric α -amylase inhibitor, sequence analysis

International Journal of Insect Science 2009:1 29–44

This article is available from <http://www.la-press.com>.

© the authors, licensee Libertas Academica Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://www.creativecommons.org/licenses/by/2.0>) which permits unrestricted use, distribution and reproduction provided the original work is properly cited.



Introduction

Scirpophaga incertulas Walker (Lepidoptera: Pyralidae), commonly known as yellow stem borer, is a predominant monophagous pest of rice, which causes 5% to 30% loss of the rice crop.¹ The larvae feed voraciously on stem parts and complete their entire growth and development within the stem and hence are not easily accessible to sprayed pesticides. Controlling yellow stem borer by insecticides is uneconomical and ecologically unsafe. Therefore, the development and use of rice varieties tolerant to this pest has been receiving increasing attention in recent years. No completely known source of resistance to this insect pest has been identified, in spite of screening all the germplasm at the International Rice Research Institute (IRRI) in the Philippines.^{2,3} Moreover, tolerance to yellow stem borer is polygenic in nature²⁻⁴ due to which, during breeding, the tolerance trait gets diluted on repeated selection. Hence, in such cases, where a source of total resistance is unavailable, the strategy to increase the host plant resistance/tolerance has been through genetic engineering, by developing rice varieties expressing genes encoding insecticidal toxins.

In case of *S. incertulas* Walker (Lepidoptera: Pyralidae), considerable success has been achieved using *Bt* genes.⁵⁻²⁷ Though presently larvae of this rice pest grown on rice varieties transformed with *Bt* genes have shown 100% mortality, there is a need to explore alternate strategies to attain durable resistance, because a major problem that could result from the extensive use of insect-resistant transgenic plants is the evolution of resistance to *Bt* toxin as has been observed in different pest populations.²⁸⁻³⁷ It is quite possible that *S. incertulas* also may develop mechanisms to overcome the toxic effects of *Bt* toxins. Hence, it is necessary to develop such alternate strategies wherein the probability of resistance evolving could be substantially reduced and resistance would be durable.

α -amylases play a central role in the carbohydrate metabolism in those insects that live on seeds and plant parts. The larvae which feed voraciously on the stem parts and complete their development within the stem depend to a large extent on their α -amylases for their survival.^{1,8} Hence, proteinaceous inhibitors to the α -amylases would be attractive candidates for the control of yellow stem borer larvae. Purified monomeric and dimeric α -amylase inhibitor fractions from

a local variety of wheat seeds (variety MP Sehore) specifically inhibit the α -amylase activity from this pest. These characteristics would make these inhibitors attractive candidates for genetic engineering to develop rice transgenics with improved tolerance to *S. incertulas*. The reported monomeric class of α -amylase inhibitors from wheat has been shown to specifically inhibit the amylases from insects,³⁸ whereas the reported dimeric α -amylase inhibitors from wheat inhibit insect amylases and also human salivary amylase and porcine pancreatic amylase.³⁸⁻⁴¹ However, the dimeric inhibitor fraction isolated from MP Sehore is specific for the *S. incertulas* amylase and does not inhibit porcine pancreatic amylase and human salivary amylase.

An understanding of the molecular basis of the specificity of inhibition between the α -amylase of this rice pest and these wheat inhibitors is an important step before initiating work for developing transgenic rice expressing such an inhibitor. As a first step towards this goal, we describe the cloning and sequence analysis of the cDNA of the α -amylase from this rice pest. The three-dimensional structure of the amylase has been predicted from its translated amino acid sequence, by homology modeling, using the structure of the amylase from *Tenebrio molitor* (Coleoptera: Tenebrionidae). The gene encoding the dimeric α -amylase inhibitor from wheat variety MP Sehore has been cloned and its sequence has been compared with the sequence of the reported dimeric inhibitor, which has been modeled with the amylase from human saliva and yellow mealworm, to find a possible explanation for this specificity.

Materials and Methods

Collection of larvae of *S. incertulas* Walker (Lepidoptera: Pyralidae)

Live larvae in the third and fourth instar stage were collected from the rice fields at Roha and Alibag (Raigad district) in Maharashtra State, India. Live larvae were dissected from infested rice stems and immediately immersed in RNAlater reagent (Qiagen) and stored at -70°C , in aliquots of 10 larvae (~ 100 mg)/vial.

Preparation of crude α -amylase from *S. incertulas* Walker

One gram of insect larvae was crushed in liquid nitrogen in a mortar and pestle. The powdered material

was suspended in five volumes of 150 mM NaCl saline or 150 mM Tris-HCl pH 7.5 and centrifuged at 10,000 rpm for 10 min at 4°C in a RC5-Sorvall centrifuge. The pellets were discarded and the supernatant was assayed for enzyme activity. It was observed that extraction with 150 mM NaCl yielded 37% more enzyme activity as compared with 150 mM Tris-HCl pH 7.5. So, all further extractions were carried out in 150 mM NaCl. The NaCl extract containing the crude amylase enzyme was dialyzed against 10 mM sodium-phosphate buffer (pH 7.0) and then concentrated by lyophilization to one fourth its volume and dissolved in 10 mM sodium-phosphate buffer pH 7.0 and then re-dialyzed against the same buffer. The recovery on lyophilization was 88%. As the extract was highly pigmented, the pigment was removed using ethyl alcohol as follows: 10 ml of the concentrated extract was precipitated by addition of three volumes of chilled ethanol. The precipitate was centrifuged at 10,000 rpm in a RC5 Sorvall centrifuge at 4°C and the resultant precipitate was suspended in a total volume of approximately 16 ml in four consecutive steps of 4 ml each. This solution was lyophilized, dialyzed against 10 mM sodium-phosphate buffer pH 7.0 and once again subjected to ethyl alcohol precipitation to remove the residual pigment. The removal of pigment was monitored spectrophotometrically at 410 nm. The recovery of the enzyme was 78%. This partially purified enzyme was used for preliminary characterization of its properties.

Assay for amylase activity

The amylase activity was determined by the dinitrosalicylic acid (DNS) method.⁴² The reaction mixture contained 0.25 ml of suitably diluted enzyme in sodium-phosphate buffer (50 mM, pH 8.0) and 0.25 ml of (1% w/v) starch solution for amylase activity. The reaction was terminated by the addition of 0.5 ml DNSA. The tubes were heated in a boiling water bath for 5 min and the color intensity was read at 540 nm after dilution with 5 ml of distilled water. A standard glucose curve was used for calculating enzyme activities. The linearity curve of the enzyme was determined for calculation of unit activity. One IU of amylase activity is defined as the amount of enzyme required to produce 1 μ mole of glucose/min.

Isolation and purification of the monomeric and dimeric α -amylase inhibitors from wheat (Variety: MP Sehore)

Extraction of α -amylase inhibitors from wheat flour
Extraction of wheat α -amylase inhibitors was carried out as described.⁴³ In a preliminary experiment it was observed that a 150 mM NaCl solution was more effective in extraction of the α -amylase inhibitors from wheat flour than 100 mM Tris/HCl buffer pH 7.6. Hence, 150 mM NaCl was used for extraction of the α -amylase inhibitors. Hundred grams of finely ground whole wheat flour (Variety MP Sehore) was stirred in 300 ml of 150 mM NaCl solutions at room temperature for 3 h. The suspension was then centrifuged for 10,000 rpm for 20 min in a RC-Sorvall centrifuge at 4°C. The supernatant liquid was subjected to ammonium sulphate fractionation to isolate the α -amylase inhibitors.

Ammonium sulphate precipitation of α -amylase inhibitors

The crude α -amylase inhibitor solution was first precipitated with 0–0.4 M $(\text{NH}_4)_2\text{SO}_4$ to remove non-proteinaceous components. The resultant supernatant liquid after centrifugation was salted out with 0.4–2.0 M $(\text{NH}_4)_2\text{SO}_4$. The precipitate was collected by centrifugation at 10,000 rpm for 10 min and dissolved in one tenth the volume of double distilled water and dialyzed against water for 24 h with four changes. The inhibitor fraction was concentrated by lyophilization. Since the crude preparation had endogenous α -amylase activity, the inhibitor solution was boiled for 1 min at 100°C to inactivate the endogenous α -amylase activity.

Purification of α -amylase inhibitors by FPLC Superose-12 Gel permeation column chromatography

The crude ammonium sulphate fraction of the wheat α -amylase inhibitor containing 6 mg protein and 7000 units of inhibitor activity in 300 μ l was loaded on a FPLC-Superose 12 gel permeation column (10 \times 300 mM). Phosphate buffer pH 7.4 of composition 137 mM-NaCl, 3 mM-KCl, 1 mM-KH₂PO₄, 8 mM-Na₂HPO₄ was used. The flow rate of the column was maintained at 18.5 ml/h. A total of twenty-five fractions of one ml each were collected and the individual fractions were assayed for inhibitor activity.

The FPLC profile revealed the presence of five distinct peaks (Fig. 1a). Inhibitor activity was detected only in peak three and no activity was detected in any of the other peaks. Native PAGE of the third peak showed the presence of two bands corresponding to the monomer and dimer. In order to separate the dimer and monomer, the individual FPLC fractions of peak three were collected from several runs, dialyzed, lyophilized and loaded on Sephadex G-50 gel permeation column chromatography. The buffer system used was the same as described for FPLC. The flow rate of the column was maintained at 15 ml/h. Two peaks were obtained as shown in Figure 1b. The two peak fractions were analyzed by native PAGE. The two samples were found to be homogeneous and corresponded to the dimer and monomer. (Fig. 1c). Other FPLC fractions were processed similarly.

Assay for inhibitor activity

The assay for determining inhibitor activity was carried out essentially as described for amylase activity excepting that the reaction mixture contained suitable aliquots of seed extract containing inhibitor.

One unit of inhibitor activity is defined as the amount of inhibitor required to inhibit 50% of enzyme activity. Commercial dimeric α -amylase inhibitor from wheat (Sigma Catalogue No. A1520) was used as a control as it inhibits human salivary α -amylase (Sigma Catalogue No. A0521) and porcine pancreatic α -amylase (Sigma Catalogue No. A3176).

PAGE electrophoresis

Non-denaturing gel electrophoresis was performed according to the procedure described by Davis⁴⁴ using a 10% separating gel at pH 8.9. Bromophenol blue was used as the tracking dye.

Molecular weight determination of the inhibitors by MALDI-TOF

Gel electrophoresis was performed.⁴⁴ After electrophoresis the protein bands were excised from the gel and transferred into two separate vials. Sufficient extraction solution (100 μ l) consisting of a mixture of formic acid/glass distilled water/2-propanol (1:3:2) was added to the crushed gel to completely cover the gel pieces. The tubes were closed and vigorously shaken at room temperature for 4 to 8 h. After vortexing,




Figure 1a. Profile of FPLC- gel permeation column chromatography (Superose 12 prep. Grade).




Figure 1b. Profile of G 50 gel permeation column chromatography.




Figure 1c. Native PAGE of purified monomeric and dimeric α -amylase inhibitors.



the sample was microfuged. The supernatant was retrieved with a micropipette. The crushed gel was washed once with an equal volume of fresh extraction solution and the washes were combined with the supernatant. The two supernatants were concentrated and subjected to MALDI-TOF.⁴⁵ MALDI-TOF analysis showed that the molecular weights of the dimer and monomer were 13,330 and 13,159 Daltons respectively, suggesting that band 1 was a dimer and band 2 was a monomer.

Cloning of the gene encoding the dimeric inhibitor from wheat (Variety: MP Sehore)

For amplifying the dimeric α -amylase inhibitor from wheat (variety MP Sehore), forward and reverse primers were designed according to Wang et al.⁴⁶ Total DNA to be used as a template in the PCR reaction was isolated according to Dellaporta et al.⁴⁷ The conditions for amplification were the same as described by Wang et al.⁴⁶ The amplification product was cloned into pCR4 TOPO vector (Invitrogen) and was transformed into *E. coli* Top10 cells (Invitrogen) as per the manufacturer's instructions. The positive clones were sequenced and the amino acid sequence of the dimeric α -amylase inhibitor was deduced from the nucleotide sequence using the 'ORF Finder' program at the NCBI site (<http://www.ncbi.nlm.nih.gov/>

projects/gorf). The Gene Bank accession number of the dimeric α -amylase inhibitor from wheat variety MP Sehore is GQ374443 (Fig. 2). The first thirty amino acids represent the signal sequence as determined by SignalP version 3.0 (<http://www.cbs.dtu.dk/services/SignalP/>).

RNA extraction

100 mg of larvae preserved in RNAlater (Qiagen) were used each time for extraction of RNA. Total RNA was extracted from the larvae using Trizol reagent (Invitrogen) according to the manufacturer's instructions. The quality of the RNA preparation was judged by formaldehyde agarose gel electrophoresis.⁴⁸

Reverse transcription—polymerase chain reaction and cDNA cloning

The full length cDNA of the α -amylase gene, henceforth designated as SiAmy, was cloned as follows:

The following four pairs of forward (F1–F4) and reverse primers (R1–R4) were designed, based on the conserved regions of insect alpha-amylases:

SiAmy F1–5'GTGCACCTCTTCGAATGG 3'
SiAmy R1–5'CTCAAATCGGCTGGCCAC 3'
SiAmy F2–5'GGGTGTCGTGGTTGTCGACGAA 3'

```

1  atgctcgtggcgacacccatagcggccgagtacgacgcatggagc
   M L V A T P I A A E Y D A W S
46  gttaacagtgggccctggatgtgctatccagggtatgcctttaag
   V N S G P W M C Y P G Y A F K
91  gtgccagcgctccctggctgtcgtccagtgtgaagctccagtgc
   V P A L P G C R P V L K L Q C
136  aatggcagccaggtgcccaggctgtcctaagggactgctgccag
   N G S Q V P E A V L R D C C Q
181  cagctcgcccacatcagcgagtgggtgcaggtgcggggccctctac
   Q L A H I S E W C R C G A L Y
226  agcatgttggacagcatgtataaggagcatggcgcgaggagga
   S M L D S M Y K E H G A Q E G
271  caggcagggacaggagcgttcccacgctgccggagggaggtggtg
   Q A G T G A F P R C R R E V V
316  aagctgacggcggcgagcatcacggcggctctgcaagctaccatc
   K L T A A S I T A V C K L P I
361  gtcattgatgcgtctggagatggagcgtatgtctgcaaggggtg
   V I D A S G D G A Y V C K G V
406  gccgcatacccggacgtctag 426
   A A Y P D V *

```

Figure 2. DNA sequence and the deduced protein sequence of the dimeric α -amylase inhibitor from wheat variety MP Sehore (Gene Bank Accession Number GQ374443). The first thirty amino acids (bold and highlighted in red) represent the signal peptide.

SiAmy R2–5'GACGTTGTGTTCAACCATATGGCTG 3'
SiAmy F3–5'GTCCATCTATTCGAATGG 3'
SiAmy R3–5'TCCAAATCTGCAGGCCAC 3'
SiAmy F4–5'GTTTCATCTTTTTGAGTGG 3'
SiAmy R4–5'TGCAGATCTGCCGGCCAC 3'

Total RNA (~1 µg) was reverse transcribed at 50°C for 30 min, using Superscript III One-Step RT-PCR System with Platinum Taq High Fidelity (Invitrogen) as per the manufacturer's instructions. The RT reaction was followed by one cycle of denaturation at 94°C for 2 min. The RT template was then subjected to 30 cycles of polymerase chain reaction. Each cycle comprised of 94°C for 30s, 50°C for 1 min and 68°C for 1 min. Final extension was done at 68°C for 5 min. The PCR was carried out in a Techne (Progene) thermal cycler.

Of the four pairs of primers, only SiAmy F1–SiAmy R1 gave a 545 bp amplification product. The 545 bp PCR products was cloned into pGEM-T vector (Promega) and transformed into NovaBlue competent cells (Novagen) according to the Manufacturer's instructions. Positive clones were sequenced in an automated DNA Sequencer. The BLAST search of the DNA sequences and translated amino acid sequence revealed that the inserts of the clones showed significant sequence homology to other insect α -amylases. Based on the sequence of the 545 bp insert, the following three pairs of forward and reverse primers were designed, to be used for rapid amplification of cDNA ends (RACE-PCR).

Race F1–5'AGCTAACCGGCCGTGGTGGGAGC
GGTACCAA 3'

Race F2–5'GCGGTACCAACCCATTTTCGTACAAA
CTCA 3'

Race F3–5'ACTCACCACACGATCAGGTGACAA
CA 3'

Race R1–5'CTGCGTCTATCCTGAAACCAGCAA
CTCCTA 3'

Race R2–5'ACAAGCTCGCAATTTTCGAACCCTC 3'

Race R3–5'CGCAATTTTCGAACCCTCCAAGCAT 3'

The 5' and 3' ends of SiAmy cDNA were amplified using the 5'/3' RACE kit from Invitrogen, according to the manufacturer's instructions. The PCR products were cloned into pCR4-TOPO vector and transformed into TOP 10 chemically competent cells (Invitrogen) according to the manufacturer's instructions. Positive

clones from the 5' and 3' RACE were sequenced to obtain the sequence of the full length alpha-amylase gene. The forward (SiAmy CF1) and reverse (SiAmy CR1) primers were designed based on the complete gene sequence derived from the 5' and 3' RACE PCR and the complete gene of 1506 bp was amplified using these primers.

SiAmyCF1: 5'ATGATTCTCTAATATTACTAGCC
GTG3'

SiAmyCR1: 5'TAGCCGGCTCTGAGGACCAAT3'

Sequence analysis and homology modeling

The cloned gene sequence was translated into the protein sequence using the 'ORF Finder' program at the NCBI site (<http://www.ncbi.nlm.nih.gov/projects/gorf>). The signal peptide cleavage site was determined using SignalP 3.0 (<http://www.cbs.dtu.dk/services/SignalP/>). The theoretical pI and Mw (molecular weight) were computed using the 'Compute pI/Mw tool' option on the ExPasy server (<http://www.expasy.ch/tools/>).

Sequence analyses were performed with the BLAST program of the National Centre for Biotechnology Information (NCBI), NIH, Bethesda, MD, USA. Sequence alignments were obtained using CLUSTALW.⁴⁹ The numbering of SiAmy is based on the mature protein (excludes the 15 residue signal peptide). The phylogenetic analysis was performed using the TreeTop Phylogenetic Tree Prediction Program (GeneBee—Molecular Biology Server).^{50,51} The phylogenetic tree was built using Phylip program option in the TreeTop Phylogenetic Tree Prediction Program and a bootstrap value of 100 was used.

The three-dimensional structure of the amylase of *S. incertulas* Walker (Lepidoptera: Pyralidae) was predicted from its amino acid sequence by homology modeling (<http://www.predictprotein.org/>) Structural model used, closest to this amylase, was that of the amylase from *T. molitor* L. (pdb:1jae). Display of the structures and superposition were carried out using QUANTA (Accelrys, USA) on a silicon graphics workstation running operating system iris 6.3. The geometry of the predicted model was checked using the program PROCHECK.⁵² When the model was checked against ideal geometry



parameters more than 99% of the residues were falling in the allowed regions of the Ramachandran plot. More than 99% of bond lengths and more than 91% of bond angles were within limits. The overall G factor -0.13 (dihedrals: -0.07 , covalent: -0.21) output by the program shows that the model has acceptable geometry.

Results and Discussion

Specificity of the dimeric α -amylase inhibitor from wheat variety MP Sehore

The activity of the purified dimeric α -amylase inhibitor from the above variety was evaluated against human salivary α -amylase and porcine pancreatic α -amylase (Table 1). The results presented in Table 1 clearly demonstrate that the dimeric α -amylase inhibitor from wheat variety MP Sehore is specific for the *S. incertulas* amylase.

Cloning and sequence analysis of SiAmy cDNA

Forward primers SiAmyF1-4 and reverse primers SiAmyR1-4 for RT-PCR were designed based on the short conserved amino acid sequences VHLFEWR/K and KHMWPA/G which are found in insect alpha-amylases.⁵³⁻⁵⁵ Of the four pairs of primers used for the RT-PCR reaction, only the primer pair SiAmyF1-SiAmyR1 amplified a 545 bp fragment. The derived amino acid sequence of this fragment showed significant homology to other insect alpha-amylases. Primers for 5' and 3' RACE were designed based on this primary fragment and the fragment was extended in both directions by RACE-PCR. SiAmyCF1 and SiAmyCR1 were designed based on the 5' and 3' RACE derived sequences and the complete gene of 1506 bp was amplified. Translation

of this sequence gave a protein with an open reading frame of 502 amino acids, which showed significant homology with α -amylases from other insects. This clone was designated as SiAmy (Gene Bank accession number EF672102). (Fig. 3). The signal peptide cleavage site was found using SignalP 3.0 (<http://www.cbs.dtu.dk/services/SignalP/>). Cleavage of the 15 amino acid long signal peptide would leave a mature protein of length 487 amino acids with a predicted molecular weight of 54,955 Daltons and theoretical pI of 5.39.

A phylogenetic tree was generated using Phylip in the TreeTop Phylogeny prediction program (Fig. 4). SiAmy shows 69%–73% identity with the α -amylases from other lepidopterans and 49%–59% identity with flies and beetles. (Fig. 4).

Figure 5a shows the clustal alignment of SiAmy with the α -amylase from *Tenebrio molitor* (L.) (Coleoptera: Tenebrionidae) (TMA) and Figure 5b shows the clustal alignment of SiAmy with the α -amylases from a few lepidopterans as well as with TMA and porcine pancreatic amylase (PPA). In the sequence alignment of SiAmy and TMA (Fig. 5a) as well as in the superposition of structures (Fig. 6), the three active site residues (Asp 185, Glu 222 and Asp 287) of *T. molitor* amylase perfectly match with the corresponding residues (Asp 196, Glu 233 and Asp 298) of *S. incertulas* amylase (The numbering in SiAmy excludes the signal peptide). This modeling and comparison has helped us to identify the putative active site residues of *S. incertulas* amylase enzyme. A comparison with the α -amylases of PPA, TMA and other insect α -amylases shows that His 104, His 200 and His 297 involved in substrate binding, Asn 103, Arg 157, Asp 166 and His 200 involved in Ca^{2+} binding and Asn 296 and Arg 194 involved in Cl^- binding are conserved in SiAmy. His 319 (in PPA) which is conserved in all mammalian amylases is present as His 303 in SiAmy.

Table 1. Specificity of the wheat dimeric alpha-amylase inhibitor from wheat variety MP Sehore.

Source	<i>S. incertulas</i> α -amylase (units/mg)	Porcine α -amylase (units/mg) (Sigma catalogue No. A3176)	Human α -amylase (units/mg) (Sigma catalogue No. A0521)
Commercial wheat inhibitor (dimer) (Sigma Catalogue No. A1520)	81.08	83.32	72.18
G-50 Purified monomer	1589.42	ND	ND

Abbreviation: ND, No Inhibition detected.



```
1 atgattcctctaataattactagccgtggccgtatttgctcgtgcg
M I P L I L L A V A V F A R A
46 gctaacgattacaagaatccttactatgtgccaggcaggtcagtg
A N D Y K N P Y Y V P G R S V
91 aatgtccacctcttcgagtggaagtgaggatagctgaagag
N V H L F E W K W E D I A E E
136 tgtgagcgccttttaggaccaatggtttgggtgggtacagatt
C E R F L G P N G F G G V Q I
181 tcaccacaaacgaaaacgtgataatgggtcagtaaccggccg
S P P N E N V I I W S A N R P
226 tgggtgggagcgtaccaacccatttcgtacaaactcaccacacga
W W E R Y Q P I S Y K L T T R
271 tcaggtgacaacacgcagctggccaacatggtgagcagatgtaac
S G D N T Q L A N M L R R C N
316 aacgttggagtcagaatataatgctgatgctgcatcaatcatatg
N V G V R I Y V D A V I N H M
361 actggggatcctccagagaatgtggggactgccggcagtgactgct
T G D P P E N V G T A G S T A
406 accttcaatgaatggcactacctaactgtaccgtacagaagagaa
T F N E W H Y P T V P Y R R E
451 cattttaactggcctcttggatcagtggtaccgactaccaa
H F N W P S C G I D G T D Y Q
496 accaatgcttggagggttcgaaattgcgagcttgggtttaaana
T N A W R V R N C E L V G L K
541 gatttagatcagtcattgatcatgtgcgcaaatgattgtggat
D L D Q S I D H V R N M I V D
586 ttcatgaacactctgattgatttaggagttgctggttcaggata
F M N T L I D L G V A G F R I
631 gacgcagcaaacacatgtggccagaagatctccgaataatctac
D A A K H M W P E D L R I I Y
676 gatcgactacataatctgagcacagaccatggcttcccattgaa
D R L H N L S T D H G F P L N
721 gcccgccatacatttaccagaagtcattgattatgggtggcga
A R P Y I Y Q E V I D Y G G E
766 gctgtcagtagagaagagtataactccaattggagctgtgactgag
A V S R E E Y T P I G A V T E
811 ttcaaggctggcatggagttgtctaactgcttcagaggacataat
F K A G M E L S N C F R G H N
856 cagctccgatggctcgtcagctggggagcgcgcttccgattgctg
Q L R W L V S W G A P F G L L
901 gagagcagagacgcttggacttattgataatcacgacaatgaa
E S R D A L T F I D N H D N E
946 agaggtcacggaggtgggtggcgggtgtgcttacttacaagcaacct
R G H G G G G V L T Y K Q P
991 agaccatataaagcagctatagccttccgtttagcacatccttat
R P Y K A A I A F L L A H P Y
1036 ggagaaccacaaatcatgagtagtttcgagtttgggaccagaa
G E P Q I M S S F E F W D P E
1081 ataggtccgcaatgaatagtcagggtaacataatccccttct
I G P P M N S Q G N I I S P S
1126 atcaactcagatggcacttggtaacgggtggatagccagcac
I N S D G T C G N G W I C Q H
1171 aggtggcgacagatctactccatgggtgctggtcaggaatgccgct
R W R Q I Y S M V S F R N A A
1216 ggcagcggggcccttaacaactgggtggacaatgggtccaaccaa
G S G G L N N W W D N G S N Q
1261 atagcattttgcccggagaaaagcgttttatagcatttaacaat
I A F C R G E S A F I A F N N
1306 gacagccgggattttaaaccgaaaatagcagacatgtctcccagct
D S R D L N E N M Q T C L P A
1351 gggcagttattgcgacgtgatcacaggcagagaaaacggaagttcc
G Q Y C D V I T G E R N G S S
1396 tgctccggtaagggtggtgacgggtgggagagaacgggtgggtcat
C S G K V V T V G E N G W A H
1441 gtctacgtaggagcgaagactacgatatgatgttagcaatacat
V Y V G A Q D Y D M M L A I H
1486 attggtcctcagagccggctatga 1509
I G P Q S R L *
```

Figure 3. cDNA sequence and the deduced protein sequence of the *S. incertulas* α -amylase SiAmy (Gene Bank Accession Number EF672102). The first fifteen amino acids (bold and highlighted in red) represent the signal peptide. The three amino acids (bold and highlighted in yellow) are conserved in α -amylases and are part of the active site.




Figure 4. Phylogenetic tree of α -amylases of the following insects. *Diatraea saccharalis* (AY333761), *S. incertulas* Walker (EF672102), *Helicoverpa armigera* (EF600049), *Spodoptera frugiperda* (AA013754), *Ostrinia nubilalis* (AAA03715), *D. pseudoobscura* (X76240), *Megaselia scalaris* (AA017923), *Bibio marci* (AAL92553), *Tribolium castaneum* (U04271), *Blaps mucronata* (AF462603), *Tenebrio molitor* (P56634), *Zabrotes subfasciatus* (AAF73435), *Callosobruchus chinensis* (AB11048), *Anthonomus grandis* (AF527876). A bootstrap value of 100 was used and the bootstrap values are expressed in percentages at the nodes.

This histidine residue is conserved in lepidopterans and flies but not in beetles. In vertebrates, beetles and flies, the Cl^- ion is coordinated by six ligands i.e. Arg 210(PPA) and Arg 351(PPA) in a bidentate mode and Asn 312(PPA) in a unidentate mode.⁵⁶ In bacterial α -amylases, Arg 351(PPA) is replaced by a lysine residue.⁵⁷ In the α -amylases of lepidopterans including SiAmy, Arg 351/Lys 351 is replaced by a glutamine residue (Q 334 in SiAmy). It has been suggested that such a replacement by a glutamine residue makes the α -amylase independent of chloride ions for activity and also shifts the optimum pH towards alkaline values^{58,62} which is an advantage for lepidopterans whose gut environment is alkaline. The α -amylases of beetles and flies have either Arg or Lys in this position, which

is suitable for activity in an acidic gut environment. Also, the binding of chloride ion to α -amylases has been suggested to broaden the pH activity profile such that the pH optimum shifts from acidic values to slight alkaline values.^{58,62}

A glycine rich 'flexible loop' (residues 318–323 in PPA) which forms the surface edge of the substrate binding cleft in PPA,⁵⁹ is also present in SiAmy (residues 302–308) and other lepidopterans (Fig. 5b), but is absent in TMA. This loop has been suggested to hold residues assisting catalysis in the appropriate orientation and to induce a trap-release mechanism of substrate and products.^{59–61} Though the amylases of *S. incertulas* Walker (Lepidoptera: Pyralidae) appear to be chloride independent (unpublished

S. INCERTULAS	1	ANDYKNPYYPGRSVNVHLFEWKWEDIAEEECERFLGPNFGGGVQISPPNE	50
T. MOLITOR	1	EKDANFASGRNSIVHLFEWKWNDADECERFLQPQGFGGVQISPPNE	47
S. INCERTULAS	51	NV I IWSANR P WWERYQPI SYKLTTRSGDNTQLANMLRRCNNVGVRIYVDA	100
T. MOLITOR	48	YLV--ADGR P WWERYQPVSYIINTRSGDESAFTDMTRRCNDAGVRIYVDA	95
S. INCERTULAS	101	VINHM TGDP PE N VGSTAGSTATFNEWHYPTVPYRREHF NWPSCGIDGTDYQ	150
T. MOLITOR	96	VINHM TG--MNG VGTSGSSADHDGMNYPVPYGS GFHSP-CEVN--NYQ	140
S. INCERTULAS	151	TNA WRVRNCELVGLKDL DQ SIDHVRNMIVDFMNTLIDLG VAGFRID AAKH	200
T. MOLITOR	141	- DA DNVRNCELVGLRDLN QGS DYVRGV LIDY MNHMIDLG VAGFRV AAKH	189
S. INCERTULAS	201	MWPEDLRIIYDRLHNLSTDHGFPLNARPYI YQ VIDYGG EAVS RREYTP I	250
T. MOLITOR	190	MSPGDLSVIFSGLKNLNTDYGFADGAR PFYQ VIDLG GEAISK NEYTG F	239
S. INCERTULAS	251	GAVTEFKAGMELSNCFR GH NQLRWLVSWGAPF GLLES RDALTFIDNH NE	300
T. MOLITOR	240	GCVLEFQFGVSLGNAF QGG NQLK NLAN WGP EWGLLE GLDAV VFV DNH Q	289
S. INCERTULAS	301	RGHGGG GGVLT YKQ PRPYKAAIAFLLAHPY GEPQ IMSSFE FWD PEIG PPM	350
T. MOLITOR	290	R--TGG SQILTYK NP KPYKMAIA FMLA HPYGTTRIMSS FDF TDND Q GG PPQ	337
S. INCERTULAS	351	NSQGNII SP SINSDGTCGNGWICQHRWRQIYSMV SFR NAAGSGGLNN WWD	400
T. MOLITOR	338	DGSGNLIS PG INDDNTCSNGY VCE HRWRQVYGM VGF RNA VEGTQ VEN WWS	387
S. INCERTULAS	401	NGSNQIA FC RGESAFIAF N NSRD LN ENMQ TCL PAGQYCDVITGER NGSS	450
T. MOLITOR	388	NDDNQIA FS RG SQ GFVA FT NGG-DLNQNL TGL PAGTYCDV IS GEL S GG S	436
S. INCERTULAS	451	CSGKVVTVGENGW AH VYVGAQDYDMMLAIHIG PQ SRL 487	
T. MOLITOR	437	CTGKSVTVGDNGSADISL GS AEDDGVLAIH VNA KL 471	

Figure 5a. ClustalW alignment of the *S. incertulas* amylase SiAmy (EF672102) with the amylase of *Tenebrio molitor* (TMA) (P56634). The three catalytic residues D196, E233 and D298 of SiAmy and the corresponding ones in *T. molitor* amylase (TMA) are highlighted in yellow. The numbering of SiAmy excludes the signal peptide.

results), residues Ser 354, His 400 and Glu 42 in PPA which are strictly conserved in chloride-dependent α -amylases, are present as Ser 337, His 375 and Glu 30 in SiAmy. This triad has been shown to mimic the active site of proteases and lipases and was first described in the α -amylase from *Alteromonas haloplanctis*.⁵⁷ Though there is no evidence for a catalytic role, this triad has been suggested to play a pivotal role in stabilizing the structure of chloride-dependent α -amylases.⁶²

Four disulfide bonds required for proper folding of the SiAmy structure are formed by the conserved cysteines ie. C31–C89, C142–C159, C367–C373 and C439–C451. These bonds are found in all chloride-dependent α -amylases.⁶² The C409–C432 disulfide

bond in SiAmy (absent in TMA) bridges the β -strands in domain C and has been implicated in improving the rigidity of the globular C domain.⁶² As seen in Figure 5b, the sequence DGX₁X₂AIH in the C-terminal tail does not completely follow the general consensus.⁶³ In SiAmy and lepidopterans, the glycine is replaced with methionine which is also a non-polar amino acid. The two amino acids preceding AIH are hydrophobic i.e. Valine, Leucine or Methionine. In SiAmy and the α -amylases of *Diatraea saccharalis* and *Ostrinia nubilalis*, AIH is followed by hydrophobic amino acids isoleucine or valine. In case of *Helicoverpa armigera* and *Spodoptera frugiperda*, AIH is followed by threonine, a hydrophilic amino acid. The next three amino acids are not necessarily hydrophilic.

Homology modeling

The sequence alignment of *S. incertulas* Walker (Lepidoptera: Pyralidae) amylase (SiAmy) with the sequence of the amylase from *Tenebrio molitor* L. (TMA) used for homology modeling is shown in Figure 5a. The chain length (487) of SiAmy is longer than that of TMA (471). Consequently, in addition to N- and C-terminals there are four major regions wherein SiAmy has insertions of amino acids compared to TMA. Manual superposition using QUANTA (Accelrys) of the modeled structure of SiAmy

on TMA shows that the sizes of four loops in *S. incertulas* Walker (Lepidoptera: Pyralidae) have increased due to insertion of additional residues (Fig. 4).

Based on Blastp, SiAmy shows only 56% homology to the amylase of *T. molitor*. The only insect amylase whose 3D structure is well studied is that of *T. molitor*. The 3D structure of *T. molitor* amylase complexed with various inhibitors is also well studied. The maximum homology between SiAmy and the amylases from other lepidopterans is 76%. Several of the pests, which infest several crops, are lepidopterans. Hence, a 3D




Figure 6. Superposition of the modeled structure of *Scirpophaga incertulas* Walker amylase (green) on the reported structure of *Tenebrio molitor* L. α -amylase (red, pdb: 1jae). The four loop regions (1) 53–59 (2) 106–112 (3) 138–152 (4) 300–306 where loops deviate due to insertion of residues are shown liquorized (numbering based on SiAmy excluding the 15 residue signal peptide). Superposition of three active site residues are also shown (residue numbers correspond to TMA/SiAmy) (Refer Fig. 5a also).



model developed for SiAmy will serve to compare lepidopteran amylases more accurately.

The wheat inhibitor reported here is sequentially placed in the class 0.19 dimer type inhibitors of wheat. Sequence comparison showed that it differs from the dimeric α -amylase inhibitor from *Triticum turgidum* subsp. *dicoccoides* (GeneBank ACP40697.1) by just three amino acids. Similarly it differs by 8 residues with respect to sequence of 0.19 inhibitor structure reported in protein data bank (1HSS) and also by the presence of 17 extra residues at the N-terminus. The wheat inhibitor characterized here has an extra C-terminal valine residue compared to others of this type. We have experimental evidence to show that the inhibitor protein isolated by us is specific for insect amylases and does not inhibit human salivary amylase (HAS) (Table 1). It is reported that among the wheat amylase inhibitors 0.19 type class shows wider specificity for both insect and mammalian α -amylases whereas 0.28 class is more specific for insect amylases.³⁸ Comparative modeling studies between 0.19 and 0.28 types inhibitors for insect and mammalian amylases⁶⁴ have shown that the residues 104–107, VVDA of 0.19 mutated to a shorter PNP in 0.28 in an insertion loop which essentially may be determining the exclusive specificity of latter type for insect amylase. Similar conclusion was drawn in a different modeling study using four different inhibitors from wheat.³⁹ We have found out that the residues of loop region 104–107 in our inhibitor are VIDA. However, it will be hard to believe that this single change of valine to isoleucine alone can change it to become non-inhibitory for human salivary amylase. Although both N- and C-terminal residues are reported to be important,⁶⁴ in a proposed global mechanism of inhibition by this class of inhibitors, whether the extra valine at C-terminus in our inhibitor decides its non-specificity for HAS is not clear from the present information. Further structural studies only can provide an exact description of the inhibitor-amylase interactions and specificity.

An ideal inhibitor should be able to specifically inhibit the insect amylases. We have identified and purified a dimeric inhibitor from a local variety of wheat which unlike other dimeric inhibitors reported to date, does not inhibit porcine α -amylase or human salivary α -amylase but specifically inhibits the *S. incertulas* amylase. Elucidation of the molecular

basis of specificity of interaction between SiAmy and the wheat inhibitor will not only help understand the specificity of the wheat inhibitor but can help to design inhibitors specific for other lepidopteran amylases.

Acknowledgement

The authors wish to acknowledge the Department of Biotechnology, New Delhi, India, for financial support, for carrying out this work. They also thank Dr. Mahesh Kulkarni for his help in the MALDI-TOF analysis.

Disclosures

The authors report no conflicts of interest.

References

- Herdt RW. Rice Biotechnology: Research priorities for rice Biotechnology. Eds: Khush GS, Toenniessen GH. Wallingford, UK, CAB International; 1991:19–54.
- Khush GS. Disease and insect resistance in rice. *Adv Agron.* 1977;29: 265–341.
- Chaudhary RC, Khush GS, Heinrichs EA. Varietal resistance to rice stem borers in Asia. *Insect Sci Appl.* 1984;5:447–63.
- Koshiary MA, Pan CL, Hak GE, et al. A study on the resistance of rice to stem borer infestation. *Int Rice Comm Newsletter.* 1957;5:23–5.
- Fujimoto H, Itoh K, Yamamoto M, Kyojuzuka J, Shimamoto K. Insect resistant rice generated by introduction of a modified delta-endotoxin gene of *Bacillus thuringiensis*. *Bio/Technology.* 1993;11:1151–5.
- Wunn J, Kloeti A, Burkhardt PK, et al. Transgenic indica rice beeding line IR 58 expressing a synthetic *cry IAb* gene from *Bacillus thuringiensis* provides effective insect pest control. *Bio/Technology.* 1996;14:171–6.
- Ghareyazie B, Alinia F, Menguito CA, et al. Enhanced resistance to two stem borers in an aromatic rice containing a synthetic *cry IA(b)* gene. *Mol Breeding.* 1997;3(5):401–14.
- Nayak P, Basu D, Das S, et al. Transgenic elite indica rice plants expressing *Cry IAc* delta-endotoxin of *Bacillus thuringiensis* are resistant against yellow stem borer (*Scirpophaga incertulas*). *Proc Natl Acad Sci U S A.* 1997;94:2111–6.
- Wu C, Fan Y, Zhang C, Oliva N, Datta SK. Transgenic fertile japonica rice plants expressing a modified *cry IA(b)* gene resistant to yellow stem borer. *Plant Cell Rep.* 1997;17(2):129–32.
- Alam MF, Datta K, Abrigo E, Vasquez A, Senadhira D, Datta SK. Production of transgenic deep water indica rice plants expressing a synthetic *Bacillus thuringiensis Cry IA (b)* gene with enhanced resistance to yellow stem borer. *Plant Sci.* 1998;35:25–30.
- Cheng X, Sardana R, Kaplan H, Altosaar I. *Agrobacterium*-transformed rice plants expressing synthetic *cry IA(b)* and *cry IA(c)* genes are highly toxic to striped stem borer and yellow stem borer. *Proc Natl Acad Sci U S A.* 1998;95(6):2767–72.
- Tu J, Datta K, Alam MF, Fan Y, Khush GS, Datta SK. Expression and function of a hybrid Bt toxin gene in transgenic rice conferring resistance to insect pest. *Plant Biotechnol.* 1998;15(4):195–203.
- Maqbool SB, Husnain T, Riazuddin S, Masson L, Christou P. Effective control of yellow stem borer and rice leaf folder in transgenic rice indica varieties Basmati 370 and M7 using the novel delta-endotoxin *cry 2A Bacillus thuringiensis* gene. *Mol Breeding.* 1998;4(6):501–7.
- Maqbool SB, Riazuddin S, Nguyen TL, Gatehouse AMR, Gatehouse JA, Christou P. Expression of multiple insecticidal genes confers broad resistance against a range of different rice pests. *Mol Breeding.* 2001;7:85–93.

15. Datta K, Vasquez A, Tu J, et al. Constitutive and tissue specific differential expression of the *cry IA(b)* gene in transgenic rice plants conferring resistance to rice insect pest. *Theor Appl Genet.* 1998;97(1–2):20–30.
16. Datta K, Baisakh N, Thet KM, Tu J, Datta S. Pyramiding transgenes for multiple resistance in rice against bacterial blight, yellow stem borer and sheath blight. *Theor Appl Genet.* 2002;106(1):1–8.
17. Sudha SN, Jayakumar R, Sekar V. Introduction and expression of the *cry IAc* gene of *Bacillus thuringiensis* in a cereal-associated bacterium *Bacillus polymyxa*. *Curr Microbio.* 1999;38(3):163–7.
18. Xiang YB, Cheng XY, Liang ZQ, Shu QY, Gao MW, Altosaar I. Agrobacterium mediated transformation of insecticidal *Bacillus thuringiensis cry IAb* and *cry IAc* genes and their expression in rice. *Chinese J Biotechnol.* 1999;15:494–500.
19. Breieler JC, Marfa V, Royer M, et al. Expression of a *Bacillus thuringiensis cry IB* synthetic gene protects Mediterranean rice against the striped stem borer. *Plant Cell Repo.* 2000;19(12):1195–202.
20. Shu Q, Ye G, Cui H, et al. Transgenic rice plants with a synthetic *cry IAb* gene from *Bacillus thuringiensis* were highly resistant to eight lepidopteran rice pests. *Mol Breeding.* 2000;6(4):433–9.
21. Ye GY, Tu J, Hu C, Datta K, Datta SK. Transgenic IR72 with fused *Bt* gene *Cry IAb/Cry IAc* from *Bacillus thuringiensis* is resistant against four lepidopteran species under field conditions. *Jpn Soc Plant Cell Mol Biol.* 2001a;18(2):25–133.
22. Ye GY, Shu QY, Yao HW, et al. Field evaluation of resistance of transgenic rice containing a synthetic *cry IAb* gene from *Bacillus thuringiensis* Berliner to two stem borers. *J Econ Entomol.* 2001b;94(1):271–6.
23. Ye G, Yao H, Shu Q, et al. High levels of stable resistance in transgenic rice with a *cry IAB* gene from *Bacillus thuringiensis* Berliner to rice leafhopper (*Cnaphalocrocis medinalis*) (guenee) under field conditions. *Crop Prot.* 2003;22:171–8.
24. Khanna HK, Raina SK. Elite indica transgenic rice plants expressing modified *Cry IAc* endotoxin of *Bacillus thuringiensis* show enhanced resistance to yellow stem borer (*Scirpophaga incertulas*). *Transgenic Res.* 2002;11(4):411–23.
25. Jiang G, Xu C, Tu J, Li X, He Y, Zhang Q. Pyramiding of insect and disease resistance genes into an elite indica, cytoplasmic male sterile restorer line of rice, Minghui 63. *Plant Breeding.* 2004;123:112–16.
26. Ramesh S, Nagadhara D, Reddy V, Rao K. Production of transgenic indica rice resistant to yellow stem borer and sap-sucking insects, using super-binary vectors of *Agrobacterium tumefaciens*. *Plant Sci.* 2004a;166:1077–85.
27. Ramesh S, Nagadhara D, Pasalu IC, et al. Development of stem borer resistant transgenic parental lines involved in the production of hybrid rice. *J Biotechnol.* 2004b;111(2):131–41.
28. Kirsch K, Schmutterer H. Low efficacy of a *Bacillus thuringiensis* (Berl.) formulation in controlling the diamondback moth, *Plutella xylostella* (L.) in the Philippines. *J Appl Entomol.* 1988;105:249–55.
29. Tabashnik BE, Cushing NL, Finson N, Johnson MW. Field development of resistance to *Bacillus thuringiensis* in the diamondback moth (Lepidoptera: Plutellidae). *J Econ Entomology.* 1990;83:1671–6.
30. Tabashnik BE, Liu YB, Finson N, Masson L, Heckel DG. One gene in diamondback moth confers resistance to four *Bacillus thuringiensis* toxins. *Proc Natl Acad Sci U S A.* 1997;94:1640–4.
31. Shelton AM Jr, Robertson L, Tang JD, et al. Resistance of the diamondback moth (Lepidoptera: Plutellidae) to *Bacillus thuringiensis* sub-species in the field. *J Econ Entomol.* 1993;86:697–705.
32. Ballester V, Escriche B, Ménsua JL, Riethmacher GW, Ferré J. Lack of cross-resistance to other *Bacillus thuringiensis* crystal proteins in a population of *Plutella xylostella* highly resistant to *Cry IA (b)*. *Biocontrol Sci Techn.* 1994;4:437–43.
33. Tabashnik E. Evolution of resistance to *Bacillus thuringiensis*. *Annu Rev Entomol.* 1994;39:47–79.
34. Oppert B, Kramer KJ, Johnson D, Upton SJ, McGaughey WH. Luminal proteinases from *Plodia interpunctella* and the hydrolysis of *Bacillus thuringiensis Cry IA(c)* protoxin. *Insect Biochem Molec.* 1996;26:571–83.
35. Huang F, Buschman LL, Higgins RA, McGaughey H. Inheritance of resistance to *Bacillus thuringiensis* toxin in the European corn borer. *Science.* 1999;284:965–7.
36. James RR, Croft BA, Strauss SH. Susceptibility of the cottonwood leaf beetle (Coleoptera: Chrysomelidae) to different strains and transgenic toxins of *Bacillus thuringiensis*. *Environ Entomol.* 1999;28:108–15.
37. Tabashnik BE, Patin AL, Dennehy TJ, et al. Frequency of resistance to *Bacillus thuringiensis* in field populations of pink bollworm. *Proc Natl Acad Sci U S A.* 2000;97:12980–4.
38. Feng G-H, Richardson M, Chen MS, Kramer KJ, Morgan TD, Reeck GR. α -amylase inhibitors from wheat: Amino acid sequences and patterns of inhibition of insect and human α -amylases. *Insect Biochem Molec.* 1996;26(5):419–26.
39. Franco OL, Rinden DJ, Melo FR, Bloch CJr, Silva CP, Grossi de Sa MF. Activity of wheat a-amylase inhibitors towards bruchid a-amylases and structural explanation of observed specificities. *Eur J Biochem.* 2000; 267:2166–73.
40. Oneda H, Lee S, Inouye K. Inhibitory effect of 0.19 alpha-amylase inhibitor from wheat kernel on the activity of porcine pancreas alpha-amylase and its thermal stability. *J Biochem.* 2004;135(3):421–7.
41. Fontanini D, Capocchi A, Muccilli V, et al. Dimeric Inhibitors of Human Salivary r-Amylase from Emmer (*Triticum dicoccon* Schrank) Seeds. *J Agr Food Chem.* 2007;55:10452–60.
42. Bernfield P. Amylases, α and β In: Colowick SP, Kaplan NO Eds. Preparation and Assay of Enzymes Methods in Enzymology, Elsevier, USA. 1955;1:149–58.
43. Petrucci T, Tomasi M, Cantagalli P, Silano V. Comparison of wheat albumin inhibitors of α -amylase and Trypsin. *Phytochemistry.* 1974;13:2487–95.
44. Davis BJ. Disc electrophoresis-method and application to human serum proteins. *Ann N Y Acad Sci.* 1964;121(2):404–27.
45. Cohen SL, Chait BT. Mass Spectrometry of Whole Proteins Eluted from Sodium Dodecyl Sulfate–Polyacrylamide Gel Electrophoresis Gels. *Anal Biochem.* 1997;247(2):257–67.
46. Wang RJ, Yan ZH, Wei YM, Nevo E, Baum BR, Zheng YL. Molecular characterization of dimeric alpha-amylase inhibitor genes in wheat and development of genome allele-specific primers for the genes located on chromosome 3BS and 3DS. *J Cereal Sci.* 2006;43:360–8.
47. Dellaporta SL, Wood J, Hicks JB. A plant minipreparation: version II. *Plant Mol Biol Rep.* 1983;1(4):19–21.
48. Sambrook J, Fritsch EF, Maniatis. Molecular Cloning. 2nd ed. Nolan C. Edr. Cold Spring Harbor Lab Press, USA; 1989.
49. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 1994;22:4673–80.
50. Chumakov KM, Yushmanov SV. The maximum topological similarity principle in molecular systematics. *Mol Genet Microbiol Virusol.* 1988;3:3–9.
51. Yushmanov SV, Chumakov KM. Algorithms of the maximum topological similarity phylogenetic trees construction. *Mol Genet Microbiol Virusol.* 1988;3:9–15.
52. Laskowski RA, MacArthur MW, Moss DS, Thornton JM. PROCHECK: a program to check the stereo-chemical quality of protein structures. *J Appl Crystallogr.* 1993;26:283–91.
53. Grossi de Sa MF, Chrispeels MJ. Molecular cloning of Bruchid (*Zabrotes subfasciatus*) α -amylase cDNA and interaction of the expressed enzymes with bean amylase inhibitors. *Insect Biochem Molec.* 1997;27(4):271–81.
54. Titarenko E, Chrispeels MJ. cDNA cloning, biochemical characterization and inhibition by plant inhibitors of the α -amylases of the Western corn rootworm, *Diabrotica virgifera virgifera*. *Insect Biochem Molec.* 2000;30:979–90.
55. MacGregor EA, Janecek S, Svensson B. Relationship of sequence and structure to specificity in the α -amylase family of enzymes. *Biochim Biophys Acta.* 2001;1546:1–20.
56. Strobl S, Maskos K, Betz M, et al. Crystal structure of yellow meal worm alpha-amylase at 1.64Å resolution. *J Mol Biol.* 1998;278:617–28.
57. Aghajari N, Feller G, Gerday C, Haser R. Crystal structures of the psychrophilic α -amylase from *Alteromonas haloplanctis* in its native form and complexed with an inhibitor. *Protein Sci.* 1998;7:564–72.
58. Feller G, Le Bussy O, Houssier C, Gerday C. Structural and functional aspects of chloride binding to *Alteromonas haloplanctis* alpha-amylase. *J Biol Chem.* 1996;271:23836–41.



59. Qian M, Haser R, Buisson G, Duee E, Payan F. The active center of a mammalian alpha-amylase. Structure of the complex of a pancreatic alpha-amylase with a carbohydrate inhibitor refined to 2.3A° resolution. *Biochemistry US*. 1994;33:6284–94.
60. Ramasubbu N, Paloth V, Luo Y, Brayer GD, Levine MJ. Structure of human salivary a-amylase at 1.6A° resolution: implications for its role in the oral cavity. *Acta Crystallogr*. 1996;D52:435–46.
61. Ultdehaag JC, Mosi R, Kalk KH, et al. X-ray structures along the reaction pathway of cyclodextrin glycosyltransferase elucidate catalysis in the alpha-amylase family. *Nat Struct Biol*. 1999;6:432–6.
62. D'Amico S, Gerday C, Feller G. Structural similarities and evolutionary relationships in chloride-dependent a-amylases. *Gene*. 2000;253:95–105.
63. Efflo PC, Folgueras-Flatschart, AV, Montor WR, Pernasetti FM, Pueyo MT, Sogayar MC. Expression of functional *Anopheles merus* α-amylases in the baculovirus/*Spodoptera frugiperda* system. *Insect Mol Biol*. 2003;12(5): 415–25.
64. Payan F. Structural basis for the inhibition of mammalian and insect α-amylases by plant protein inhibitors. *Biochim Biophys Acta*. 2004; 1696:171–80.

Publish with Libertas Academica and every scientist working in your field can read your article

“I would like to say that this is the most author-friendly editing process I have experienced in over 150 publications. Thank you most sincerely.”

“The communication between your staff and me has been terrific. Whenever progress is made with the manuscript, I receive notice. Quite honestly, I've never had such complete communication with a journal.”

“LA is different, and hopefully represents a kind of scientific publication machinery that removes the hurdles from free flow of scientific thought.”

Your paper will be:

- Available to your entire community free of charge
- Fairly and quickly peer reviewed
- Yours! You retain copyright

<http://www.la-press.com>