



## **Floristic Traits and Biogeographic Characterization of the Gennargentu Massif (Sardinia)**

Authors: Bacchetta, Gianluigi, Fenu, Giuseppe, Guarino, Riccardo, Mandis, Giovanni, Mattana, Efisio, et al.

Source: *Candollea*, 68(2) : 209-220

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2012v682a4>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Floristic traits and biogeographic characterization of the Gennargentu massif (Sardinia)

Gianluigi Bacchetta, Giuseppe Fenu, Riccardo Guarino, Giovanni Mandis, Efsio Mattana, Giovanni Nieddu<sup>†</sup> & Carmine Scudu

## Abstract

BACCHETTA, G., G. FENU, R. GUARINO, G. MANDIS, E. MATTANA, G. NIEDDU & C. SCUDU (2013). Floristic traits and biogeographic characterization of the Gennargentu massif (Sardinia). *Candollea* 68: 209-220. In English, English and French abstracts.

A study on the vascular flora of the Gennargentu Massif (Central-Eastern Sardinia) is presented. According to our results, the flora consists of 948 taxa: 686 species, 249 subspecies, 10 varieties and 3 hybrids, belonging to 97 families and 427 genera. Three taxa are new findings for the flora of Italy and eight for that of Sardinia. Life form analysis revealed, in particular, dominance of 35.65% hemicryptophytes, 34.6% therophytes, 12.13% geophytes and 11.6% (nano)-phanerophytes. As concerns chorology, the Mediterranean element is largely prevailing (68.14%), mainly represented by circum-Medit. (29.1%) and Euro-Medit. (23.07%). Endemics are 14.87% of the whole flora (141 taxa), with a large prevalence of Sardo-Corsican (39.01%) and Sardinian taxa (35.46%), i.e. 74.47% of the total. Due to the high number of taxa (9) of Gennargentu exclusive endemics and the geologic and geomorphologic peculiarities, it is here proposed a biogeographic classification for these territories serving to the identification of an autonomous sector.

## Key-words

Sardinia – Gennargentu – Floristics – Biogeography – Endemics – Mediterranean vascular flora

## Résumé

BACCHETTA, G., G. FENU, R. GUARINO, G. MANDIS, E. MATTANA, G. NIEDDU & C. SCUDU (2013). Caractérisations floristiques et biogéographiques du massif du Gennargentu (Sardaigne). *Candollea* 68: 209-220. En anglais, résumés anglais et français.

L'étude de la flore vasculaire du massif du Gennargentu (Sardaigne centre-est) est présentée. Les investigations floristiques ont établi la présence de 948 taxa, dont 686 espèces, 249 sous-espèces, 10 variétés et 3 hybrides, appartenant à 97 familles et 427 genres. Trois nouveaux taxa sont signalés pour la flore d'Italie et huit pour la flore Sarde. L'analyse du spectre biologique établit, en particulier, la dominance d'hémicryptophytes (35,65%), de thérophytes (34,6%), de géophytes (12,13%) et de (nano-)phanérophytes (11,6%). L'analyse du spectre chorologique met en évidence la prédominance de composantes méditerranéennes (68,14%), principalement représentées par des taxa circum-méditerranéens (29,1%) et euro-méditerranéens (23,07%). Les entités endémiques représentent 14,87% (141 taxa) de la flore, avec une majorité d'éléments corsico-sardes (39,01%) et sardes (35,46%) formant ensemble 74,47% des endémiques. En raison du nombre élevé de taxa (9) d'espèces endémiques exclusives du Gennargentu et des particularités géologiques et géomorphologiques, une classification biogéographique est proposée pour ces territoires servant à l'identification d'un secteur autonome.

Addresses of the authors: GB, GF, GM, EM: Centre for the Conservation of Biodiversity (CCB), Dept of Environmental and Life Sciences – University of Cagliari. Viale S. Ignazio da Laconi, 11-13 - 09123 Cagliari, Italy. Email (GB): [bacchet@unica.it](mailto:bacchet@unica.it)

RG: Dept. of Environmental Sciences and Biodiversity – University of Palermo. Via Archirafi, 38 - 90123 Palermo, Italy.

GN, CS: Via Vittorio Emanuele II, 8 - 08049 Villagrande, Italy.

Submitted on July 10, 2012. Accepted on July 18, 2013.

Edited by P. Bungener

## Introduction

The Mediterranean basin has been recognised as one of the 34 most important biodiversity hotspots, also because of its high number of endemic plant species (MITTERMEIER & al., 2004). This area not only constitutes a refuge for many relic species, but the relatively short distance of many islands and peninsulas promotes floristic exchanges and active plant speciation. MÉDAIL & DIADEMA (2009) identified the Central-Northern Sardinia as one of the 52 putative floristic refugia within the Mediterranean, i.e. places facilitating the long-term persistence of a species (one or more glacial-interglacial cycles) or of one or more of its meta-populations in a well-defined geographical area (e.g. mountain range, gorge).

Sardinia, with its 24,090 km<sup>2</sup>, is the second-largest island in the Mediterranean Sea. The prolonged isolation and high geological diversity created a wide range of habitats rich in endemic species, particularly on its mountain massifs, where the insularity is strengthened by the altitude and diversity of terrains (MÉDAIL & QUÉZEL, 1997). The Sardinian flora consists of 2408 taxa including 2295 species (CONTI & al., 2005) 168 of which are exclusive endemics (BACCHETTA & al., 2012b).

According to the biogeographic classification of the Mediterranean region proposed by RIVAS-MARTÍNEZ & al. (2002), the Italo-Tyrrhenian province is composed by three subprovinces: the Sardinian, the Corsican and the Tuscano-Calabrian. Owing to the many similarities, not only in the floristic aspects, it is here preferred to recognize the rank of biogeographical province to Corsica and Sardinia, in the frame of an Italo-Tyrrhenian superprovince extended to all over the western coast of the Italian Peninsula, from Tuscany to Calabria, as formerly proposed by LADERO ALVAREZ & al. (1987). The Sardo-Corsican province, on the contrary, can be furtherly divided into a Sardinian and a Corsican subprovince, as stated by BACCHETTA & PONTECORVO (2005). These authors, basing on their studies on the vascular endemic flora of Sulcis-Iglesiente, conferred the rank of biogeographic sector to these territories. Furthermore, a Sinisico subsector (included in the Campidano Sector) has been identified by FENU & BACCHETTA (2008) for the Sinis Peninsula (Central-Western Sardinia), while FENU & al. (2010) proposed a new biogeographic sector for the Supramontes region (Central-Eastern Sardinia). Other parts of the island, including Gennargentu massif, still remain poorly investigated from a biogeographic viewpoint.

Gennargentu is the main mountain complex of Sardinia and since the beginning of the 18<sup>th</sup> century it became a popular destination for botanical investigations (MORIS, 1827, 1837-1859; BARBEY, 1885; MARTELLI, 1896-1904; HERZOG, 1909; SCHMID, 1933; DESOLE, 1948, 1966; ARRIGONI, 1966, 1986; BRULLO & al., 2001), that led to the description of several new taxa (PIGNATTI & FEOLI, 1974; PIGNATTI & PIGNATTI, 1974; ARRIGONI & al., 1977-1991; PIGNATTI & al., 1980; BACCHETTA

& BRULLO, 2006; BACCHETTA & al., 2000, 2006, 2010) and to the analysis of the conservation status of some threatened taxa (FENU & al., 2011, 2012). In spite of the relevant information provided by the abundant literature, an organic review on the vascular flora of Gennargentu was still missing, as well as floristic checklists for the whole massif or single parts of it. Furthermore, relatively big portions of this territory were little known or even unexplored. For these reasons, up to now, a detailed biogeographic framework has never been proposed for the concerned area.

Aims of this work were a comprehensive checklist of the vascular flora of Gennargentu and the analysis of its endemic component, in order to set the area in the Sardinian biogeographic subprovince.

## Study Area

Gennargentu (Fig. 1) is located in the central-eastern part of Sardinia and borders the “Barbagia di Ollolai” to the North, the “Mandrolisai” to the North-West and West, the “Sarcidano” and “Barbagia di Belvi” to the South, the “Ogliastra” to the East, the “Supramontes of Urzulei and Orgosolo” to the North-East. The study area has a surface of 50,000 ha and consists of a system of summits and windy ridges at 1400-1500 m, with four culminations at more than 1800 m: Punta La Marmora (1834 m), Bruncu Spina (1828 m), Su Sciusciu (1823 m) and Punta Florisa (1822 m). Other important landmarks are Punta Paolinu (1792 m), Monte Spada (1596 m) and Mont’Arbu (1568 m). The heights of Tonneri de Sa Irgini and Arcu Correboi have also been included in the study area (Fig. 1).

Metamorphic rocks are by far the most represented outcrops. They include Carboniferous metasiltstones and metasandstones, regularly superimposed to schists, limestones and dating, respectively, to Devonian-Silurian and Ordovician (CARMIGNANI & al., 2001). As by-products of the late Hercinic orogeny, intrusions of granites and porphyrites are also frequent (CARMIGNANI & al., 2001). In the area of Arcu Correboi, thick Ordovician quartzitic and foliated silicates are interstratified, with Devonian-Silurian black schists and limestones. The study area covers the whole layered structure that identifies the Gennargentu lithostratigraphic unit (CARMIGNANI & al., 2001).

According to the Rivas-Martínez’s bioclimatic classification, most of the Gennargentu massif has a temperate-submediterranean climate, with thermotypes ranging from the lower supratemperate to the lower orotemperate, and ombrotypes from the upper subhumid to the upper humid. The Mediterranean climate is only found on the eastern and southern slopes of the massif, with a lower supramediterranean thermotype and ombrotype ranging from the upper subhumid to the lower humid (BACCHETTA & al., 2009a).




Fig. 1. – Study area in Sardinia.

## Methods

Years of floristic researches have been carried out between 2004 and 2011; field trips were effectuated from February to November. Specimens and seeds collected in the field are stored in CAG and in the Sardinian Germplasm Bank (BG-SAR), respectively. Bibliographic and herbariological researches have been fulfilled in BOLO, CAG, CAT, FI, NAP, PAL, RO, SASSA, SS, TO, W, Z.

Collected specimens have been identified using PIGNATTI (1982), TUTIN & al. (1964-80; 1993), ARRIGONI & al. (1977-1991), JEANMONOD & GAMISANS (2007), BOLÓS & VIGO (1984-2001) and CASTROVIEJO & al. (1986-2011).

The delimitation of the plant families listed in our checklist followed the APG III proposal (APG III, 2009; CHASE & REVEAL, 2009; PERUZZI, 2010). Taxonomic nomenclature followed mainly CONTI & al. (2005, 2007) and GREUTER & al. (1984-2008), JALAS & SUOMINEN (1972-1994), JALAS & al. (1996-1999) and KURTTO & al. (2004).

Growth and life forms have been determined in the field, following the classification of RAUNKIAER (1934), and expressed with the abbreviations proposed by PIGNATTI (1982).

In addition to the consulted floras, chorotypes refer to the classification proposed by BRULLO & al. (1996). For the chorological classification of the endemics, the nomenclature proposed by ARRIGONI & TOMMASO (1991) and modified by BACCHETTA & PONTECORVO (2005) is followed.

For the non-native species, the place of origin is reported, as well as the abbreviations on the “status” proposed by RICHARDSON & al. (2000) and modified by PYŠEK & al. (2004): “Cas = casual”, “Nat = naturalized”, “Inv = invasive”.

Taxa known from literature but not found during our field investigations have not been considered in the floristic analysis.

For the biogeographic analysis of the investigated area, the methodological framework proposed by RIVAS-MARTÍNEZ (2007) has been followed. The same has been recently applied for several areas of Sardinia by BACCHETTA & PONTECORVO (2005), FENU & BACCHETTA (2008), ANGIUS & BACCHETTA (2009), BACCHETTA & al. (2009a) and FENU & al. (2010).

## Results

The vascular flora of Gennargentu consists of 948 taxa, of which 686 are species, 249 subspecies, 10 varieties and 3 hybrids, belonging to 97 families and 427 genera. Among the Angiosperms, dicots (incl. Laurales and Piperales) prevail, with 707 taxa (74.58% of the whole flora); monocots count for 199 taxa (20.99%). Pteridophytes and Gymnosperms include 28 (2.95%) and 14 (1.48%) taxa, respectively.


The most represented families are *Asteraceae* (118 taxa), followed by *Poaceae* (99) and *Fabaceae* (79); significant are also *Caryophyllaceae* (55), *Lamiaceae* (34), *Rosaceae* (31), *Apiaceae* (30), *Brassicaceae* (28) and *Orchidaceae* (26).

The richest genera are *Trifolium* L. (22 taxa), *Ranunculus* L. (18), *Sedum* L., *Vicia* L. and *Carex* L. (13), *Poa* L. (12), *Silene* L. (11), *Euphorbia* L., *Galium* L., *Orchis* L. and *Rumex* L. (10), *Allium* L., *Filago* L., *Quercus* L., *Juncus* L. and *Orobancha* L. (8), *Geranium* L., *Epilobium* L., *Veronica* L., *Vulpia* C. C. Gmel., *Cerastium* L., *Hieracium* L. and *Medicago* L. (7).

The following taxa are new records for the Sardinian and the Italian flora: *Anarrhinum corsicum* Jord. & Fourr., *Hypericum corsicum* Steud. and *Sagina procumbens* subsp. *muscosa* (Jord.) Nyman; the following ones are new for the flora of the Island: *Asplenium adiantum-nigrum* L. subsp. *adiantum-nigrum*, *A. trichomanes* subsp. *pachyrachis* (Christ) Lovis & Reichst., *Carex depauperata* Curtis & With., *C. flava* L., *Rubus caesius* L., *Rumex aquaticus* L., *Senecio erucifolius* L., and *Tragopogon dubius* Scop.

Life form analysis (Fig. 2) highlighted a clear prevalence of hemicryptophytes (35.65%), followed by therophytes (34.6%), geophytes (12.13%) and (nano-)phanerophytes (11.6%).

As concerns chorology (Fig. 3), the flora of Gennargentu is mostly constituted by Mediterranean taxa (68.14%); the only other significant chorotype is the palaeotemperate (9.07%), the rest ranging between 0.21 and 7.28%. In the Mediterranean quota, the circum-Mediterranean elements prevail (188 taxa, 29.1%), followed by the Euro-Medit. (149 taxa,



**Fig. 2.** – Life forms percentages, referred to the whole flora. H = hemicryptophytes; C = chamaephytes; G = geophytes; NP = nanophanerophytes; P = phanerophytes; T = therophytes; Hy = hydrophytes.

Table 1. – List of endemic taxa.

No.	Taxon	Family	Biological form	Chorology	Biogeographic unit
1	<i>Allium parviflorum</i> Viv.	Amaryllidaceae	G	SA-CO	Sardo-Cors. Prov.
2	<i>Allium roseum</i> var. <i>insulare</i> Gennari	Amaryllidaceae	G	SA-CO	Sardo-Cors. Prov.
3	<i>Anarrhinum corsicum</i> Jord. & Fourr.	Plantaginaceae	H	SA-CO	Sardo-Cors. Prov.
4	<i>Aquilegia barbaricina</i> Arrigoni & E. Nardi	Ranunculaceae	G	SA	Sard. Subprov.
5	<i>Aquilegia nugorensis</i> Arrigoni & E. Nardi	Ranunculaceae	G	SA	Sard. Subprov.
6	<i>Arenaria balearica</i> L.	Caryophyllaceae	Ch	SA-CO-AT-BL	W-Medit. Subregion
7	<i>Aristolochia rotunda</i> subsp. <i>insularis</i> E. Nardi & Arrigoni	Aristolochiaceae	G	SA-CO-AT	Ital.-Tyrr. Superprov.
8	<i>Aristolochia tyrrhena</i> E. Nardi & Arrigoni	Aristolochiaceae	G	SA-CO-AT	Ital.-Tyrr. Superprov.
9	<i>Armeria morisii</i> Boiss.	Plumbaginaceae	H	SA	Sard. Subprov.
10	<i>Armeria sardoa</i> subsp. <i>genargentea</i> Arrigoni	Plumbaginaceae	H	SA	Gennargentu Sector
11	<i>Armeria sardoa</i> Spreng. subsp. <i>sardoa</i>	Plumbaginaceae	H	SA	Sard. Subprov.
12	<i>Arum pictum</i> L. f. subsp. <i>pictum</i>	Araceae	G	SA-CO	Sardo-Cors. Prov.
13	<i>Astragalus genargenteus</i> Moris	Fabaceae	Ch	SA	Sard. Subprov.
14	<i>Barbarea rupicola</i> Moris	Brassicaceae	Ch	SA-CO	Sardo-Cors. Prov.
15	<i>Bellium bellidioides</i> L.	Asteraceae	H	SA-CO-BL	W-Medit. Subregion
16	<i>Berberis vulgaris</i> subsp. <i>aetnensis</i> (C. Presl) Rouy & Foucaud	Berberidaceae	NP	SA-CO-SHTM	Ital.-Tyrr. Superprov.
17	<i>Bituminaria morisiana</i> (Pignatti & Metlesics) Greuter	Fabaceae	Ch	SA-TN (La Galite)	W-Medit. Subregion
18	<i>Borago pygmaea</i> (DC.) Chater & Greuter	Boraginaceae	H	SA-CO-AT	Ital.-Tyrr. Superprov.
19	<i>Brimeura fastigiata</i> (Viv.) Chouard	Asparagaceae	G	SA-CO-BL	W-Medit. Subregion
20	<i>Bunium corydalinum</i> DC. subsp. <i>corydalinum</i>	Apiaceae	G	SA-CO	Sardo-Cors. Prov.
21	<i>Campanula forsythii</i> (Arcang.) Bég.	Campanulaceae	H	SA	Sard. Subprov.
22	<i>Carduus sardous</i> DC.	Asteraceae	H	SA-CO-AT-ITC	Ital.-Tyrr. Superprov.
23	<i>Carex caryophyllea</i> subsp. <i>insularis</i> (Barbey) Arrigoni	Cyperaceae	H	SA-CO-(SI)	Ital.-Tyrr. Superprov.
24	<i>Carex microcarpa</i> Moris	Cyperaceae	G	SA-CO-AT	Ital.-Tyrr. Superprov.
25	<i>Carlina macrocephala</i> Moris subsp. <i>macrocephala</i>	Asteraceae	H	SA-CO	Sardo-Cors. Prov.
26	<i>Centaurea magistrorum</i> Arrigoni & Camarda	Asteraceae	Ch	SA	Gennargentu Sector
27	<i>Cephalaria mediterranea</i> (Viv.) Szabó	Caprifoliaceae	Ch	SA	Sard. Subprov.
28	<i>Cerastium palustre</i> Moris	Caryophyllaceae	T	SA	Sard. Subprov.
29	<i>Cistus creticus</i> subsp. <i>corsicus</i> (Loisel.) Greuter	Cistaceae	NP	SA-CO	Sardo-Cors. Prov.
30	<i>Clinopodium sardoum</i> (Asch. & Levier) Peruzzi & F. Conti	Lamiaceae	Ch	SA	Sard. Subprov.
31	<i>Clypeola jonthlaspi</i> subsp. <i>microcarpa</i> (Moris) Arcang.	Brassicaceae	T	SA-CO-SI	Ital.-Tyrr. Superprov.
32	<i>Colchicum alpinum</i> subsp. <i>parvulum</i> (Ten.) Nyman	Colchicaceae	G	SA-CO-SHTC	Ital.-Tyrr. Superprov.
33	<i>Colchicum gonarei</i> Camarda	Colchicaceae	G	SA	Sard. Subprov.
34	<i>Crepis vesicaria</i> subsp. <i>hyemalis</i> (Biv.) Babç.	Asteraceae	H	SA-SI	Ital.-Tyrr. Superprov.
35	<i>Crocus minimus</i> DC.	Iridaceae	G	SA-CO-AT	Ital.-Tyrr. Superprov.
36	<i>Cuscuta epithimum</i> subsp. <i>corsicana</i> (Yunck.) Lambinon	Convolvulaceae	T	SA-CO	Sardo-Cors. Prov.
37	<i>Cymbalaria aequitriloba</i> (Viv.) A. Chev. subsp. <i>aequitriloba</i>	Plantaginaceae	Ch	SA-CO-AT-BL	W-Medit. Subregion
38	<i>Cymbalaria muelleri</i> (Moris) A. Chev.	Plantaginaceae	Ch	SA	Sard. Subprov.
39	<i>Cynoglossum barbaricinum</i> Arrigoni & Selvi	Boraginaceae	G	SA	Gennargentu Sector
40	<i>Delphinium pictum</i> Willd. subsp. <i>pictum</i>	Ranunculaceae	H	SA-CO-BL-H	W-Medit. Subregion
41	<i>Dianthus genargenteus</i> Bacch., Brullo, Casti & Giusso	Caryophyllaceae	H	SA	Gennargentu Sector
42	<i>Dianthus oliastreae</i> Bacch., Brullo, Casti & Giusso	Caryophyllaceae	H	SA	Sard. Subprov.
43	<i>Dianthus sardous</i> Bacch., Brullo, Casti & Giusso	Caryophyllaceae	H	SA	Sard. Subprov.
44	<i>Digitalis purpurea</i> var. <i>gyspergerae</i> (Rouy) Fiori	Plantaginaceae	H	SA-CO	Sardo-Cors. Prov.
45	<i>Dipsacus ferox</i> Loisel.	Caprifoliaceae	H	SA-CO	Sardo-Cors. Prov.
46	<i>Echium anchusoides</i> Bacch., Brullo & Selvi	Boraginaceae	H	SA	Sard. Subprov.
47	<i>Euphorbia amygdaloides</i> subsp. <i>semiperfoliata</i> (Viv.) Radcl.-Sm.	Euphorbiaceae	H	SA-CO	Sardo-Cors. Prov.
48	<i>Euphorbia gayi</i> Salis	Euphorbiaceae	G	SA-CO-BL	W-Medit. Subregion
49	<i>Euphorbia hyberna</i> subsp. <i>insularis</i> (Boiss.) Briq.	Euphorbiaceae	G	SA-CO-ITC (Liguria e Toscana)	Ital.-Tyrr. Superprov.

Cont.

Table 1. – *Cont.*

No.	Taxon	Family	Biological form	Chorology	Biogeographic unit
50	<i>Euphorbia meuselii</i> Geltman	Euphorbiaceae	Ch	SA-SHTM (Calabria)	Ital.-Tyrr. Superprov.
51	<i>Euphrasia nana</i> Rouy	Orobanchaceae	T	SA-CO	Sardo-Cors. Prov.
52	<i>Festuca morisiana</i> Parl.	Poaceae	H	SA	Sard. Subprov.
53	<i>Galium corsicum</i> Spreng.	Rubiaceae	H	SA-CO	Sardo-Cors. Prov.
54	<i>Galium schmidii</i> Arrigoni	Rubiaceae	Ch	SA	Sard. Subprov.
55	<i>Genista aetnensis</i> (Biv.) DC.	Fabaceae	P	SA-CO-SI	Ital.-Tyrr. Superprov.
56	<i>Genista corsica</i> (Loisel.) DC.	Fabaceae	NP	SA-CO	Sardo-Cors. Prov.
57	<i>Genista desoleana</i> Vals.	Fabaceae	NP	SA-CO-ITC (Liguria e Toscana)	Ital.-Tyrr. Superprov.
58	<i>Genista pichisermoliana</i> Vals.	Fabaceae	NP	SA	Gennargentu Sector
59	<i>Glechoma sardoa</i> (Bég.) Bég.	Lamiaceae	H	SA	Sard. Subprov.
60	<i>Helichrysum microphyllum</i> subsp. <i>tyrrhenicum</i> Bacch., Brullo & Giusso	Asteraceae	Ch	SA-CO-BL	W-Medit. Subregion
61	<i>Helichrysum saxatile</i> Moris subsp. <i>saxatile</i>	Asteraceae	Ch	SA	Sard. Subprov.
62	<i>Helleborus lividus</i> subsp. <i>corsicus</i> (Briq.) P. F. Yeo	Ranunculaceae	G	SA-CO	Sardo-Cors. Prov.
63	<i>Herniaria litardierei</i> (Gamisans) Greuter & Burdet	Caryophyllaceae	H	SA-CO	Sardo-Cors. Prov.
64	<i>Hieracium iolai</i> Arrigoni	Asteraceae	H	SA	Sard. Subprov.
65	<i>Hieracium soleirolianum</i> Arv.-Touv. & Briq.	Asteraceae	H	SA-CO	Sardo-Cors. Prov.
66	<i>Hieracium zizianum</i> subsp. <i>sardonium</i> Zahn	Asteraceae	H	SA-CO	Sardo-Cors. Prov.
67	<i>Hypericum annulatum</i> Moris	Hypericaceae	H	SA	Sard. Subprov.
68	<i>Hypericum corsicum</i> Steud.	Hypericaceae	H	SA-CO	Sardo-Cors. Prov.
69	<i>Hypericum hircinum</i> L. subsp. <i>hircinum</i>	Hypericaceae	NP	SA-CO-AT	Ital.-Tyrr. Superprov.
70	<i>Hypochaeris sardoa</i> Bacch., Brullo & Terrasi	Asteraceae	H	SA-CO	Sardo-Cors. Prov.
71	<i>Iberis integerrima</i> Moris	Brassicaceae	Ch	SA	Sard. Subprov.
72	<i>Juniperus communis</i> var. <i>corsicana</i> Lebreton, Mossa & Gallet	Cupressaceae	NP	SA-CO	Sardo-Cors. Prov.
73	<i>Lamium garganicum</i> subsp. <i>corsicum</i> (Gren. & Godr.) Mennema	Lamiaceae	H	SA-CO	Sardo-Cors. Prov.
74	<i>Lamyropsis microcephala</i> (Moris) Dittrich & Greuter	Asteraceae	G	SA	Gennargentu Sector
75	<i>Limonium morisianum</i> Arrigoni	Plumbaginaceae	Ch	SA	Sard. Subprov.
76	<i>Lonicera cyrenaica</i> Viv.	Caprifoliaceae	NP	SA-LI	W-Medit. Subregion
77	<i>Luzula spicata</i> subsp. <i>italica</i> (Parl.) Arcang.	Juncaceae	H	SA-CO	Sardo-Cors. Prov.
78	<i>Mentha requienii</i> Benth. subsp. <i>requienii</i>	Lamiaceae	H	SA-CO	Sardo-Cors. Prov.
79	<i>Mentha suaveolens</i> subsp. <i>insularis</i> (Req.) Greuter	Lamiaceae	H	SA-CO-AT-BL	W-Medit. Subregion
80	<i>Mercurialis corsica</i> Coss. & Kralik	Urticaceae	Ch	SA-CO	Sardo-Cors. Prov.
81	<i>Micromeria filiformis</i> subsp. <i>cordata</i> (Bertol.) Pignatti	Lamiaceae	Ch	SA	Sard. Subprov.
82	<i>Morisia monanthos</i> (Viv.) Asch.	Brassicaceae	H	SA-CO	Sardo-Cors. Prov.
83	<i>Myosotis soleirolii</i> Godr.	Boraginaceae	H	SA-CO	Sardo-Cors. Prov.
84	<i>Noccaea brevistyla</i> (DC.) Steud.	Brassicaceae	H	SA-CO	Sardo-Cors. Prov.
85	<i>Odontites corsicus</i> (Loisel.) G. Don	Orobanchaceae	T	SA-CO	Sardo-Cors. Prov.
86	<i>Oenanthe lisae</i> Moris	Apiaceae	H	SA	Sard. Subprov.
87	<i>Orchis mascula</i> subsp. <i>ichnusae</i> Corrias	Orchidaceae	G	SA-CO	Sardo-Cors. Prov.
88	<i>Ornithogalum corsicum</i> Jord. & Fourr.	Asparagaceae	G	SA-CO	Sardo-Cors. Prov.
89	<i>Orobanche denudata</i> Moris	Orobanchaceae	G	SA	Gennargentu Sector
90	<i>Orobanche rapum-genistae</i> subsp. <i>rigens</i> (Loisel.) Arcang.	Orobanchaceae	G	SA-CO	Sardo-Cors. Prov.
91	<i>Paeonia corsica</i> Tausch	Paeoniaceae	G	SA-CO	Sardo-Cors. Prov.
92	<i>Pancratium illyricum</i> L.	Amaryllidaceae	G	SA-CO-AT	Ital.-Tyrr. Superprov.
93	<i>Petrohragia saxifraga</i> subsp. <i>bicolor</i> (Jord. & Fourr.) Gamisans	Caryophyllaceae	H	SA-CO	Sardo-Cors. Prov.
94	<i>Phalaroides arundinacea</i> subsp. <i>rotgesii</i> (Husn.) Valdés & H. Scholz	Poaceae	H	SA-CO	Sardo-Cors. Prov.
95	<i>Plantago subulata</i> subsp. <i>insularis</i> (Gren. & Godr.) Nyman	Plantaginaceae	Ch	SA-CO	Sardo-Cors. Prov.
96	<i>Poa balbisii</i> Parl.	Poaceae	H	SA-CO	Sardo-Cors. Prov.
97	<i>Polygala sardoa</i> Chodat	Polygalaceae	H	SA	Sard. Subprov.
98	<i>Potentilla caulescens</i> subsp. <i>nebrodensis</i> (Zimm.) Arrigoni	Rosaceae	Ch	SA-SI	Ital.-Tyrr. Superprov.

*Cont.*

Table 1. – Cont.

No.	Taxon	Family	Biological form	Chorology	Biogeographic unit
99	<i>Potentilla crassinervia</i> Viv.	Rosaceae	Ch	SA-CO	Sardo-Cors. Prov.
100	<i>Potentilla rupestris</i> subsp. <i>corsica</i> (Lehm.) Rouy & E. G. Camus	Rosaceae	H	SA-CO	Sardo-Cors. Prov.
101	<i>Prospero autumnale</i> var. <i>corsicana</i> (Boullu) Briq.	Asparagaceae	G	SA-CO	Sardo-Cors. Prov.
102	<i>Phlostemon casabonae</i> (L.) Greuter	Asteraceae	H	SA-CO-H-AT	W-Medit. Subregion
103	<i>Ptychotis sardoa</i> Pignatti & Metlesics	Apiaceae	H	SA	Sard. Subprov.
104	<i>Pulicaria vulgaris</i> var. <i>sardoa</i> Fiori	Asteraceae	T	SA	Sard. Subprov.
105	<i>Quercus ichnusae</i> Mossa, Bacch. & Brullo	Fagaceae	P	SA	Sard. Subprov.
106	<i>Ranunculus cordiger</i> Viv. subsp. <i>cordiger</i>	Ranunculaceae	H	SA-CO	Sard. Subprov.
107	<i>Ranunculus cymbalarifolius</i> Moris	Ranunculaceae	H	SA	Sard. Subprov.
108	<i>Rhamnus persicifolia</i> Moris	Rhamnaceae	P	SA	Sard. Subprov.
109	<i>Ribes multiflorum</i> subsp. <i>sandaloticum</i> Arrigoni	Grossulariaceae	NP	SA	Sard. Subprov.
110	<i>Romulea requienii</i> Parl.	Iridaceae	G	SA-CO	Sardo-Cors. Prov.
111	<i>Rumex pulcher</i> subsp. <i>suffocatus</i> (Bertol.) Nyman	Polygonaceae	H	SA	Sard. Subprov.
112	<i>Rumex scutatus</i> subsp. <i>glaucescens</i> (Guss.) Brullo, Scelsi & Spamp.	Polygonaceae	H	SA-SHTM (Calabria)	Ital.-Tyrr. Superprov.
113	<i>Ruta lamarmorae</i> Bacch., Brullo & Giusso	Rutaceae	Ch	SA	Gennargentu Sector
114	<i>Sagina pilifera</i> (DC.) Fenzl	Caryophyllaceae	H	SA	Sardo-Cors. Prov.
115	<i>Salix arrigonii</i> Brullo	Salicaceae	P	SA	Sard. Subprov.
116	<i>Salix purpurea</i> subsp. <i>eburnea</i> (Borzì) Cif. & Giacom.	Salicaceae	P	SA	Sard. Subprov.
117	<i>Santolina insularis</i> (Fiori) Arrigoni	Asteraceae	NP	SA	Sard. Subprov.
118	<i>Saponaria ocymoides</i> subsp. <i>alsinoides</i> (Viv.) Arcang.	Caryophyllaceae	H	SA-CO	Sardo-Cors. Prov.
119	<i>Saxifraga corsica</i> Gren. & Godr. subsp. <i>corsica</i>	Saxifragaceae	H	SA-CO	Sardo-Cors. Prov.
120	<i>Saxifraga pedemontana</i> subsp. <i>cervicornis</i> (Viv.) Engl.	Saxifragaceae	Ch	SA-CO	Sardo-Cors. Prov.
121	<i>Scorzonera callosa</i> Moris	Asteraceae	H	SA	Sard. Subprov.
122	<i>Scrophularia oblongifolia</i> Loisel. subsp. <i>oblongifolia</i>	Scrophulariaceae	H	SA-CO	Sardo-Cors. Prov.
123	<i>Scrophularia trifoliata</i> L.	Scrophulariaceae	H	SA-CO-AT	Ital.-Tyrr. Superprov.
124	<i>Sedum villosum</i> subsp. <i>glandulosum</i> Moris	Crassulaceae	T	SA	Gennargentu Sector
125	<i>Sesleria insularis</i> subsp. <i>barbaricina</i> Arrigoni	Poaceae	H	SA	Sard. Subprov.
126	<i>Silene nodulosa</i> Viv.	Caryophyllaceae	H	SA-CO	Sardo-Cors. Prov.
127	<i>Silene requienii</i> Otth	Caryophyllaceae	Ch	SA-CO	Sardo-Cors. Prov.
128	<i>Sorbus aucuparia</i> subsp. <i>praemorsa</i> (Guss.) Nyman	Rosaceae	P	SA-CO-SHTM	Ital.-Tyrr. Superprov.
129	<i>Stachys corsica</i> Pers.	Lamiaceae	H	SA-CO	Ital.-Tyrr. Superprov.
130	<i>Stachys glutinosa</i> L.	Lamiaceae	Ch	SA-CO-AT	Ital.-Tyrr. Superprov.
131	<i>Tanacetum audibertii</i> (Req.) DC.	Asteraceae	H	SA-CO	Sardo-Cors. Prov.
132	<i>Taraxacum garbarianum</i> Peruzzi, Aquaro, Caparelli & Raimondo	Asteraceae	H	SA-SI	Ital.-Tyrr. Superprov.
133	<i>Teucrium marum</i> L.	Lamiaceae	Ch	SA-CO-BL-AT-H	W-Medit. Subregion
134	<i>Thesium italicum</i> A. DC.	Santalaceae	H	SA	Sard. Subprov.
135	<i>Thymus herba-barona</i> Loisel. subsp. <i>herba-barona</i>	Lamiaceae	Ch	SA-CO	Sardo-Cors. Prov.
136	<i>Trisetum gracile</i> (Moris) Boiss.	Poaceae	H	SA-CO	Sardo-Cors. Prov.
137	<i>Urtica atrovirens</i> Loisel.	Urticaceae	H	SA-CO-AT	Ital.-Tyrr. Superprov.
138	<i>Verbascum conocarpum</i> Moris subsp. <i>conocarpum</i>	Scrophulariaceae	H	SA-CO-AT	Ital.-Tyrr. Superprov.
139	<i>Veronica verna</i> subsp. <i>brevistyla</i> (Moris) Rouy	Plantaginaceae	T	SA-CO	Sardo-Cors. Prov.
140	<i>Vinca difformis</i> subsp. <i>sardoa</i> Stearn	Apocynaceae	Ch	SA	Sard. Subprov.
141	<i>Viola corsica</i> subsp. <i>limbarae</i> Merxm. & W. Lippert	Violaceae	H	SA	Sard. Subprov.






Fig. 3. – Percentages of the chorologic units, referred to the whole flora.




Fig. 4. – Life forms percentages of the endemic flora of Gennargentu. H = hemicyptophytes; C = chamaephytes; G = geophytes; NP = nanophanerophytes; P = phanerophytes; T = therophytes.

23.07%) and the endemic (141 taxa, 21.83%). The West-Medit. (56 taxa, 8.67%), together with the Medit.-Atlantic (29 taxa, 4.49%) count for 13.16% of the whole flora.

In our census, only 38 non-native taxa have been found, i.e. 4.01% of the whole flora: 1 of them is considered invasive (*Xanthium spinosum* L.), 19 casual and 18 naturalized.

The endemic component (Table 1) includes 141 taxa: 84 species, 51 subspecies and 6 varieties, belonging to 46 families and 108 genera. The most represented families are *Asteraceae* (18 taxa), *Caryophyllaceae* (11), *Lamiaceae* (10) *Ranunculaceae*, *Plantaginaceae* and *Fabaceae* (6). The genera richest in endemic taxa are: *Genista* L. and *Euphorbia* L. (4), *Armeria* Willd., *Dianthus* L., *Hieracium* L., *Hypericum* L., *Potentilla* L. (3), followed by 17 genera with 2 taxa each, such as for example *Aquilegia* L., *Ranunculus* L., *Saxifraga* L., *Carex* L. and *Scrophularia* L.

Nearly the half of the endemics (Fig. 4) are hemicyptophytes (46.1%), the rest are chamaephytes (19.15%), geophytes (17.02%) and (nano-)phanerophytes (11.35%).

The Sardo-Corsican elements (Fig. 5) prevail (39.01%), followed by the Sardinian ones (35.46%) and by the Italo-Tyrrhenian (18.44%). The last group includes taxa in common with the Tuscan Archipelago (7.8%) and some others ranging up to limited portions of the western coast of the Italian peninsula and Sicily (10.64%).




Fig. 5. – Percentages of the chorologic units of the endemic flora of Gennargentu. EMOI = W-Mediterranean insular endemics; ETI = Tyrrhenian insular endemics; ET = Tyrrhenian endemics; ETI-NA = Tyrrhenian insular and N-Africa endemics; ESS = Sardinia and Sicily endemics; ESC = Sardo-Corsican endemics; ESA = Sardinian endemics.

## Discussion

In the area of Gennargentu, corresponding roughly to 1% of Sardinia, more than 30% of the regional flora is occurring. Although the available floristic data do not allow to estimate species-area relationships, this high number of taxa, combined with more general considerations on the uniqueness of Gennargentu in terms of largeness and altitude highlight the floristic importance of this area.

The floristic richness of Gennargentu can be explained hypothesizing that along the Quaternary age, the less drastic climate changes on large Mediterranean islands favoured the local persistence of high plant richness and the co-existence of distinct genetic lineages (VALIENTE BANUET & al., 2006; MÈDAIL & DIADEMA, 2009), confirming the identification of this area as one of the Mediterranean putative refugia. The observed pattern in plant family distribution, which is only partially correlated with the pattern of endemic species, may testify that Gennargentu acted as a “climatic island” during the Quaternary climatic variations. Floristic changes were more severe and drastic in the lowlands and re-colonisation was particularly important for shaping the modern floras at lower altitudes. Not all taxa were able to recolonise and the empty niches were filled by new adaptive radiation of migrating taxa, with a strong tendency towards the annual life strategy (GUARINO, 2006). The result is that families are generally larger, but with fewer endemic species in the lowlands than in the summit areas of Gennargentu, where the elevation facilitated the survival or, eventually, an independent adaptive radiation of locally surviving taxa. For example, a high potential for adaptive radiation of the *Ranunculaceae* may be seen in the fact that this family has a significant number of endemic species on Gennargentu, even if it does not account for the most representative families in that area, nor in the whole Sardinian flora. Likewise, the *Asteraceae* seem to have a strong potential for both migrations and local adaptive radiation, whereas the *Poaceae* are remarkable for a high migration potential but may have a low potential for local, independent adaptive radiation. Families like *Fabaceae*, *Asteraceae* and *Euphorbiaceae* were likely to be able to diversify on the Gennargentu summits not only in the Pleistocene, but also in the Post-Messinian phase.

The role of Gennargentu as a “climatic island” was also confirmed by the life form analysis where the ratio between hemicyptophytes and therophytes is much higher than in the rest of Sardinia (28.1% and 39.9% for H and T, respectively; BOCCHIERI, 1995). Even the number of nano-phanerophytes is well above the average value for Sardinia (8.8%; BOCCHIERI, 1995). The percentage of geophytes is also pretty high; this is probably linked to the pastoral land-use of the whole area, frequently affected by periodical fires as for the whole Island (12.1%; BOCCHIERI, 1995). Chamaephytes display values

slightly under the average value for Sardinia (8.1%; BOCCHIERI, 1995) and the few hydrophytes testify the lack of backwater on Gennargentu.

In the chorologic analysis, the relatively high percentage of palaeotemperate elements outlines the temperate-sub-mediterranean bioclimate in the summit areas of the massif, as well as on its northern slopes. The low percentage of non-native taxa denotes the high naturalistic value of the inspected area and suggests that the mountain flora of Sardinia is probably less prone to competition by allochthonous taxa (BACCHETTA & al., 2009b).

Sardinia is well known for its richness in endemic taxa, due to the isolation and high topographic diversity (MÈDAIL & QUÉZEL, 1997). The Gennargentu massif, even if its elevation is not comparable with that of the Corsican mountains and has never been affected by glacial perturbation, hosts several distinctive ecological niches as highlighted in the case of *Lamyropsis microcephala* (Moris) Dittrich & Greuter by MATTANA & al. (2009). The ecologic and bioclimatic isolation, added to the geographic insularity of Sardinia lets to identify the Gennargentu massif as one of the main “micro hotspot” of Sardinia sensu FENU & al. (2010).

Moreover, like the Supramontes region (FENU & al., 2010), Gennargentu represents a southern European “refugium” (sensu TZEDAKIS & al., 2002) for some temperate tree species (e.g. *Acer monspessulanum* L. subsp. *monspessulanum*, *Ilex aquifolium* L., *Ostrya carpinifolia* Scop., *Quercus congesta* C. Presl, *Rhamnus alpina* L. subsp. *alpina*, *Sambucus nigra* L., *Sorbus aria* (L.) Crantz subsp. *aria*, *S. aucuparia* subsp. *praemorsa* (Guss.) Nyman, *S. torminalis* Crantz, *Taxus baccata* L.). Therefore, it represents an area of special value for the long-term persistence of biodiversity (TABERLET & CHEDDADI, 2002), as further testified by the occurrence of several relict endemic taxa, like *Astragalus genargentus* Moris, *Lamyropsis microcephala*, *Rhamnus persicifolia* Moris, *Ribes multiflorum* subsp. *sandalioticum* Arrigoni, *Ruta lamarmorae* Bacch., Brullo & Giusso, *Tanacetum audibertii* (Req.) DC.

In particular, rocky habitats and windy summit areas, in spite of their limited extension, form a very important reservoir for the local biodiversity (MÈDAIL & QUÉZEL, 1997). The high percentage of Sardinian endemics (35.46%) and, within these, of the taxa exclusively growing on Gennargentu (6.38%) testifies the floristic autonomy of the massif. The relatively high percentage of Sardo-Corsican endemics (39.01%), as well as of endemic taxa in common with the Tuscan Archipelago (7.8%) turns out to be so high because of the prevalence of siliceous rocks, that enhances the floristic affinity of Gennargentu with the Corsican and the Ilvensian territories (BACCHETTA & PONTECORVO, 2005). On the other hand, in the Sardinian areas where limestones and carbonatic rocks in general prevail, and particularly in the so-called

“carbonate-metalliferous ring” (i.e. S-W Sardinia), or in the Supramontes area (i.e. C-E Sardinia, just around Gennargentu), the number of Sardinian exclusive endemics tends to be higher than in Gennargentu (BACCHETTA & PONTECORVO, 2005; FENU & al., 2010).

The nine endemic taxa in common with the Balearic Islands, added to the 8.51% of West-Mediterranean elements, confirmed the biogeographical affiliation of Sardinia to the so-called “W-Mediterranean subregion” (BACCHETTA & PONTECORVO, 2005). The 11 taxa in common with Corsica and the Tuscan Archipelago justify the biogeographical identity of an Italo-Tyrrhenian Superprovince, as proposed by LADERO ALVAREZ & al. (1987) and followed by BACCHETTA & al. (2012a).

Due to the relatively high number of exclusive endemics of Gennargentu massif, and the geologic and geomorphologic peculiarities, it is here proposed a biogeographic classification for these territories with the identification of an autonomous sector named “Gennargentu”. According to RÍOS RUIZ & al. (2003), the floristic autonomy of this sector is highlighted not only by the presence of nine exclusive taxa (*Armeria sardoa* subsp. *genargentea* Arrigoni, *Centaurea magistrorum* Arrigoni & Camarda, *Cynoglossum barbaricinum* Arrigoni & Selvi, *Dianthus genargenteus* Bacch., Brullo, Casti & Giusso, *Genista pichisermolliana* Vals., *Lamyropsis microcephala*, *Orobanche denudata* Moris, *Sedum villosum* subsp. *glandulosum* Moris, *Ruta lamarmorae*) but also by several differential taxa, with a wider distribution range and limited in Sardinia to the study area, such as *Anarrhinum corsicum*, *Euphorbia hyberna* subsp. *insularis* (Boiss.) Briq., *Euphrasia nana* Rouy, *Herniaria litardierei* (Gamisans) Greuter & Burdet, *Hypericum corsicum*, *Gentiana lutea* L. s.l., *Myosotis soleirolii* Godr., *Noccaea brevistyla* (DC.) Steud., *Sorbus aucuparia* subsp. *praemorsa* (Guss.) Nyman, *Tanacetum audibertii*, *Trisetum gracile* (Moris) Boiss. In addition, the Gennargentu massif shares with the surrounding sector of Supramontes some relictual endemic calcifugous taxa like *Genista aetnensis* (Biv.) DC., *Morisia monanthos* (Viv.) Asch., *Cerastium palustre* Moris, *Astragalus genargenteus*, *Juniperus nana* var. *corsicana* Lebreton, Mossa & Gallet and *Carex microcarpa* Moris (FENU & al., 2010).

In conclusion, the overall floristic richness of Gennargentu massif, determined by its ecological insularity and the richness of its endemic flora, highlighted the peculiarity of this territory and its identification as one of the Mediterranean putative “refugia”, as stated by MEDAIL & DIADEMA (2009), justifying the set up of an autonomous biogeographic sector.

## References

- ANGIUS, R. & G. BACCHETTA (2009). Boschi e boscaglie ripariali del Sulcis-Iglesiente (Sardegna Sud-Occidentale). *Braun-Blanquetia* 45.
- APG III (2009). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Bot. J. Linn. Soc.* 161: 105-121.
- ARRIGONI, P. V. (1966). Il Governo dell'istituendo Parco Nazionale del Gennargentu, in Sardegna, in rapporto ai caratteri geobotanici del territorio. *Arch. Bot. Biogeogr. Ital.* 42: 218-233.
- ARRIGONI, P. V. (1986). Contributo alla conoscenza della vegetazione del Monte Gennargentu, in Sardegna. *Boll. Soc. Sarda Sci. Nat.* 25: 63-96.
- ARRIGONI, P. V., I. CAMARDA, B. CORRIAS, S. DIANA CORRIAS, E. NARDI, M. RAFFAELLI & F. VALSECCHI (1977-1991). Le piante endemiche della Sardegna. 1-202. *Boll. Soc. Sarda Sci. Nat.* 16-28.
- ARRIGONI, P. V. & P. L. DI TOMMASO (1991). La vegetazione delle montagne calcaree della Sardegna centro-orientale. *Boll. Soc. Sarda Sci. Nat.* 28: 201-310.
- BACCHETTA, G., S. BAGELLA, E. BIONDI, E. FARRIS, R. FILIGHEDDU & L. MOSSA (2009a). Vegetazione forestale e serie di vegetazione della Sardegna (con rappresentazione cartografica alla scala 1:350.000). *Fitosociologia* 46(1).
- BACCHETTA, G. & S. BRULLO (2006). Taxonomic revision of *Astragalus genargenteus* Moris species complex (Fabaceae). *Willdenowia* 36: 157-167.
- BACCHETTA, G., S. BRULLO, M. CASTI & G. P. GIUSSO DEL GALDO (2010). Taxonomic revision of the *Dianthus sylvestris* group (Caryophyllaceae) in central-southern Italy, Sicily and Sardinia. *Nordic J. Bot.* 28: 137-173.
- BACCHETTA, G., S. BRULLO & G. GIUSSO DEL GALDO (2006). *Ruta lamarmorae* a new species from Sardinia. *Edinburgh J. Bot.* 63: 153-160.
- BACCHETTA, G., S. BRULLO & F. SELVI (2000). *Echium anchusoides* (Boraginaceae), a new species from Sardinia (Italy). *Nordic J. Bot.* 20: 271-278.
- BACCHETTA, G., E. FARRIS & C. PONTECORVO (2012a). A new method to set conservation priorities in biodiversity hotspots. *Pl. Biosyst.* 146: 638-648.
- BACCHETTA, G., G. FENU & E. MATTANA (2012b). A checklist of the exclusive vascular flora of Sardinia with priority rankings for conservation. *Anales Jard. Bot. Madrid* 69: 81-89.
- BACCHETTA, G., O. MAYORAL GARCÍA-BERLANGA & L. PODDA (2009b). Catálogo de la flora exótica de Cerdeña (Italia). *Fl. Montiber.* 41: 35-61.
- BACCHETTA, G. & C. PONTECORVO (2005). Contribution to the knowledge of the endemic vascular flora of Iglesias (SW Sardinia-Italy). *Candollea* 60: 481-501.
- BARBEY, W. (1885). *Flora Sardoae compendium*. Lausanne.
- BOCCHIERI, E. (1995). La connaissance et l'état de conservation de la flore en Sardaigne. *Ecol. Medit.* 21: 71-81.

- BOLÒS, O. & J. VIGO (1984-2001). *Fl. Països Catalans* 1-4. Editorial Barcino.
- BRULLO, S., G. GIUSSO DEL GALDO & R. GUARINO (2001). The orophilous communities of the Pino-Juniperetea class in the Central and Eastern Mediterranean area. *Feddes Repert.* 112: 261-308.
- BRULLO, S., M. GRILLO & A. GUGLIELMO (1996). Considerazioni fitogeografiche sulla flora iblea. *Boll. Acc. Gioenia Sci. Nat.* 29(352): 45-111.
- CARMIGNANI, L., G. OGGIANO, S. BARCA, P. CONTI, A. ELTRUDIS, A. FUNEDDA & S. PASCÌ (2001). *Note illustrative della Carta Geologica della Sardegna in scala 1:200.000*. – Memorie descrittive della Carta Geologica d'Italia. Istituto Poligrafico e Zecca dello Stato, Roma.
- CASTROVIEJO, S. (ed.) (1986-2011). *Fl. Iber.* Real Jardín Botánico, C.S.I.C., Madrid.
- CHASE, M.W. & J. L. REVEAL (2009) A phylogenetic classification of the land plants to accompany APG III. *Bot. J. Linn. Soc.* 161: 122-127.
- CONTI, F., G. ABBATE, A. ALESSANDRINI & C. BLASI (ed.) (2005). *An annotated checklist of the Italian Vascular Flora*. Palombi Editori, Roma.
- CONTI, F., A. ALESSANDRINI, G. BACCHETTA, E. BANFI, G. BARBERIS, F. BARTOLUCCI, L. BERNARDO, S. BONACQUISTI, D. BOUVET, M. BOVIO, G. BRUSA, E. DEL GUACCHIO, B. FOGGI, S. FRATTINI, G. GALASSO, L. GALLO, C. GANGALE, G. GOTTSCHLICH, P. GRÜNANGER, L. GUBELLINI, G. IIRITI, D. LUCARINI, D. MARCHETTI, B. MORALDO, L. PERUZZI, L. POLDINI, F. PROSSER, M. RAFFAELLI, A. SANTANGELO, E. SCASSELLATI, S. SCORTEGAGNA, F. SELVI, A. SOLDANO, D. TINTI, D. UBALDI, D. UZUNOV & M. VIDALI (2007). Integrazioni alla checklist della flora vascolare italiana. *Nat. Vicentina* 10: 5-74.
- DESOLE, L. (1948). Distribuzione geografica dell'*Ilex aquifolium* L. e del *Taxus baccata* L. in Sardegna. Prima Nota. *Atti Soc. Tosc. Sci. Nat. Pisa, Mem.* 55: 3-38.
- DESOLE, L. (1966). Distribuzione geografica dell'*Ilex aquifolium* L. e del *Taxus baccata* L. in Sardegna (Seconda e ultima nota.). *Boll. Ist. Bot. Reale Univ. Sassari* 7: 5-67.
- FENU, G. & G. BACCHETTA (2008). La flora vascolare della penisola del Sinis (Sardegna occidentale). *Acta Bot. Malac.* 33: 91-124.
- FENU, G., E. MATTANA & G. BACCHETTA (2011). Distribution, status and conservation of a Critically Endangered, extremely narrow endemic: *Lamyropsis microcephala* (Asteraceae) in Sardinia. *Oryx* 42: 180-186.
- FENU, G., E. MATTANA & G. BACCHETTA (2012). Conservation of endemic insular plants: the genus *Ribes* L. (Grossulariaceae) in Sardinia. *Oryx* 46: 219-222.
- FENU, G., E. MATTANA, A. CONGIU & G. BACCHETTA (2010). The endemic vascular flora of Supramontes: a priority plant conservation area in Sardinia. *Candollea* 65: 347-358.
- GREUTER, W., H. M. BURDET & G. LONG (ed.) (1984-2008). *Med-Checklist* 1, 2, 4. Conservatoire & Jardin botaniques de Genève, Genève.
- GUARINO, R. (2006). On the origin and evolution of the Mediterranean dry grasslands. *Ber. Reinhold-Tüxen Ges.* 18: 195-206.
- HERZOG, T. (1909). Über die Vegetationsverhältnisse Sardiniens. *Bot. Jahrb. Syst.* 42: 341-436.
- JALAS, J. & J. SUOMINEN (ed.) (1972-1994). *Atlas Fl. Eur.* 1-10. Helsinki University Printing House.
- JALAS, J., J. SUOMINEN & R. LAMPINEN (ed.) (1996-1999). *Atlas Fl. Eur.* 11-12. Helsinki University Printing House.
- JEANMONOD, D. & J. GAMISANS (2007). *Fl. Corsica*. Edisud.
- KURTTO, A., R. LAMPINEN & L. JUNIKKA (ed.) (2004). *Atlas Fl. Eur.* 13. Helsinki University Printing House, Helsinki.
- LADERO ALVAREZ, M., T. E. DÍAZ GONZÁLEZ, A. PENAS MERINO, S. RIVAS-MARTÍNEZ & C. VALLE GUTIÉRREZ (1987). Datos sobre la vegetación de las Cordillera Central y Cantábrica. *Itin. Geobot.* 1.
- MARTELLI, U. (1896-1904). *Monocotyledones Sardoae sive ad floram sardoam Josepho Hyacinthi Moris per Ugolino Martelli continuatio*. Vol. 1-3. Firenze.
- MATTANA, E., M. I. DAWS & G. BACCHETTA (2009). Seed dormancy and germination ecology of *Lamyropsis microcephala*: a mountain endemic species of Sardinia (Italy). *Seed Sci. Technol.* 37: 491-497.
- MÉDAIL, F. & K. DIADEMA (2009). Glacial refugia influence plant diversity patterns in the Mediterranean Basin. *J. Biogeogr.* 36: 1333-1345.
- MÉDAIL, F. & P. QUÉZEL (1997). Hot-spots analysis for conservation of plant biodiversity in the Mediterranean Basin. *Ann. Missouri Bot. Gard.* 84: 112-127.
- MORIS, G. G. (1827). *Stirpium sardoarum Elenchus*. Carali.
- MORIS, G. G. (1837-1859). *Flora Sardoae seu Historia Plantarum in Sardinia et adiacentibus insulis*. Vol. 1-3. Taurini.
- MITTERMEIER, R. A., P. ROBLES GIL, M. HOFFMANN, J. PILGRIM, T. BROOKS, C. G. MITTERMEIER, J. LAMOREUX & G. A. B. DA FONSECA (2004). *Hotspots Revisited*. CEMEX, Mexico.
- PERUZZI, L. (2010). Checklist dei generi e delle famiglie della flora vascolare Italiana. *Inform. Bot. Ital.* 42: 151-170.
- PIGNATTI, S. (1982). *Fl. Ital.* 1-3. Edagricole, Bologna.
- PIGNATTI, E. & E. FEOLI (1974). *Euphrasia minima* var. *genargentea*, nuova per la flora sarda. *Boll. Soc. Sarda Sci. Nat.* 14: 31-35.
- PIGNATTI, E. & S. PIGNATTI (1974). Osservazioni fitosociologiche sulla vegetazione rupestre delle montagne silicee della Sardegna. *Boll. Soc. Sarda Sci. Nat.* 14: 19-30.
- PIGNATTI, E., S. PIGNATTI, S. NIMIS & A. AVANZINI (1980). *La vegetazione ad arbusti spinosi emisferici: Contributo alla interpretazione delle fasce di vegetazione delle alte montagne dell'Italia mediterranea*. Collana del programma finalizzato Promozione della qualità dell'ambiente, C. N. R., Roma.

- PYŠEK, P., D. M. RICHARDSON, M. REJMÁNEK, G. L. WEBSTER, M. WILLIAMSON & J. KIRSCHNER (2004). Alien plants in checklist and floras: towards better communication between taxonomist and ecologists. *Taxon* 53: 131-143.
- RAUNKIAER, C. (1934). *The life forms of plants and statistical plant geography*. The Clarendon Press.
- RICHARDSON, D. M., P. PYŠEK, M. REJMÁNEK, M. G. BARBOUR, F. D. PANETTA & C. J. WEST (2000). Naturalization and invasion of alien plants: concepts and definitions. *Diversity & Distrib.* 6: 93-107.
- RÍOS RUIZ, S., F. ALCAZAR ARIZA & A. VALDÉS FRANZI (2003). *Vegetación de sotos y riberas de la Provincia de Albacete (España)*. Instituto de Estudios Albacetenses “Don Juan Manuel” De la Excma. Deputación de Albacete. Serie 1 - Estudios - Núm. 148, Albacete.
- RIVAS-MARTÍNEZ, S. (ed.) (2007). Mapa de series, geoserias y geopermaseries de vegetación de España. *Itin. Geobot.* 17: 5-436.
- RIVAS-MARTÍNEZ, S., T. E. DÍAZ, F. FERNÁNDEZ-GONZALES, J. IZCO, J. LOIDI, M. LOUSÁ & Á. PENAS (2002). Vascular plant communities of Spain and Portugal. *Itin. Geobot.* 15: 5-432.
- SCHMID, E. (1933). Beitrage zur Flora der Insel Sardinien. *Mitt. Bot. Mus. Univ. Zürich* 146: 232-255.
- TABERLET, P. & R. CHEDDADI (2002). Quaternary refugia and persistence of biodiversity. *Science* 297: 2009-2010.
- TUTIN, T. G., N. A. BURGESS, A. O. CHATER, G. R. EDMONDSON, W. H. HEYWOOD, D. M. MOORE, D. H. VALENTINE, S. M. WALTERS & D. A. WEBB (ed.) (1993). *Fl. Eur.* 1. Second edition. Cambridge University Press.
- TUTIN, T. G., N. A. BURGESS, D. H. VALENTINE, S. M. WALTERS & D. A. WEBB (ed.) (1964-80). *Fl. Eur.* 1-5. Cambridge University Press.
- TZEDAKIS, P. C., I. T. LAWSON, M. R. FROGLEY, G. M. HEWITT & R. C. PREECE (2002). Buffered tree population changes in a Quaternary refugium: evolutionary implications. *Science* 297: 2044.
- VALIENTE BANUET, A., A. VITAL RUMEBE, M. VERDU & R. M. CALLAWAY (2006). Modern quaternary plant lineages promote diversity through facilitation of ancient Tertiary lineages. *Proc. Natl. Acad. Sci. U.S.A.* 103: 812-817.