



An Earlier Name for *Lindernia Andringitrae* Eb. Fischer (Linderniaceae)

Author: Mats, Thulin

Source: *Candollea*, 68(1) : 63-64

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2013v681a6>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

27. THULIN, Mats: An earlier name for *Lindernia andringitrae* Eb. Fischer (Linderniaceae)

Lindernia All. was revised for Madagascar by FISCHER (1995), who recognized 16 species, seven of which are endemic. One of the endemic species is *L. andringitrae* Eb. Fischer, restricted to elevations above 2000 m in the Andringitra Massif. *Lindernia andringitrae* is a *nom. nov.* for *Craterostigma cerastioides* Bonati, a name that cannot be transferred to *Lindernia* because of the existence of *L. cerastioides* T. Yamaz. (YAMAZAKI, 1978), an Asian species. One further synonym, *Craterostigma perrieri* Bonati, was also cited by FISCHER (1995: 236) with the comment that its transfer to *Lindernia* “would also produce a homonym”. However, this is not correct and accordingly *Craterostigma perrieri* is transferred to *Lindernia*. An update of the synonymy and distribution of the species, including details of a new collection made in 2010 and a preliminary IUCN risk of extinction assessment (IUCN, 2012), is provided.

Lindernia perrieri (Bonati) Thulin, **comb. nova.**

= *Craterostigma perrieri* Bonati in Bull. Soc. Bot. Genève ser. 2, 15: 108. 1924.

Typus: MADAGASCAR. **Prov. Fianarantsoa:** Massif d’Andringitra, rocailles humides, 2200 m, II.1922, fl. & fr., *Perrier de la Bâthie 14423* (holo-: P [P00057076]; iso-: P [P00057077]).

= *Craterostigma cerastioides* Bonati in Bull. Soc. Bot. Genève ser. 2, 15: 106. 1924 [non *Lindernia cerastioides* T. Yamaz. in J. Jap. Bot. 53: 97. 1978]. = *Lindernia andringitrae* Eb. Fischer in Bull. Mus. Natl. Hist. Nat., sect. B, Adansonia 17: 236. 1995. **Typus:** MADAGASCAR. **Prov. Fianarantsoa:** Massif d’Andringitra, étangs temporaires de la saison des pluies, après le retrait des eaux, 2000-2400 m, IV.1921, fl. & fr., *Perrier de la Bâthie 13600* (holo-: P [P00057074]; iso-: P [P00057075]).

Observations. – *Homolle 1148* and *1294* have no locality information other than “Madagascar”, but *Homolle* is known to have collected in the Andringitra Massif in the 1940’s. Also the habitat description “bord de tourbières” strongly indicates that her collections were made in the higher parts of this Massif.

Two further collections were cited by FISCHER (1995), *Perrier de la Bâthie 14429* (P [P00346047, P00346048, P00346049]) and *Rakotovao 174* (P). However, *Perrier de la Bâthie 14429* is a syntype of *Carex andringitrensis* Cherm., and no collection of *Lindernia* corresponding to *Rakotovao 174* seems to be present in P.

Conservation status. – With an area of occupancy (AOO) of 9 km² and no more than three subpopulations, all encompassed in a protected area (Andringitra Massif), *Lindernia perrieri* is assigned a preliminary status of “Vulnerable” (VU D2) (calculation following CALLMANDER & al., 2007).

Additional collections examined. – **MADAGASCAR. Prov. Fianarantsoa:** Andringitra Massif, plateau d’Andohariana, dépression humide des plaques rocheuses, 2000-2100 m, 12.I.1971, fl. & fr., *Guillaumet 3737* (P [P00221519]); Andringitra Massif, plateau d’Andohariana, 22°09’03”S 46°53’19”E, open grassland in thin soil over rocks, 2100 m, 4.IV.2010, fl. & fr., *Thulin & Razafindraibe 11781* (TAN, UPS). **S. loc.:** bord de tourbières, fl. & fr., *Homolle 1148* (P [P03416462]); fl. & fr., *Homolle 1294* (P [P03416463]).

Acknowledgements

Field work in Madagascar was supported by the Swedish Research Council. *Direction Générale des Eaux et Forêts* and Madagascar National Parks are thanked for issuing collecting permits, Missouri Botanical Garden, Madagascar, for logistical support, Hanta Razafindraibe for excellent company in the field, and Martin Callmänder for assistance with the conservation assessment.

References

- CALLMANDER, M. W., G. E. SCHATZ, P. P. LOWRY II, M. O. LAIVAO, J. RAHARIMAMPIONONA, S. ANDRIAMBOLOLONERA, T. RAMINOSOA & T. K. CONSIGLIO (2007). Application of IUCN Red List criteria and assessment of priority areas for plant conservation in Madagascar: rare and threatened Pandanaceae indicate new sites in need of protection. *Oryx* 41: 168-176.
- FISCHER, E. (1995). Revision of the Lindernieae (Scrophulariaceae) in Madagascar. 1. The genera *Lindernia* All. and *Crepidorhopalon* E. Fischer. *Bull. Mus. Natl. Hist. Nat. Paris, sect. B, Adansonia* 17: 227-257.
- IUCN (2012). *IUCN Red List Categories and Criteria: Version 3.1*. 2nd Edition. IUCN Species Survival Commission. IUCN Gland & Cambridge.
- YAMAZAKI, T. (1978). New or noteworthy plants of the Scrophulariaceae from Indo-China (2). *J. Jap. Bot.* 53: 97-100.