

## **Dolichomitus irritator (Hymenoptera: Ichneumonidae): A New Parasite of *Dectes texanus* (Coleoptera: Cerambycidae) in Soybeans**

Authors: Tindall, Kelly V., and Fothergill, Kent

Source: Florida Entomologist, 95(1) : 238-240

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.095.0143>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

**DOLICHOMITUS IRRITATOR (HYMENOPTERA: ICHNEUMONIDAE): A NEW PARASITE OF *DECTES TEXANUS* (COLEOPTERA: CERAMBYCIDAE) IN SOYBEANS**

KELLY V. TINDALL<sup>1,2</sup> AND KENT FOTHERGILL<sup>2</sup>

<sup>1</sup>University of Missouri Division of Plant Sciences, PO Box 160, Portageville, MO 63873 USA

<sup>2</sup>Conservation Seeding and Restoration, Inc. 506 Center Street West, Kimberly, ID 83341 USA

Corresponding author; E-mail: ktindall@csr-inc.com

The Ichneumonidae is a large insect family with over 4,900 species in the Nearctic region (Yu et al. 2005). Ichneumonids are obligate parasitoids, or hyperparasitoids, but they have not been used as successfully as classical biological control agents as hymenopteran families within Chalcidoidea and the Braconidae (Daly et al. 1998). *Dolichomitus irritator* (F.) (Ichneumonidae: Pimplinae) is common in woodland and brush habitats in eastern North America (Townes & Townes 1960) and occurs south to Costa Rica (Cancino et al. 2010). *Dolichomitus irritator* larvae are external parasites (Chittenden 1893) of larvae of wood boring insects (Townes & Townes 1960). Table 1 lists previously known insect hosts on which development of *D. irritator* has been witnessed or implied and the associated plant relationship. Unequivocal host records are very difficult to attain for parasites of wood boring insects and these records are often inferred from rearing parasite and host in the same logs.

*Dectes texanus* LeConte, is a native, univoltine, cerambycid beetle that, as a larva, primarily feeds within herbaceous plants in the Asteraceae over much of North America (Lingafelter 2007). Falter (1969) and Hatchett et al. (1973) first documented a host switch from plants in Asteraceae to soybeans (*Glycine max* (L.) Merr.; Fabales: Fabaceae). Since then, *D. texanus* larvae have been noted as a pest of soybeans in 14 U.S. states (Buschman & Sloderbeck 2010). Several species of braconids, ichneumonids, and pteromalids are known to parasitize *D. texanus* larvae in giant ragweed (*Ambrosia trifida* L.) (Hatchett et al. 1975); however, in soybean *D. texanus* larvae are known to be parasitized only by pteromalid wasps (Tindall, unpublished data) and a tachinid fly, *Zelia tricolor* (Coquillett) (Tindall & Fothergill 2010). Discovery of parasitoids of *D. texanus* larvae within soybeans offers opportunity for producers to manage for biological control within soybean production systems.

Soybean stems (i.e., stubble) were collected from a soybean field harvested in the fall of 2009 on 23 Mar 2010 in New Madrid County, Missouri (N 36.42482° W -89.64933°). From this set, 480 were selected based on the presence of a frass plug, an indicator of occupancy by *D. texanus* (Hatchett et al. 1975), and subsequently kept in

an insect rearing room (16:8 h L:D, 24 °C). Five *D. irritator* were recovered from these stems (Table 2). Three additional individuals were found during other stubble-based *D. texanus* survey work from soybean stems from the same field (Table 2). The *D. irritator* specimens recovered were sent to the American Entomological Institute (Gainesville, Florida) for identification by Dr. David Wahl. Six of the specimens were retained by the American Entomological Institute collection and 2 are deposited in the collection of KVT. *D. texanus* as the insect host for these *D. irritator* is confirmed by:

- 1) the timing of stem harvest (Fothergill et al. 2010),
- 2) morphology of stem tunneling (Fothergill et al. 2010),
- 3) discovery of a *D. texanus* larva carcass in a soybean stem with a *D. irritator* pupa (Table 2, specimen #6), and
- 4) emergence of only *D. texanus* and a pteromalid wasp (data not shown) from this cohort of soybeans.

This is, to our knowledge, the first documented case of *D. irritator* parasitizing *D. texanus* and using a host boring within an annual, herbaceous plant. It is likely that further research will find other insect hosts and plant associations utilized by *D. irritator*.

These records indicate that *D. irritator* can overwinter with *D. texanus* larvae in soybean stubble. The five *D. irritator* adults recovered from the above mentioned cohort of 480 soybean stubble, which contained overwintering *D. texanus* larvae, represent a field parasitism rate of 1%. Additional work is needed to determine the parasitism rate of *D. irritator* in other populations of soybean utilizing *D. texanus* and what role it may play in *D. texanus* population dynamics.

*Dolichomitus irritator* has been found visiting flowers of *Salix discolor* Muhl. (Graenicher 1900) (Malpighiales: Salicaceae) and the term: "Ichneumon flowers" (Knuth 1906) has been coined to describe flowers that attract Ichneumonids and other similar insects. Bianchi et al. (2006) concluded that field margins and other non-crop habitats can enhance the abundance and diversity of natural enemy species within an agricul-

TABLE 1. PREVIOUSLY KNOWN INSECT HOSTS OF *DOLICHOMITUS IRRITATOR* AND ASSOCIATED PLANTS.

Host Insect		Host Plant			Source
Family	Species	Family	Species		
Buprestidae	<i>Chrysobothris</i> sp.	Fabaceae	<i>Cercis canadensis</i> L.	Champlain 1922	
Cerambycidae	"Leptura"	Fagaceae	<i>Quercus</i> sp.	Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Anelaphus parallelus</i> (Newman)			Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Astyleiopus variegatus</i> (Hald.)	Aceraceae	<i>Acer negundo</i> L.	Chittenden 1893	
Cerambycidae	<i>Elaphidion</i> sp.	Fagaceae	<i>Quercus</i> sp.	Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Graphisurus fasciatus</i> (DeGeer)			Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Megacyllene caryae</i> (Drury)	Juglandaceae	<i>Carya</i> sp.	Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Megacyllene caryae</i> (Drury)			Champlain 1922	
Cerambycidae	<i>Megacyllene robiniae</i> (Forster)	Fabaceae	<i>Robinia pseudoacacia</i> L.	MacAndrews 1933	
Cerambycidae	<i>Obrium rufulum</i> Gahan	Oleaceae	<i>Fraxinus</i> sp.	Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Pachyta lamed</i> (L.)			Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Parelaphidion incertum</i> (Newman)	Juglandaceae	<i>Carya</i> sp.	Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Rhagium inquisitor</i> (L.)			Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Saperda discoidea</i> F.			Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Trigonarthris proxima</i> (Say)			Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	Unknown	Aceraceae	<i>Acer</i> sp.	Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Xylotrechus annosus</i> (Say)	Fagaceae	Oak	Chittenden 1893	
Cerambycidae	<i>Anelaphus villosus</i> (F.)	Fagaceae	<i>Castanea dentata</i> (Marsh.) Borkh.	Townes and Townes 1960 <sup>1</sup>	
Cerambycidae	<i>Anelaphus villosus</i> (F.)	Fagaceae	<i>Quercus</i> sp.	Townes and Townes 1960 <sup>1</sup>	
Curculionidae	<i>Cryptorynchus lapathi</i> (L.)			Chittenden 1904	
Melandryidae	<i>Melandrya striata</i> Say	Fagaceae	<i>Fagus</i> sp.	Townes and Townes 1960 <sup>1</sup>	
Synchroidae	<i>Synchroa punctata</i> Newman			Townes and Townes 1960 <sup>1</sup>	
Sesiidae	<i>Synanthedon exitosa</i> Say	Rosaceae	<i>Prunus persica</i> (L.) Batsch.	Gossard and King 1918	
Sesiidae	<i>Synanthedon tipuliformis</i> (Clerck)			Townes and Townes 1960 <sup>1</sup>	
Unknown	Unknown	Ulmaceae	Elm	Zimmer 1909	

<sup>1</sup>Based on label data from collections.

TABLE 2. *DOLICHOMITUS IRRITATOR* SPECIMENS REARED FROM 2009 GROWING SEASON SOYBEAN STUBBLE INHABITED BY *DECTES TEXANUS* AND HARVESTED FROM NEW MADRID COUNTY, MISSOURI.

Specimen #	Stem harvest date	Discovery Date	Comment
1	3/23/2010	3/29/2010	adult in rearing room with stems
2	3/23/2010	3/30/2010	adult in 2 L container with stems
3	3/23/2010	3/30/2010	adult in 2 L container with stems
4	3/23/2010	4/2/2010	adult in 2 L container with stems
5	3/23/2010	4/3/2010	adult in 2 L container with stems
6	3/24/2010	3/24/2010	pupa in stem w/ <i>D. texanus</i> larva carcass -3/27/2010 adult eclosed
7	4/5/2010	4/7/2010	adult within a plastic bag of stems
8	4/5/2010	4/5/2010	pupa in <i>D. texanus</i> tunneled soybean -4/6/2010 adult eclosed

tural landscape. These findings suggest it may be possible for soybean producers to manage non-crop habitats to attract and/or produce a source of natural enemies to aid in control of *D. texanus*. Further research may elucidate these processes and discover other parasites of *D. texanus*.

#### SUMMARY

*Dolichomitus irritator* (Fabricus) was discovered utilizing *Dectes texanus* LeConte larvae as hosts within the previous year's soybean stubble in early spring in southeast Missouri. This represents the first records of *D. irritator* utilizing hosts within herbaceous plants and the first records of an Ichneumonid wasp parasitizing *D. texanus* in soybeans.

#### REFERENCES CITED

- BIANCHI, F. J. J. A., BOOLJ, C. J. H., AND TSCHARNTKE, T. 2006. Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proc. Royal Soc. B* 273: 1715-1727.
- BUSCHMAN, L. L., AND SLODERBECK, P. E. 2010. Pest status and distribution of the stem borer, *Dectes texanus*, in Kansas. *J Insect Sci* 10:198 available online: insect-science.org/10.198.
- CANCINO, E. R., KASPARYAN, D. R., CORONADO BLANCO, J. M. A., MYARTSEVA, S. N., TRJAPITZIN, V. A., HERNANDEZ, S. G., AND JIMENEZ, S. G. 2010. Himenópteros de la Reserva "El Cielo", Tamaulipas, México. *Dugesiana* 17.1: 53-71
- CHAMPLAIN, A. B. 1922. Records of Hymenopterous parasites in Pennsylvania. *Psyche* 29: 95-100
- CHITTENDEN, F. H. 1893. Observations of some hymenopterous parasites of Coleoptera. *Insect life* 5:247-251.
- CHITTENDEN, F. H. 1904. Insects injurious to the basket willow, *In* W. F. Hubbard [ed.], *The Basket Willow*. Farmers' Bulletin 46, USDA, Government Printing Office. Washington DC., 100 pp.
- DALY, H. V., DOYEN, J. T., AND PURCELL, A. H. III. 1998. Introduction to insect biology and diversity, 2nd edition. Oxford Univ. Press: New York. 680 pp.
- FALTER, J. M. 1969. *Dectes* sp. (Coleoptera: Cerambycidae): A unique and potential important pest of soybeans. *J. Elisha Mitchell Sci. Soc.* 85: 123
- FOTHERGILL, K., CROSS, C. B., TINDALL, K. V., WRATHER, J. A., LORENZ, G., AND YOUMANS, C. 2010. Clover stem borer, *Languria mozardi* (Coleoptera: Languriidae), on soybeans, *Glycine max*: a new host record. *Florida Entomol.* 93:125-127.
- GOSSARD, H. A., AND KING, J. L. 1918. The peach tree borer *Sanninoidea exitiosa* Say. *Bull. Ohio Agric. Exp. Stn.* 329: 55-87
- GRAENICHER, S. 1900. The fertilization and insect visitors of our earliest entomophilous flowers. *Bull. Wisconsin Nat. Hist. Soc.* 1: 73-84.
- HATCHETT, J. H., JACKSON, R. D., BARRY, R. M., AND HOUSER, E. C. 1973. Rearing a weed cerambycid, *Dectes texanus*, on an artificial medium, with notes on biology. *Ann. Entomol. Soc. Am.* 66: 519-522.
- HATCHETT, J. H., DAUGHERTY, D. M., ROBBINS, J. C., BARRY, R. M., AND HOUSER, E. C. 1975. Biology in Missouri of *Dectes texanus*, a new pest of soybean. *Ann. Entomol. Soc. Am.* 68: 209-213.
- KNUTH P. 1906. Handbook of flower pollination. Clarendon Press: Oxford, MA. 382 pp.
- LINGAFELTER, S. W. 2007. Illustrated key to the long-horned woodboring beetles of the eastern United States. North Potomac, MD: The Coleopterists Society 206 pp.
- MACANDREWS, A. H. 1933. The control of the locust borer by forest management. *Rep. Entomol. Soc. Ontario* 63: 48-50.
- TINDALL, K. V., AND FOTHERGILL, K. 2010. *Zelia tricolor* (Coquillett) (Diptera: Tachinidae): first record of an internal parasite of *Dectes texanus* Leconte (Coleoptera: Cermabycidae) in soybean. *Florida Entomol.* 93: 635-636.
- TINDALL, K. V., STEWART, S., MUSSER, F., LORENZ, G., BAILEY, W., HOUSE, J., HENRY, R., HASTINGS, D., WALLACE, M., AND FOTHERGILL, K. 2010. Distribution of *Dectes texanus* in Soybeans of Missouri, Western Tennessee, Mississippi, and Arkansas, USA. <http://www.insectscience.org/10.178/i1536-2442-10-178.pdf>. *J Insect Sci* 10: Article 178.
- TOWNES, H., AND TOWNES, M. 1960. Ichneumon-flies of America north of Mexico: 2. Subfamilies Ephialtinae, Xoridinae, and Acaenitinae. *United States Nat. Mus. Bull.* 216(2): 1-676.
- YU, D., VAN ACHTERBERG, K., AND HORSTMANN, K. 2005. World Ichneumonoidea 2004. Taxonomy, biology, morphology and distribution. [CD/DVD]. Taxapad®, Vancouver, Canada. [Available at <http://www.taxapad.com>].
- ZIMMER, J. F. 1909. List of insects affecting the maple. *Ohio Nat.* 10(2): 36-38.