



## **Cylindrocopturus (Coleoptera: Curculionidae: Conoderinae) Species Associated with Opuntia (Caryophyllales: Cactaceae) Species**

Authors: Martínez, Néstor Bautista, Riquelme, Carlos Patricio Illescas, and Jones, Robert W.

Source: Florida Entomologist, 99(1) : 126-127

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.099.0125>

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# ***Cylindrocopturus* (Coleoptera: Curculionidae: Conoderinae) species associated with *Opuntia* (Caryophyllales: Cactaceae) species**

**Néstor Bautista Martínez<sup>1</sup>, Carlos Patricio Illescas Riquelme<sup>1,\*</sup>, and Robert W. Jones<sup>2</sup>**

Some of the plants that belong to the genus *Opuntia* (Caryophyllales: Cactaceae) are highly valued vegetables in Mexican culture. The fruits (prickly pears) and the so-called “nopalitos” (young cladodes) are mainly used for human consumption, but also are used for livestock feed and as raw material for various industrial products (Sáenz et al. 2006). Nopal exploitation is done under 3 systems, namely, wild plantations, family gardens, and commercial plantations (Flores-Valdez 2001). This plant sometimes is affected by several insect pests that hinder production, making it necessary to have a correct identification of each of them to make handling decisions.

In Sep 2013, nopal growers of San Sebastian Xololapa, Teotihuacan Municipality, state of Mexico, Mexico, reported a strong presence of resinous secretions emerging from the base of the spines of mature cladodes. This study initially was conducted to identify the cause of this symptomatology. Twenty cladodes were collected and transported to the Agricultural Entomology laboratory of the Colegio de Postgrados, campus Montecillo and confined in plastic containers covered with organdy gauze to await the emergence of the insects that caused the damage.

Adults started emerging on 2 Jan 2014. They were collected and preserved in 70% ethyl alcohol for later identification. Extraction of genitalia from the males was done through tissue maceration in KOH (10%). The results obtained from the taxonomic study were that 2 spine weevils coexisted in these *Opuntia* plants. Twenty-one adults of *Cylindrocopturus biradiatus* Champion and 16 of *Cylindrocopturus ganglbaueri* Heller (Coleoptera: Curculionidae) emerged from the collected cladodes. Distinctive characteristics can be seen to differentiate both species with the naked eye (Fig. 1). *Cylindrocopturus biradiatus* is smaller, with a mean length of 3.32 mm, it has frontal and lateral areas of the pronotum with orange color and light brown lines running parallel on the inner margin of the elytra with small perpendicular lines of the same color forming crosses that do not reach the outer margin of the elytra. *Cylindrocopturus ganglbaueri* is larger with a total mean length of 4.82 mm, with predominantly dark coloring and a black transverse dorsomedial line on its elytra.

Dampf (1929) reported on the presence of the nopal spine weevil, *C. biradiatus*. Later publications continued to mention the same species affecting nopal in several regions in Mexico, occasionally as an important pest for the crop (Flores-Valdez 2001; Sáenz et al. 2006; Var-

gas-Mendoza et al. 2008; Angeles-Nuñez et al. 2014). We detected that most reports are based only on symptomatology of the cladodes, and it was taken for granted that it was only due to *C. biradiatus* without confirming the taxonomy of the species. For example, Badii & Flores (2001) mentioned the incidence of *C. biradiatus* and described it as follows: “It is 4–4.5 mm long and has a dark dorsal mark in the form of a cross.” At least to our perception, this corresponds to the description of *C. ganglbaueri*.

On the other hand, Muñiz (1976) and Salas-Araiza et al. (2001) mentioned both species of spine borers; however, the reports show little information about these weevils. In this note, the aedeagus of each is shown and compared for the first time (Fig. 1). Both species show smooth margins with a round, slightly inclined tip, with greater sclerotization in *C. biradiatus* than *C. ganglbaueri*.

## **Summary**

In Sep 2013, nopal growers in Teotihuacan, Mexico, reported resinous secretions emerging from the base of the spines of mature cladodes. Cladodes with symptoms were collected, and the weevil species *Cylindrocopturus biradiatus* Champion and *Cylindrocopturus ganglbaueri* Heller (Coleoptera: Curculionidae) emerged. This work confirms the presence of both species, and it provides illustrations and descriptions for their diagnosis.

Key Words: spine borer; prickly pear cactus; nopal

## **Sumario**


En Sep 2013, productores de nopal en Teotihuacán, México, reportaron escurreimientos de resina que emergían de la base de las espinas de cladodios maduros. Se colectaron cladodios con síntomas y emergieron los picudos *Cylindrocopturus biradiatus* Champion y *Cylindrocopturus ganglbaueri* Heller (Coleoptera: Curculionidae). El presente trabajo confirma la presencia de ambas especies, además proporciona ilustraciones y descripciones para su diagnóstico.

Palabras Clave: barrenador de la espina; nopal

<sup>1</sup>Programa de Entomología y Acarología, Colegio de Postgrados Campus Montecillo, Km 36.5 carretera México-Texcoco Montecillo Texcoco estado de México, México

<sup>2</sup>Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro, 76010, Querétaro, México

\*Corresponding author; E-mail: entoillescas@gmail.com



**Fig. 1.** *Cylindrocopturus* species associated with *Opuntia*. A) *Cylindrocopturus biradiatus*, lateral view; B) *C. biradiatus*, dorsal view; C) *C. ganglbaueri*, lateral view; D) *C. ganglbaueri*, dorsal view; E) *C. biradiatus*, aedeagus; F) *C. ganglbaueri*, aedeagus.

## References Cited

- Angeles-Nuñez JG, Anaya-Lopez JL, Arévalo-Galarza ML, Leyba-Ruelas G, Anaya-Rosales S, Martínez-Martínez TO. 2014. Análisis de la calidad sanitaria de nopal verdura en Otumba, Estado de México. Revista Mexicana de Ciencias Agrícolas 5: 129–141.
- Badii MH, Flores AE. 2001. Prickly pear cacti pests and their control in Mexico. Florida Entomologist 84: 503–505.
- Dampf A. 1929. Una nueva plaga del nopal (*Opuntia* sp.): *Cylindrocopturus biradiatus* Champion (Ins. Col., Fam. Curculionidae). Talleres Gráficos de la Secretaría de Agricultura y Fomento, Mexico.
- Flores-Valdez CA. 2001. Producción, Industrialización y Comercialización de Nopalitos. No. 58. Centro de Investigaciones Económicas, Sociales y Tecnológicas de la Agroindustria y la Agricultura Mundial (CIESTAAM). Universidad Autónoma de Chapingo, Chapingo, Mexico.
- Muñiz R. 1976. Algunas plagas de cactáceas, pp. 71–73 In Órgano de la Sociedad Mexicana de Cactología, A.C. Cactáceas y Suculentas Mexicanas, Mexico D.F.
- Sáenz C, Berger H, Corrales J, Galletti L, García V, Higuera I, Mondragón C, Rodríguez-Félix A, Sepúlveda E, Varnero M. 2006. Utilización Agroindustrial del Nopal. Boletín de Servicios Agrícolas de la FAO. Organización de las Naciones Unidas para la Agricultura y la Alimentación, Rome, Italy.
- Salas-Araiza MD, O'Brien CW, Romero-Nápoles J. 2001. Curculionoidea (Insecta: Coleoptera) from the state of Guanajuato, Mexico. Insecta Mundia 15: 45–57.
- Vargas-Mendoza A, Flores-Hernández A, Basaldua-Suarez JF. 2008. Dinámica poblacional de las principales plagas de nopal *Opuntia* spp. en la zona semiarida de Querétaro. Revista Chapingo Serie Zonas Áridas 7: 21–27.