



Examining the Relationship between Flower Thrips (Thysanoptera: Thripidae) Spatial Distribution and Blueberry (Ericales: Ericaceae) Flower Density

Authors: Rhodes, Elena M., Liburd, Oscar E., and Grunwald, Sabine

Source: Florida Entomologist, 99(1) : 128-129

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.099.0126>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Examining the relationship between flower thrips (Thysanoptera: Thripidae) spatial distribution and blueberry (Ericales: Ericaceae) flower density

Elena M. Rhodes^{1,*}, Oscar E. Liburd¹, and Sabine Grunwald²

Southern highbush blueberry (*Vaccinium corymbosum* L. × *V. darrowii* Camp) (Ericales: Ericaceae) is an important crop in Florida, where it is grown for the highly profitable early-season fresh market (USDA 2010). Flower thrips (*Frankliniella* species) are one of the key pests of blueberry. In Florida, *Frankliniella bispinosa* (Morgan) (Thysanoptera: Thripidae) is the dominant species collected (Arévalo & Liburd 2007). The thrips injure blueberry flowers when they feed and oviposit in developing flower tissues. This injury can cause scarring on developing fruit that makes the fruit unmarketable (Arévalo-Rodríguez 2006).

Thrips populations tend to form one or a few “hot spots,” small areas of comparatively high thrips numbers, on blueberry plantings (Arévalo & Liburd 2007). These hot spots begin forming about 7 to 10 d after bloom initiation, peak when the majority of the flowers are open (12–15 d after initiation), and decline until most of the flowers have become fruit (approx. 22 d after initiation) and very few thrips remain (Arévalo & Liburd 2007). The objective of this study was to determine if thrips density is related to flower density. The hypothesis was that thrips population density in space has a positive linear relationship with flower density. Two experiments were conducted, one on a farm in Inverness, Florida, in 2009 and another on a farm in Windsor, Florida, in 2010. White traps were used as the standard monitoring tool for thrips because Liburd et al. (2009) found them to be an effective device for monitoring thrips in southern highbush blueberries. Arévalo & Liburd (2007) found a strong correlation ($r = 0.76$) between thrips per flower and thrips per trap in rabbiteye blueberries (*Vaccinium virgatum* Aiton).

On the Inverness farm, 100 white sticky traps (Great Lakes IPM, Vestaburg, Michigan) were distributed throughout a 1 ha, 4- to 7-yr-old southern highbush blueberry planting in a regular grid at 7.61 m increments. An additional 30 traps were placed randomly throughout the plot. Traps were replaced weekly over a 5 wk period from 30 Jan to 26 Feb 2009 and taken to the University of Florida, Small Fruit and Vegetable Laboratory in Gainesville, Florida, where the number of thrips per trap was counted and recorded.

The percentage of open flowers present in each blueberry row in the study was recorded when traps were collected. The sampling area consisted of 38 rows with a dirt road in the middle (19 rows on either side of the dirt road). Each row contained blueberry plants of the same variety and age.

Linear regression analysis was used to determine if the number of thrips (dependent variable) was related to the percentage of open flowers (independent variable). All 130 sample points were input into

the analysis by assigning to each trap the percentage of open flowers recorded from the row in which the trap was hung. Because some of the assumptions of least squares regression could not be met even after transformation, Theil regression (Hollander & Wolfe 1999) was used and Kendall's tau, a nonparametric correlation statistic (Hollander & Wolfe 1999), was calculated (Wessa 2008) for all sampling dates.

At the Windsor farm, a smaller study was conducted due to space constraints: 20 white sticky traps were placed in a 2,464 m² area of a 7-yr-old southern highbush blueberry planting. Traps were spaced approximately 15 m apart within each of 5 blueberry rows. The spacing between rows was 10 m. Each trap was replaced weekly. Thrips data were collected as in the Inverness study. Percentage of open flowers was recorded from each sampled plant where traps were collected. Traps and flower data were collected for 6 wk from 18 Feb to 25 Mar 2010.

Least squares regression analysis (SAS Institute 2002) was used to determine if the number of thrips per trap (dependent variable) was related to the percentage of open flowers (independent variable). The thrips per trap (x) data were $\log_{10}(x + 1)$ transformed to meet assumptions of least squares regression.

On the Inverness farm, a significant positive linear relationship between percentage of open flowers and thrips per trap occurred on 30 Jan ($\tau = 0.36$, $C = 2,594$, $n = 130$, $P_{\text{slope}} < 0.0001$; Fig. 1a) and 5 Feb ($\tau = 0.24$, $C = 1,791$, $n = 130$, $P_{\text{slope}} = 0.0002$; Fig. 1b). No relationship was found between the 2 variables on any other date (all $\tau \leq 0.07$, $C \leq 556$, $n = 130$, $P_{\text{slope}} \geq 0.25$).

On the Windsor farm, there was a significant negative linear relationship between percentage of open flowers and \log_{10} thrips per trap on 18 Mar ($R^2 = 0.24$, $t = -2.67$, $df = 19$, $P_{\text{slope}} = 0.0156$; Fig. 2). No relationship was found between the 2 variables on any other date (all $R^2 < 0.03$, $|t| \leq 1.23$, $df = 19$, $P_{\text{slope}} > 0.23$).

According to Arévalo-Rodríguez (2006), flower thrips population density is strongly correlated with the percentage of open flowers over time. The results from the Inverness study indicate that this relationship may exist in space, also. Differences in percentages of open flowers in space and time probably exist because multiple varieties are grown on the same farm to maximize cross pollination (Childers & Lyrene 2006). Different varieties begin to flower at different times and for different lengths of time (Rhodes et al. 2012).

An intense cold snap that occurred from 4 to 6 Feb 2009 (FAWN 2009) may explain the low thrips numbers recorded on 6 Feb on the Inverness farm. The lack of a relationship between thrips and flowers on 13 Feb probably happened because peak flowering occurred dur-

¹Entomology and Nematology Department, University of Florida, Gainesville, Florida 32611, USA

²Soil and Water Sciences Department, University of Florida, Gainesville, Florida 32611, USA

*Corresponding author; E-mail: erhodes@ufl.edu




Fig. 1. Graphs showing percentage of open blueberry flowers vs. thrips per trap from the Inverness farm on a) 30 Jan and b) 5 Feb 2009. The black lines represent regression lines fitted by Theil regression.

ing this week. The thrips population continued to increase after peak flowering for 2.5 wk, possibly due to delayed development (transition from larvae and pupae to adults), which may have resulted in the lack of a relationship on 20 and 26 Feb.

On the Windsor farm, a negative linear relationship between thrips and flowers occurred on 18 Mar. This, and the lack of any other significant relationships, may have been a result of the unusually cold winter weather that occurred throughout Jan and Feb 2010 (FAWN 2010).

Although numbers of thrips on sticky traps are correlated with thrips per flower (Arévalo & Liburd 2007), using flower samples might produce a stronger correlation with flower density. Sticky traps only monitor for adults, whereas larvae are present in flowers along with adults. More accurate measures of flower density are also needed.

Results from the Inverness study indicate that flower thrips density is related to flower density. A better understanding of the relationship




Fig. 2. Graph of percentage of open blueberry flowers vs. \log_{10} thrips per trap from the Windsor farm on 18 Mar 2010. The black line represents a regression line fitted by least squares regression.

between thrips density and flower density could help in revising management programs. For example, scouting could be concentrated in areas where thrips density is likely to be high, and insecticide applications could be made to those sites where population density is high as opposed to whole field applications.

We thank Drs. Robert McSorley and Joe Funderburk for their initial review of this manuscript. We also thank the Florida Blueberry Grower's Association for allowing us to do this study on one of their farms. Many thanks also go to the staff and students of the Small Fruit and Vegetable IPM laboratory for their help in collecting samples.

Summary

Flower thrips tend to form areas of high population, termed "hot spots," in blueberry plantings. This study provided evidence that thrips density may be correlated with flower density in space. Further understanding of the relationship between flower density and thrips numbers could lead to better scouting and management programs.

Key Words: *Frankliniella*; hot spot; nonparametric correlation; Theil regression; southern highbush

Sumario

Los trips de las flores suelen formar zonas de alta población, que se llaman "puntos calientes", en las plantaciones de arándanos. Este estudio provee evidencia de que la densidad de trips se puede correlacionar con la densidad de las flores en espacio. Una mayor comprensión de la relación entre la densidad de las flores y el número de trips podría resultar en mejores programas de muestreo y de manejo.

Palabras Clave: *Frankliniella*; punto caliente; correlación no paramétrica; Regresión Theil; arándanos de variedad highbush de sur

References Cited

- Arévalo HA, Liburd OE. 2007. Horizontal and vertical distribution of flower thrips in southern highbush and rabbiteye blueberry plantings, with notes on a new sampling method for thrips inside blueberry flowers. *Journal of Economic Entomology* 100: 1622–1632.
- Arévalo-Rodríguez HA. 2006. A study of the behavior, ecology, and control of flower thrips in blueberries towards the development of an integrated pest management (IPM) program in Florida and southern Georgia. Ph.D. dissertation, University of Florida, Gainesville, Florida.
- Childers NF, Lyrene PM. 2006. *Blueberries for Growers, Gardeners, Promoters*. Dr. Norman F. Childers Publications, Gainesville, Florida.
- FAWN. 2009. Florida Automated Weather Network Report Generator. University of Florida, Gainesville, Florida, <http://fawn.ifas.ufl.edu/data/reports/> (last accessed 9 Nov 2015).
- FAWN. 2010. Florida Automated Weather Network Report Generator. University of Florida, Gainesville, Florida, <http://fawn.ifas.ufl.edu/data/reports/> (last accessed 9 Nov 2015).
- Hollander M, Wolfe DA. 1999. *Nonparametric Statistical Methods*. John Wiley & Sons, Inc., New York, New York.
- Liburd OE, Sarzynski EM, Arévalo HA, MacKenzie K. 2009. Monitoring and emergence of flower thrips species in rabbiteye and southern highbush blueberries. *Acta Horticulturae* 810: 251–258.
- SAS Institute. 2002. *The SAS system 9.0 for Windows*. SAS Institute, Cary, North Carolina.
- Rhodes EM, Liburd OE, England GK. 2012. Effects of southern highbush blueberry variety and treatment threshold on flower thrips populations. *Journal of Economic Entomology* 105: 480–489.
- USDA. 2010. *Noncitrus fruits and nuts 2009 summary*, pp. 31–32. United States Department of Agriculture, National Agricultural Statistics Service. www.nass.usda.gov/ (last accessed 9 Nov 2015).
- Wessa. 2008. Kendall tau Rank Correlation (v1.0.10) in Free Statistics Software (v1.1.23–r6). Office for Research Development and Education, http://www.wessa.net/rwasp_kendall.wasp/ (last accessed 9 Nov 2015).