



Influence of Gamma-Irradiation on Flight Ability and Dispersal of *Conopomorpha sinensis* (Lepidoptera: Gracillariidae)

Authors: Zhang, Ke, Fu, Haohao, Zhu, Shaowen, Li, Zhibin, Weng, Qun-fang, et al.

Source: Florida Entomologist, 99(sp1) : 79-86

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.099.sp111>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Influence of gamma-irradiation on flight ability and dispersal of *Conopomorpha sinensis* (Lepidoptera: Gracillariidae)

Ke Zhang, Haohao Fu, Shaowen Zhu, Zhibin Li, Qun-fang Weng* and Mei-ying Hu*

Abstract

Assessment of quality of male insects has been done primarily in area-wide integrated pest management (AW-IPM) programs that have a sterile insect technique (SIT) component. Routine monitoring of sterile male quality needs to be carried out both in the mass-rearing facility and in the field. Simple bioassays, which can be conducted in the laboratory or under semi-field conditions, would be potential surrogates for laborious field tests that are usually very costly. In the laboratory, a flight mill system was used to assess the quality of males of the litchi stem-end borer, *Conopomorpha sinensis* (Lepidoptera: Gracillariidae), in terms of flight distance, flight duration and speed. Flight distance, duration, mean speed and greatest speed of non-irradiated adult males were 13,926 m, 29,365 s, 0.42 m/s, and 1.01 m/s, respectively, during a 24 h period. Although the values of these parameters of non-irradiated males were greater than corresponding values for moths irradiated with 150 and 200 Gy, there were no significant differences between the various treatments and the non-irradiated control group. These data suggest that irradiation with 150 and 200 Gy did not significantly affect the flight ability of male litchi stem-end borers. This study also included 2 field release and recapture experiments. The data of the field experiments indicated that recapture rates, dispersal distances, and dispersal directions of the 150 and 200 Gy irradiated males were not significantly different from those of non-irradiated males. These data indicate that the ability of litchi stem-end borer males irradiated with doses of 150 and 200 Gy to disperse in the field was not impaired in comparison with non-irradiated males.

Key Words: irradiation; sterile insect technique; release/recapture; flight mill; dispersal

Resumen

La evaluación de la calidad de los insectos machos estériles o parcialmente estériles se ha hecho principalmente en relación con programas de manejo de plagas integrado en toda la área (MIP-TA) que tienen un componente de técnica del insecto estéril (TIE). El monitoreo rutinario de la calidad de los machos esteriles necesita ser realizado tanto en las instalaciones de cría masiva como en el campo. Bioensayos simples, que pueden ser realizados en el laboratorio o en condiciones de semi-campo, serían sustitutos potenciales para las pruebas laboriosas de campo para monitorear el desempeño de los machos estériles en el campo que por lo general son costosos. En el laboratorio, se utilizó un sistema de molino de vuelo para evaluar la calidad de machos del barrenador del tallo de litchi (*Conopomorpha sinensis*; Lepidoptera: Gracillariidae) en términos de distancia, tiempo y velocidad de vuelo. La distancia del vuelo, la duración del vuelo, el promedio de la velocidad y la mayor velocidad de vuelo de los machos adultos no irradiados fueron 13,926 m, 29,365 s, 0.42 m/s, y 1.01 m/s, respectivamente, dentro de un período de 24 h. Aunque los valores de estos parámetros de los machos no irradiados fueron mejores que los de las polillas que habían sido irradiadas con 150 y 200 Gy, no hubo diferencia significativa entre los diferentes tratamientos y el grupo control no tratado. Estos datos sugieren que la irradiación con 150 Gy y 200 Gy no afectó significativamente la capacidad de vuelo de los machos del barrenador del tallo de litchi. Este estudio también incluyó 2 experimentos de liberación de campo y recaptura. Los datos de los experimentos de campo indicaron que la tasa de recaptura, la distancia de dispersión, y la dirección dispersión de las polillas tratadas con 150 Gy y 200 Gy no fueron significativamente diferentes de las de los machos no irradiados. Esto indica que la capacidad de dispersión en el campo de los machos del barrenador del tallo de litchi irradiados con dosis de 150 y 200 Gy no difieren significativamente de los de los machos no irradiados.

Palabras Clave: irradiación; técnica del insecto estéril; liberar/recaptura; molino de vuelo, dispersión

The litchi stem-end borer, *Conopomorpha sinensis* Bradley (Lepidoptera: Gracillariidae) is an important pest of litchi, *Litchi chinensis* Sonn. (Sapindales: Sapindaceae) and longan, *Dimocarpus longan* Lour. (Sapindales: Sapindaceae). Litchi stem-end borers not only damage the fruit, but also damage the shoots and young leaves due to larval boring and feeding. This results in a large number of dropped fruits ("dung fruits"), fallen flowers and dieback (FAO 2002). The adult moths become active soon after sunset (18:00–18:30 h), although a few moths may become active as early as 16:30 h on dark rainy d. Adult moths

cease to be active near the dawn, and they remain quiescent during the photophase (Cai et al. 2011). As the litchi stem-end borer larvae are interval feeders, they are difficult to control with chemical pesticides. Also, in organic litchi production, most chemicals are strictly prohibited, which makes control by traditional methods very difficult (Tsang et al. 2007, 2011).

The sterile insect technique (SIT) is a method of pest control based on area-wide releases of sterile males in a ratio of sterile to wild males of the same species that is sufficiently great to drasti-

Department of Plant Protection, South China Agricultural University, Guangzhou, 510642 China

*Corresponding author; E-mail: humy@scau.edu.cn; huabao@scau.edu.cn

Copyright © International Atomic Energy Agency 2016. Published by the Florida Entomological Society. All rights reserved.

cally reduce the fertility of the targeted field population. The SIT depends greatly on the production of good quality sterile males that can locate and compete for wild females of the wild target population. Since the New World screwworm, *Cochliomyia hominivorax* (Coquerel) (Diptera: Calliphoridae) was eradicated from the Island of Curacao in 1954 using this technology, the SIT has received increased attention.

Many lepidopteran species are destructive pests of annual and perennial crops, forests, and stored products throughout the world. Because lepidopteran species are usually more radiation resistant than most other insect species (Dyck et al. 2005), a fully sterilizing radiation dose would be so great that it would reduce the quality of the irradiated males and reduce their ability to disperse in the field and compete for wild females. To increase the competitiveness of irradiated Lepidoptera, Proverbs (1962) investigated the potential of using sub-sterilizing doses of radiation on the codling moth, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae), and discovered that at less than fully sterilizing doses, the sons of irradiated males were fully sterile. This discovery of inherited (IS) or F_1 sterility promoted numerous investigations in many Lepidopteran pests.

Inherited sterility is a derivative of the SIT that is applicable to lepidopteran pests (Proverbs 1962; LaChance 1985; Carpenter et al. 2005; Kean et al. 2008), and—when compared with full sterility, provides several advantages for lepidopteran pest control. As explained by Carpenter et al. (2005), lepidopteran females are usually more radio-sensitive than males of the same species. Therefore in many cases it is possible to find a radiation dose that induces complete sterility in females but only partial sterility in males. Thus large numbers of irradiated males and females can be released in the field, but the released females cannot produce progeny that can damage the crop. However, when the partially sterile parental generation mates with wild females the radiation induced deleterious genes are inherited by the F_1 generation. As a result the F_1 egg hatch is reduced and the F_1 offspring tend to be both sterile and predominantly male. Because the F_1 progeny are produced in the field they tend to be better adapted and distributed to compete for wild mates. The advantages of IS derive from the lowering of the radiation dose, which allows the release of more competitive and longer living males. In addition adult F_1 males tend to be stronger fliers that mate more frequently and are more competitive than moths irradiated with greater radiation doses (Carpenter 2000).

In a previous study, females irradiated with 200 Gy did not oviposit when mated with non-irradiated males (Fu et al. 2016), indicating that 200 Gy was the lowest fully sterilizing dose for females, and 200 Gy was confirmed to be the dose needed to induce complete sterility in the F_1 generation offspring from the parental cross untreated female (UF) × treated male (TM). Radiation-induced partial sterility of the litchi stem-end borer was shown to be inherited in at least 3 consecutive generations beginning with the parental males that were irradiated with 150 Gy as pupae shortly before emergence. There were no significant differences in longevity between the parental generation and the F_1 generation when the male in the parental generation was irradiated with 150 Gy. Therefore, 150 Gy applied before emergence could be used to irradiate moths for large scale release programs, although a dose of 200 Gy may be more appropriate for use in SIT/IS programs. Very little work has been done on quality control aspects of *C. sinensis*, however, many workers have reported on various quality parameters used for other lepidopteran pests (Rogers & Winks 1993; Bloem et al. 1997; Vreysen & Hendrichs 2005). This study was carried out to assess the effect 150 and 200 Gy irradiation on the flight ability of male litchi stem-end borers in the laboratory and to assess their dispersal ability in the field.

Materials and Methods

SOURCE OF INSECTS AND PROCEDURE FOR MARKING MALES WITH DIFFERENT FLUORESCENT DYES

The litchi stem-end borers were collected from litchi orchards at the Guangdong Academy of Agricultural Sciences, Hainan, Maoming and Zengcheng, China. We gathered infested fallen fruit and collected the pupae from leaves on trees and fallen leaves. These insects were used to establish a laboratory colony (according to the procedure described by Fu et al. (2016) in this issue) that provided moths for the experiments.

After irradiation the male adult moths were marked with 1 of 3 fluorescent dyes (Kunyou Illumination Technology Co., Ltd. Jinan, China) to distinguish non-irradiated from irradiated males after recapture in the release-recapture experiment. The effect of the fluorescent dye on the longevity of adult male moths was first assessed. One d old adult males were marked with 3 different fluorescent dyes (blue, orange and green) with 30 males in each color treatment. Adult males without dye were used as the control. Their longevities were observed and recorded.

RADIATION SOURCE

The experimental material was irradiated in a ^{60}Co source (Nordion Inc., Ottawa, Ontario, Canada) located at the Guangzhou Furui High-Energy Technology Co., Nansha District, Guangzhou, Guangdong Province, China. The dose rate was 3.2 Gy/min.

FLIGHT MILL AND TETHERED FLIGHT

The flight performance of *C. sinensis* adults was tested using a 26-channel, computer monitored flight mill system at the Institute of Plant Protection (Zhengzhou, China). Mature pupae of uniform size were irradiated with either 150 or 200 Gy. Adult males which emerged from these irradiated pupae were anesthetized with CO_2 for 30 s when 1 d old. By means of a drop of cyanoacrylate Super Glue (Yuyao Kexing Adhesive Co., Zhejiang, China) placed on the ventral side of the mesothorax, each male was attached to the end of the flight mill arm. The flight mill was placed in a dark room at $30 \pm 2^\circ\text{C}$ and 75–85% RH. One tethered flight test took 24 h to complete, i.e., from 21.00 h to 21.00 h of the next day.

The software system recorded the time of flight initiation and cessation, as well as the number of mill revolutions every 5 s during the assay period. If flight stopped for ≥ 1 min, it was considered the end of a flight. Based on the number of mill revolutions over a given time period, total flight distance and duration, average speed, as well as distance and duration of each flight were computed for each individual using a custom-made software package (Cheng et al. 1997).

WEATHER CONDITIONS DURING FIELD RELEASE STUDIES

Weather data during the release trials were obtained from a station of the Guangdong Meteorological Service 5 km from the release orchard.

RELEASE AND RECAPTURE EXPERIMENT

Irradiation and Marking

Mature pupae of uniform size were irradiated with either 150 or 200 Gy. Each of the adult males that emerged from these irradiated pupae was dyed when 1 d old with 1 of the 3 different fluorescent dyes

to identify the 3 treatments (non-irradiated control was blue, 150 Gy was orange, and 200 Gy was green).

Sex Pheromone Bait and Triangle Traps

The sex pheromone used to bait the traps and the triangle traps were provided by Zhangzhou Enjoy Agriculture Technology Co., Ltd., China. The traps were open on both sides and had an adhesive cardboard insert on the bottom of the trap to capture adult males that entered the trap. The sex pheromone attractant was placed in a small plastic cup inside the trap.

Release and Recapture

Two release-recapture experiments were conducted to assess the dispersal ability of adult males. The releases were carried out in a litchi orchard of the South China Agricultural University, Guangzhou, Guangdong Province, China. The litchi orchard had a rectangular shape with dimensions of 60 m (E-W) × 160 m (S-N). The release point was located at the intersection of the 2 diagonals of the orchard rectangle. Twenty-four traps were deployed and hung in litchi trees and aligned according to 8 compass directions—with radial angles of 45°—at 5, 15 and 30 m from the release point. An additional 6 traps were deployed at 80 m distances to the S, N, SW, SE, NW and NE. The release point of adult males, the position of each trap, and the number of traps were the same throughout the experimental period.

The first experiment was conducted from 3 to 11 Jul, and the second from 20 to 28 Jul 2013. In each experiment a total of 500 dyed male adults of each treatment (150 and 200 Gy) and of the non-irradiated control were released into the orchard around 16:30–17:00 h. During the 5 d after the release, the triangle traps were checked every 24 h. The adhesive cardboard base was removed and replaced with a new one if moths were trapped and the cardboard base was taken to the laboratory to record the catch. Normally the color of dye on the surface of the captured male could be readily determined with the aid of a UV lamp.

DATA ANALYSIS

For each individual *C. sinensis* male, flight activity was characterized by total flight distance, flight duration, average flight speed, number of flights, mean duration of flights, and distance and duration of the longest flight. Flight parameters were analyzed using a two-way analysis of variance (ANOVA), with wind direction, temperature, relative humidity, and dye as factors. Flight data of individual tests were analyzed and calculated using the Matlab software package (MathWorks, Inc., Natick, Massachusetts, USA). All other data were analyzed with SPSS Version 19.0 (SPSS Inc., Chicago, Illinois, USA) software. The meteorological data were presented as mean values for each 24 h period.

Results

INFLUENCE OF IRRADIATION ON FLIGHT ABILITY OF ADULT MALE LITCHI STEM-END BORERS

The flight distance, flight time, mean speed, and the greatest speed of non-irradiated adult litchi stem-end borer males were 13,926 m, 29,365 s, 0.42 m/s, and 1.01 m/s, respectively, during a 24 h period using the flight mill system in the laboratory. The values of each of these parameters were greater for the non-irradiated control males as than for the irradiated males, but the differences were not significantly different (Table 1). The flight mill data did not reveal any significant physiological deterioration as the result of either the 150 or 200 Gy treatment on the flight parameters of adult litchi stem-end borer males.

EFFECT OF FLUORESCENT DYES ON LONGEVITY OF ADULT MALE LITCHI STEM-END BORERS

The observed mean longevity of each treatment was in the range of 13–15 d (Table 2). Analyses using the proportional hazard model indicated that none of the dyes significantly increased the hazard of dying (Table 2) in comparison with the unmarked control. Even after 11 days, the survival rates in all treatments were greater than 90%, but survival in all treatments declined steeply soon thereafter (Fig. 1).

RELEASE AND RECAPTURE EXPERIMENTS

Recaptures of *Conopomorpha sinensis* Males

Weather conditions during the first release experiment were favorable with almost no precipitation, and steady light winds (1.6–1.9 m/s) mostly from the SE (Table 3) under a mostly clear sky. In the first release experiment, a total of 1,500 adult male moths were released, and a total of 87 males (5.8%) were recaptured. The recapture rates of the non-irradiated control males, and males irradiated with either 150 or 200 Gy in the first release experiment were 6.2%, 5.4%, and 5.8%, respectively. The irradiation treatments (150 and 200 Gy) had no significant effect on the recapture rates as compared with the non-irradiated control ($\chi^2 = 0.29$; $P = 0.86$) (Table 4).

Weather conditions during the second release experiment were somewhat turbulent with precipitation on all 5 d of the experiment, and light but variable winds (0.8–2.0 m/s) of quite variable directions fluctuating mostly between SE and NE under a mostly cloudy sky (Table 3). In the second release experiment, a total of 1,500 males were released, and a total of 123 males (8.2%) were recaptured. The results were similar to those of the first release in that no significant differences were found in recapture rates of the males irradiated either with either 150 or 200 Gy and the non-irradiated control males ($\chi^2 = 1.49$; $P = 0.48$) (Table 4).

Table 1. Flight parameters of *Conopomorpha sinensis* males tethered on a flight mill system. The males were non-irradiated or irradiated either with 150 or 200 Gy and allowed to fly for 24 h. If a flight stopped for ≥ 1 min, it was considered the end of that flight.

Radiation dose (Gy)	Number of tested insects	Flight distance \pm SE (m)	Flight duration \pm SE (s)	Mean flight speed \pm SE (m/s)	Maximum speed \pm SE (m/s)
0 (control)	8	13,926 \pm 1,419 a	29,365 \pm 2,970 a	0.4175 \pm 0.026 a	1.0075 \pm 0.067 a
150	8	11,732 \pm 2,254 a	24,934 \pm 3,636 a	0.4073 \pm 0.019 a	1.0119 \pm 0.064a
200	8	9,257 \pm 1,116 a	26,328 \pm 2,180 a	0.4088 \pm 0.022 a	0.656 \pm 0.019 a

Note: Mean values in the same column followed by the same letters are not significantly different at $P = 0.05$ level according to the Duncan's multiple ranger test (DMRT)

Table 2. The influence of fluorescent dyes on longevity in the laboratory of adult *Conopomorpha sinensis* males.

Color of fluorescent dye	Mean longevity (days) \pm SE	Proportional hazard model			
		T _i	SE	Wald chi-square	P
None (control)	14.1 \pm 1.6	—	—	—	—
Orange	14.2 \pm 2.0	-0.045	0.366	0.015	0.902
Blue	13.7 \pm 2.1	-0.130	0.366	0.126	0.723
Green	13.9 \pm 2.0	0.062	0.366	0.029	0.865

Dispersal Direction and Dispersal Distance

The maximum distances that the adult males dispersed were 30 m and 80 m in the first and second release experiments, respectively.

Most of the recaptured males (67.8% in the first experiment and 62.8% in the second experiment) were caught in traps located only 5 m from the release point. The number of recaptured males decreased exponentially with distance from the release point (Fig. 2).

A Kruskal-Wallis test indicated that there were no significant differences in the dispersal distances of males of the 2 treatment groups as compared with the non-irradiated control males in the first ($W = 0.00$, $P = 1$) and second experiment ($W = 0.2$, $P = 0.905$). In view that no effect of irradiation could be detected with respect to recapture rate and dispersal distance, all data in each experiment was analyzed by regression analysis. The analysis indicated a median dispersal distance of 11.3 m and an arithmetic mean of dispersal distance of 8.7 m in the first release experiment. In the second experiment the median dispersal distance was estimated at 10.6 m and the arithmetic mean of the dispersal distance was estimated at 15.6 m (Table 4).




Fig. 1. Survival (days) of non-irradiated *Conopomorpha sinensis* males either dyed with 1 of 3 different fluorescent colors or undyed (control) in the laboratory. Each treatment involved 30 males.

Table 3. The weather conditions during the field release experiments with *Conopomorpha sinensis* adult males in a litchi orchard of the South China Agricultural University, Guangzhou, China. Meteorological data were obtained from a station of the Guangdong Meteorological Service located 5 km from the release orchard. The data presented are the mean values for each 24 h period.

	Day of release				
	1st	2nd	3rd	4th	5th
First Experiment					
Mean temperature (°C)	28.8	28.9	29.3	30.2	29.0
Max/min temperature (°C)	34.0/24.9	35.0/26.2	34.8/26.9	34.6/27.2	33.2/26.8
Dominant wind direction	SE	SE	SSE	SE	SSE
Mean speed of wind (m/s)	1.6	1.9	1.6	1.8	1.9
Rainfall (mm/d)	0.1	0.0	0.0	0.0	0.0
Second Experiment					
Mean temperature (°C)	27.1	26.7	27.7	28.5	28.8
Max/min, temperature (°C)	32.1/24.5	31.0/24.2	33.7/23.3	35.2/25.0	35.0/25.2
Dominant wind direction	SE	ESE	SE	NE	ENE
Mean speed of wind (m/s)	2.0	1.5	0.8	0.9	1.0
Rainfall (mm/d)	16.6	18.5	0.2	8.6	0.8

Litchi stem-end borer males dispersed in all directions, but in both experiments most males were recaptured in 3 directions (S, W and SW) (Fig. 3). These dispersal directions were similar to the wind directions in each experimental period (Table 3). The dispersal direction was not significantly different for the males of the 2 treatment groups as compared with the non-irradiated control males; for the first release ($P = 0.50$) and for the second release ($P = 0.80$). Analyzed percentages of recaptured males in each treatment and experiment were significantly different among the 8 directions (Table 5).

Discussion

Our study showed an average flight speed of non-irradiated male litchi stem-end borers of 0.42 m/s, as measured in a flight mill system in the laboratory. This recorded speed was lower than the one

observed for 1 d old Asian corn borers, *Ostrinia furnacalis* (Guenée) (Lepidoptera: Crambidae), i.e., 1.01 m/s at 24 °C and 70–75% RH (Lu et al. 2005), but similar to the one observed for 1 d old pink bollworms, *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae), i.e., 0.54 m/s at 28 °C and 80% RH (Wu 2006). In addition, the greatest total flight distance flown by a *C. sinensis* male using the flight mill system was 13.93 km.

The flight ability of released sterile insects is an important factor for the success of the SIT/IS. Although we cannot expect that the flight performance of insects with a flight mill will accurately reflect their performance under field conditions, the flight mill does provide standardized and quantitative estimates of the insect's intrinsic flight capacity (Taylor et al. 1992). Within the known limitations of the system (Armes & Cooter 1991; Beerwinkle et al. 1995; Riley et al. 1997), flight mills have been used to examine insect flight activity as a function of age, sex, size, mating status, and physiological state (Colvin & Gatehouse

Table 4. Analysis of parameters of field release experiments with adult *Conopomorpha sinensis* males in a litchi orchard of the South China Agricultural University, Guangzhou, China.

	First experiment			Second experiment		
	0	150	200	0	150	200
Dose (Gy)						
No. of males released	500	500	500	500	500	500
No. of males recaptured	31	27	29	39	47	37
Recapture rate (%)	6.2	5.4	5.8	7.8	9.4	7.4
Effect of irradiation on the recapture rate ^a	$\chi^2 = 0.293$ df = 2 $P = 0.864$			$\chi^2 = 1.488$ df = 2 $P = 0.475$		
Arithmetic mean of dispersal distance(m) ^b	9.2	8.89	8.10	14.10	15.96	16.49
Effect of irradiation on the dispersal ability ^c	$W = 0.000$ $P = 1$			$W = 0.2$ $P = 0.905$		
Regression coefficient ^d	-1.283			-2.964		
P^e	0.000			0.002		
Correlation coefficient (r)	0.965			0.788		
Estimates of mean dispersal distance (m) ^f	11.33			10.61		

^aComparison of the recapture rates across 3 treatments; Chi-square test

^bSum of dispersal distance of every recaptured male per number of recaptured males

^cComparison of the dispersal distances across 3 treatments; Kruskal-Wallis test

^dRegression analysis of the number of male individuals on distance from released point

^eTest of significance for regression; P is probability

^fMedian dispersal distance calculated by the regression




Fig. 2. Numbers of adult *Conopomorpha sinensis* males recaptured in traps deployed at various distances (m) from the release point in 2 release/recapture experiments in a litchi orchard of the South China Agricultural University, Guangzhou, China. The males were either non-irradiated or irradiated either with 150 or 200 Gy. A: First release, B: Second release.

1993; Rankin et al. 1994; Lu et al. 2007; Sarvary et al. 2008; Tsang et al. 2011), as well as lipid utilization in flight (Williams & Robertson 2008).

Irradiation of adult litchi stem-end borer males with either 150 or 200 Gy did not impair their flight ability as measured with a flight mill. Bloem et al. (2006) reported that the mobility of adult *C. pomonella* males was significantly influenced by an interaction of radiation and ambient temperature, but that the relatively low dose of 150 Gy had little effect on the flight activity of the codling moths. In contrast wind tunnel studies with the light brown apple moth, *Epiphyas postvittana* Walker (Lepidoptera: Tortricidae), indicated that either 100 or 250 Gy reduced upwind flight by 14% and 46%, respectively (Suckling et al. 2011). Previous studies had shown that relatively low doses of gamma irradiation did not significantly affect the flight ability of *C. sinensis*, and the results of this study are consistent with the previous studies.

Li et al. (2009) reported that 80.5% of litchi stem-end borer adults in 1 experiment flew 3–7 m, with an average flight distance of 4.97 m and a maximum distance of 12 m. These data are similar to the results of our release recapture experiment, where 59.1% of the released

adult males were recaptured in traps that were located at 5 m from the release point, and the median dispersal distance as obtained with the regression analysis was about 11 m. However, we need to recognize that the released males were recaptured in traps baited with the female's sex pheromone, and it is very likely that the released males would have dispersed much longer distances if pheromone sources had been absent.

In the 2 release experiments, the main dispersal direction of the adult male litchi stem-end borers corresponded with the main wind direction, and it was skewed toward the west. We hypothesize that there are 2 other possible factors that may have affected the results. First, our study site was located on an east-west slope, and the experimental area was surrounded by trees except on the north-eastern side. This topography may have enhanced the effect of the wind. Second, the odor of vegetation including oleander, *Nerium indicum* Mill. (Gentianales: Apocynaceae) and wax apple, *Syzygium samarangense* Merr. & Perry (Myrtales: Myrtaceae) outside of the study area may have affected the dispersal direction. Generally, these factors—as well as the effect of some artificial lights to which insects may display a negative




Fig. 3. Frequency distributions of the dispersal directions of adult *Conopomorpha sinensis* males in 2 release/recapture experiments in a litchi orchard of the South China Agricultural University, Guangzhou, China. The males were either non-irradiated or irradiated either with 150 or 200 Gy. A: First release, B: Second release.

Table 5. Tests of dispersal direction of adult *Conopomorpha sinensis* males in field release experiments in a litchi orchard of the South China Agricultural University, Guangzhou, China.

	Test of dispersal direction			Comparison of dispersal direction across 3 treatments
	0 Gy	150 Gy	200 Gy	
First experiment	$P < 0.01^b$	$P < 0.01^b$	$P = 0.009^b$	$P = 0.496^b$
Second experiment	$P < 0.01^b$	$\chi^2 = 37.708, P = 0 \text{ df} = 7^a$	$P < 0.01^b$	$P = 0.800^b$

^aChi-square test^bFisher exact test

phototaxis (Li et al. 2008)—may combine to affect the dispersal direction (Kumano et al. 2009).

In the present study, we demonstrated that irradiating adult litchi stem-end borer males with either 150 or 200 Gy did not affect their flight ability as measured with a flight mill in the laboratory, nor did these levels of gamma irradiation affect their dispersal ability in the field as measured in 2 release recapture studies. It is clear that operational and commercial insect pest management programs with a SIT component can only be successful when there is consistent production and release of sterile insects of high biological quality and performance (Calkins & Parker 2005). Effective quality assurance tests for monitoring and providing feedback on the performance of insects during each step of production, handling and release are crucial to insure quality control and success of these programs (Huettel 1976; Singh & Ashby 1985; Calkins & Parker 2005). Additional studies are needed to evaluate the quality and performance of litchi stem-end borers released in the field, and to develop mass rearing and mass handling procedures.

Acknowledgments

This work was part of the FAO/IAEA Coordinated Research Project on Increasing the Efficiency of Lepidoptera SIT by Enhanced Quality Control. The researchers gratefully acknowledge the grants from the International Atomic Energy Agency under Research Contract No. 15059/RQ, as well as the invaluable support of the Department of Plant Protection, South China Agricultural University.

References Cited

- Armes NJ, Cooter RJ. 1991. Effects of age and mated status on flight potential of *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Physiological Entomology* 16: 131-144.
- Beerwinkle KR, Lopez JD, Cheng D, Lingren PD, Meola RW. 1995. Flight potential of feral *Helicoverpa zea* (Lepidoptera: Noctuidae) males measured with a 32-channel, computer-monitored, flight-mill system. *Environmental Entomology* 24: 1122-1130.
- Bloem S, Bloem KA, Fielding LS. 1997. Mass-rearing and storing codling moth larvae in diapause: A novel approach to increase production for sterile insect release. *Journal of the Entomological Society of British Columbia* 94: 75-81.
- Bloem S, Carpenter JE, Dorn S. 2006. Mobility of mass-reared diapaused and nondiapaused *Cydia pomonella* (Lepidoptera: Tortricidae): Effect of mating status and treatment with gamma radiation. *Journal of Economic Entomology* 99(3): 699-706.
- Cai ST, Qi XY, Lin GC, Deng XY. 2011. The meteorological forecast method of rating damage caused by *Conopomorpha sinensis* (Lepidoptera: Gracillariidae) in Gaozhou City. *Guangdong Meteorology* 33(3): 46-48. In Chinese.
- Calkins CO, Parker AG. 2005. Sterile insect quality, pp. 269-296 *In* Dyck VA, Hendrichs J, and A.S. Robinson [eds.], *Sterile Insect Technique. Principles and Practice in Area-Wide Integrated Pest Management*. Springer, Dordrecht, The Netherlands.
- Carpenter JE. 2000. Area-wide integration of lepidopteran F_1 sterility and augmentative biological control, pp. 193-200 *In* Tan KH [ed.], *Area-wide control of fruit flies and other insect pests*. *Proceedings International Conference on Area-Wide Control of Insect Pests, and the 5th International Symposium on Fruit Flies of Economic Importance, 28 May-5 June 1998, Penang, Malaysia*. Penerbit Universiti Sains Malaysia, Pulau Pinang, Malaysia.
- Carpenter JE, Bloem S, Marec F. 2005. Inherited sterility in insects, pp. 115-146 *In* Dyck VA, Hendrichs J, Robinson AS [eds.], *Sterile Insect Technique. Principles and Practice in Area-Wide Integrated Pest Management*. Springer, Dordrecht, The Netherlands.
- Cheng D, Tian Z, Sun J, Ni H, Li G. 1997. A computer monitored flight mill system for tiny insects such as aphids. *Acta Entomologica Sinica* 40: 172-179. In Chinese.
- Colvin J, Gatehouse AG. 1993. Migration and genetic regulation of the pre-reproductive period in the cotton bollworm moth, *Helicoverpa armigera*. *Heredity* 70: 407-412.
- Dyck VA, Hendrichs J, Robinson AS [eds.]. 2005. *Sterile Insect Technique*. Springer, Dordrecht, The Netherlands.
- Food and Agriculture Organization of the United Nation (FAO). 2002. Lychee production in the Asia-Pacific region. Rome, Italy. pp. 9-13.
- Fu H, Zhu F-W, Deng Y-Y, Weng Q-F, Hu M-Y, Zhang T-Z. 2016. Development, reproduction and sexual competitiveness of *Conopomorpha sinensis* (Lepidoptera: Gracillariidae) gamma-irradiated as pupae and adults. *Florida Entomologist* 99 (special issue #1): 66-72.
- Huettel MD. 1976. Monitoring the quality of mass-reared insects: A biological and behavioral perspective. *Environmental Entomology* 5: 807-814.
- Kean JM, Wee SL, Stephens AEA, Suckling DM. 2008. Modelling the effects of inherited sterility for application of the sterile insect technique. *Agricultural and Forest Entomology* 10: 101-110.
- Kumano N, Kawamura F, Haraguchi D, Kohama T. 2009. Irradiation does not affect field dispersal ability in the West Indian sweet potato weevil, *Euscepes postfasciatus*. *Entomologia Experimentalis et Applicata* 130: 63-72.
- LaChance LE. 1985. Genetic methods for the control of lepidopteran species. United States Department of Agriculture-Agricultural Research Service, Series 28. Washington DC. 40 pp.
- Li ZQ, Qiu YP, Ou LX, Chen JZ, Xiang X, Sun QM. 2008. Survey of damage caused by the litchi stem-end borer in longan orchards in various environments. *Guangdong Agricultural Sciences* 12: 93-94. In Chinese.
- Li ZQ, Qiu YP, Ou LX, Chen JZ, Xiang X, Sun QM. 2009. Observations on the flight range of the litchi stem-end borer and its quiescent habits indoors. *South China Fruits* 38(2): 54-55. In Chinese.
- Lu X, Tian ZL, Zhang GH, Li L. 2005. Research on the flight capability and effective accumulated temperature of different voltine ecotypes of Asian corn borer. *Acta Phytophylactica Sinica* 32(3): 333-334. In Chinese.
- Lu Y, Wu K, Guo Y. 2007. Flight potential of the green plant bug, *Lygus lucorum* (Meyer-Dur) (Heteroptera: Miridae). *Environmental Entomology* 36: 1007-1013.
- Proverbs MD. 1962. Progress on the use of induced sexual sterility for the control of the codling moth *Carpocapsa pomonella* (L.) (Lepidoptera: Olethreutidae). *Proceedings of the Entomological Society of Ontario* 92: 5-11.
- Rankin MA, Hampton EN, Summy KR. 1994. Investigations of the oogenesis-flight syndrome in *Anthonomus grandis* (Coleoptera: Curculionidae) using tethered flight tests. *Journal of Insect Behavior* 7: 795-810.
- Riley JR, Downham MCA, Cooter RJ. 1997. Comparison of the performance of *Cicadulina* leafhoppers on flight mills with that to be expected in free flight. *Entomologia Experimentalis et Applicata* 83: 317-322.
- Rogers DL Winks CJ. 1993. Quality control in laboratory-reared codling moth at Mt. Albert Research Centre, Auckland, New Zealand, pp.13-21 *In* Nicoli G, Benuzzi M, Leppla NC [eds.], *Proceedings of the 7th Workshop of the IOBC Global Working Group "Quality Control of Mass Reared Arthropods"*, 13-16 September 1993, Rimini, Italy (available at <http://users.ugent.be/~padclerc/AMRQC/proceedings.html>).
- Sarvary MA, Bloem KA, Bloem S, Carpenter JE, Hight SD, Dorn S. 2008. Diel flight pattern and flight performance of *Cactoblastis cactorum* (Lepidoptera: Pyralidae) measured on a flight mill: Influence of age, gender, mating status, and body size. *Journal of Economic Entomology* 101: 314-324.

- Singh P, Ashby MD. 1985. Insect rearing management, pp.185-215 In Singh P, Moore RF [eds.], Handbook of Insect Rearing, Vol. I, Elsevier, Amsterdam, The Netherlands.
- Suckling DM, Stringer LD, Mitchell VJ, Sullivan TES, Sullivan NJ, Simmons GS, Barrington AM, El-Sayed AM. 2011. Comparative fitness of irradiated light brown apple moths (Lepidoptera: Tortricidae) in a wind tunnel, hedgerow, and vineyard. *Journal of Economic Entomology* 104(4): 1301-1308.
- Taylor RAJ, Nault LR, Styer WE, Cheng Z. 1992. Computer-monitored, 16-channel flight mill for recording the flight of leafhoppers (Homoptera: Auchenorrhyncha). *Annals of the Entomological Society of America* 85: 627-632.
- Tsang WS, Liang GW, Liu WH, Chen QX. 2007. The new record of selecting effective species of egg parasitoids of *Conopomorpha sinensis* Bradley (Lepidoptera: Gracillariidae). *Natural Enemies of Insects* 29(1): 6-11.
- Tsang W, You LS, Achterberg C van, Liang GW. 2011. A new species of *Phanerotoma* Wesmael (Hymenoptera: Braconidae: Cheloniinae), a parasitoid of *Conopomorpha sinensis* Bradley (Lepidoptera: Gracillariidae) from South China. *Zootaxa* 28: 53-58.
- Vreysen MJB, Hendrichs J. 2005. The potential of integrating the sterile insect technique as an environmentally friendly method for area-wide management of the codling moth (*Cydia pomonella*), pp. 56-62 In Cross J, Ioriatti C [eds.], Integrated Fruit Protection in Fruit Crops and Use of Pheromones and other Semiochemicals in Integrated Control. IOBC WPRS Bulletin 28.
- Williams WI, Robertson IC. 2008. Using automated flight mills to manipulate fat reserves in Douglas-fir beetles (Coleoptera: Curculionidae). *Environmental Entomology* 37: 850-856.
- Wu HS. 2006. Studies on flight potential and its relationship with reproduction in pink bollworm (*Pectinophora gossypiella* Saunders). Agricultural College of Xinjiang Agricultural University, 32-39. In Chinese.