



Acrotomopus atropunctellus (Coleoptera: Curculionidae) Preference for Large Sugarcane Shoots Mitigates Damage to Sugarcane Crop

Authors: Pérez, María L. del P., Isas, Marcos G., Salvatore, Analía R., García, José M., and Trumper, Eduardo V.

Source: Florida Entomologist, 100(3) : 678-679

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.100.0332>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Acrotomopus atropunctellus (Coleoptera: Curculionidae) preference for large sugarcane shoots mitigates damage to sugarcane crop

María L. del P. Pérez^{1,*}, Marcos G. Isas¹, Analía R. Salvatore², José M. García³, and Eduardo V. Trumper⁴

The sugarcane stem weevil, *Acrotomopus atropunctellus* (Boheman) (Coleoptera: Curculionidae), affects sugarcane in the principal Argentinian sugarcane-producing area (Pérez et al. 2016). Although its damage potential has been known since at least 1929 (Box 1929), significant crop injury was not observed until 2003 (Salvatore et al. 2006). The oviposition and feeding punctures made by the adults, and the tunnels made by the larvae, cause the death of sugarcane shoots (Pérez et al. 2012a).

Although the production of shoots is constant across the whole sugarcane cycle, the proportion of small shoots is highest in the early phenological stages, mainly the sprouting and tillering phases (according to the classification of Romero 2002). Preliminary studies indicated that the *A. atropunctellus* emergence period overlaps with early and middle phenological stages (tillering and rapid growth) (Pérez et al. 2012b). The objective of this research was to assess the food preference of *A. atropunctellus* in relation to shoot size and to determine the effect of damage on shoot survival.

Adults used for experiments were collected from emergence cages placed in commercial sugarcane fields planted with variety LCP 85-384 located in Ranchillos (26.9847°S, 65.0100°W), Tucumán, Argentina. Only individuals emerging within the same week were used in each experiment.

Using forest soil as substrate, 300 vegetative seed pieces of sugarcane (each 10 cm long and bearing 1 lateral bud) of variety LCP 85-384, the dominant variety in this area (Ostengo et al. 2014), were planted in 250 cm³ plastic cups, in batches of 100 cups, at 2 wk intervals to obtain shoots of different sizes. The cups were placed in a greenhouse under ideal conditions for seedling development (21–35 °C).

The shoots were classified according to diameter and height. The diameter was measured at the mid-point of its length, which in turn was measured from soil level to the leaf +1 ligule. Ranges were 5 to 7 mm wide and 5 to 11 cm long (small), 7 to 9 mm wide and 12 to 15 cm long (medium), and 10 to 12 mm wide and greater than 15 cm long (large). These size categories represent the most common sizes of shoots in the early phenological stages of sugarcane (Romero 2002).

The experimental unit consisted of an 8 L pot containing 3 shoots (1 shoot per size category) covered by a voile cage and forest soil as substrate. To determine feeding preference, 8 experimental unit were infested with 1 male and 8 were infested with 1 female (total $n = 16$).

After 5 d, the damage was assessed through the number of punctures on shoots.

A second experiment was performed to study the impact of *A. atropunctellus* on survival of sugarcane shoots. The experimental unit consisted of a 20 L pot containing 2 shoots in the same size category covered by a voile cage. Only medium and large size categories were used because 100% of small size shoots died in a preliminary experiment. Pots were infested with 0, 1 (female), 2 (1 couple), 4 (2 couples), 6 (3 couples), and 9 weevils (4 couple plus 1 female), with 3 replicates per weevil density. After 2 wk, the numbers of dead shoots were counted.

Linear mixed effects models were used (Pinheiro & Bates 2000), and the analyses were carried out in Infostat R interface (Di Rienzo et al. 2009) with pot as a random effect and size category of shoots, sex of weevil, their interaction, and the density of weevil as fixed effects. Heteroscedasticity was modeled using a constant variance function (varIdent function, package nlme) (Pinheiro et al. 2015). Several models were fitted to the data. Means were compared with Fisher's least significant difference (LSD) test.

The model that included shoot size, sex, and their interaction as fixed effects and pot as random effect was selected. *Acrotomopus atropunctellus* preferred to feed on large and medium shoots instead of small ones (Fig. 1). Significant effects of sex ($F = 7.39$; $df = 1$; $P = 0.016$)




Fig. 1. Mean numbers (\pm SE) of feeding punctures made by *Acrotomopus atropunctellus* on sugarcane shoots of different sizes. Different letters represent significant differences between means ($P < 0.05$) (Fisher's LSD test).

¹Instituto de Tecnología Agroindustrial del Noroeste Argentino (ITANOVA), Estación Experimental Agroindustrial Obispo Colombres (EEAOC)-Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Av. William Cross 3150, C.P. T4101XAC, Las Talitas, Tucumán, Argentina; E-mail: mlperez@eeaoc.org (M. L. P. P.); mgisas@hotmail.com (M. G. I.)

²Facultad de Agronomía y Zootecnia, Universidad Nacional de Tucumán, Avenida Néstor Kirchner 1900, Argentina; E-mail: asalvatore@hotmail.com (A. R. S.)

³Estación Experimental Agroindustrial Obispo Colombres (EEAOC), Las Talitas, Tucumán, Argentina; E-mail: joegarcía_220@hotmail.com (J. M. G.)

⁴Instituto Nacional de Tecnología Agropecuaria (INTA), Ruta Nac. Nro. 9 km 636 (C.P. 5988), Manfredi, Córdoba, Argentina; E-mail: eduardo.trumper@gmail.com (E. V. T.)

*Corresponding author; E-mail: mlperez@eeaoc.org.ar (M. L. P. P.)

and shoot size ($F = 9.23$; $df = 2$; $P = 0.001$) on the number of punctures were identified. Size by sex interaction was not significant ($F = 0.98$; $df = 2$; $P = 0.387$) indicating that shoot preference does not vary according to sex. However, the number of punctures made by females (10.4 ± 1.2) was 1.6 times greater than that produced by males (6.3 ± 0.9). This behavior may be due to the different nutritional demands associated with reproduction and oviposition (Behmer & Joern 1994).

In shoots of medium size, shoot mortality with 1 weevil per pot was 15% whereas densities of 2 weevils per pot and higher caused shoot mortalities of 100% ($P = 0.001$). In large shoots, only densities of 9 weevils per pot caused mortality (25%) ($P = 0.001$).

Although large- and medium-sized shoots were preferred by *A. atropunctellus* adults, the low numbers of punctures found on small shoots were enough to cause their death. Considering the susceptibility of small shoots, this pest would cause greater impact in early than in late phenological stages.

Sugarcane planting and harvesting dates affect the emergence dynamics of shoots. In Tucumán, these operations are simultaneously conducted from May to Nov (Digonzelli et al. 2005; Romero et al. 2005). Despite its negative impact on yield (Digonzelli et al. 2006), producers sometime use late planting dates (Oct–Nov) due to machinery availability. The manipulation of sowing or planting dates to avoid the main pulse of pest emergence or immigration is a widely used tactic in integrated pest management (Ghosh 1989). Beuzelin et al. (2011) recommended this management option to protect sugarcane yields against *Diatraea saccharalis* F. sensu Guenée (Lepidoptera: Crambidae). Avoidance of late planting or harvest dates could be a useful tactic for mitigating *A. atropunctellus* incidence in sugarcane crops because it would avoid the overlap of crop sprouting or ratooning and emergence of *A. atropunctellus* adults.

This study was supported by Estacion Experimental Agro-industrial Obispo Colombes (EEAOC) and Instituto de Tecnología Agroindustrial del Noroeste Argentino (ITANOA) of Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

Summary

The sugarcane weevil, *Acrotomopus atropunctellus* (Boheman) (Coleoptera: Curculionidae) is an increasingly important sugarcane pest in Argentina. Its emergence period overlaps with early stages of sugarcane phenology. This research assessed the preference of *A. atropunctellus* for shoot size and the effect of damage on shoot survival. *Acrotomopus atropunctellus* preferred to feed on large- and medium-sized shoots, regardless of sex. However, older plants were resistant to injury. Also, the number of punctures produced by females was 1.6 times greater than that produced by males. Avoiding late planting or harvest dates could be a useful tactic for mitigating *A. atropunctellus* incidence in sugarcane crops.

Key Words: weevil; late harvest date; integrated pest management; puncture

Sumario

El picudo de la caña de azúcar, *Acrotomopus atropunctellus* (Boheman) (Coleoptera: Curculionidae) es una plaga de la caña de azúcar cada vez más importante en Argentina. El periodo de emergencia de los adultos de *A. atropunctellus* coincide con estados fenológicos tempranos de la caña de azúcar (macollaje y gran crecimiento). El objeti-

vo de este trabajo fue conocer la preferencia de alimentación de *A. atropunctellus* según distintos tamaños de brotes de caña de azúcar y el efecto del daño sobre la sobrevivencia de los brotes. *Acrotomopus atropunctellus* prefirió alimentarse de brotes medianos y grandes y la preferencia no varió según el sexo. Sin embargo, las plantas más viejas fueron resistentes al daño. Además, el número de perforaciones producidas por las hembras fue 1.6 veces mayor que las producidas por los machos. Evitar fechas de siembra y cosechas tardías puede contribuir a la reducción de incidencia de *A. atropunctellus* en los cultivos de caña de azúcar.

Palabras Clave: picudo; fecha tardía de cosecha; manejo integrado de plagas; perforaciones

References Cited

- Behmer ST, Joern A. 1994. The influence of proline on diet selection: sex-specific feeding preferences by the grasshoppers *Ageneotettix deorum* and *Phoetaliotes nebrascensis* (Orthoptera: Acrididae). *Oecologia* 98: 76–82.
- Beuzelin JM, Mészáros A, Akbar W, Reagan TE. 2011. Sugarcane planting date impact on fall and spring sugarcane borer (Lepidoptera: Crambidae) infestations. *Florida Entomologist* 94: 242–252.
- Box HE. 1929. Observaciones sobre taladradores de la caña de azúcar. Una plaga nueva de la caña de azúcar: El gorgojo taladrador (Coleoptera: Curculionidae). *Revista Industrial Agrícola de Tucumán* 19: 319–322.
- Di Rienzo AJ, Macchiavelli R, Casanoves F. 2009. Modelos Mixtos en Infostat. Tutorial, Grupo InFoStat, Facultad de Ciencias Agropecuarias, Universidad Nacional de Córdoba, Córdoba, Argentina.
- Digonzelli PA, Romero ER, Scandaliaris J, Giardina J, Arce O. 2005. Efecto de la época de plantación en la dinámica de la emergencia de caña semilla de alta calidad (termotratada y micropropagada) de las variedades CP 65-357 y LCP 85-384. *Revista Industrial Agrícola de Tucumán* 82: 45–53.
- Digonzelli PA, Romero ER, Scandaliaris J, Arce O, Fernández de Ullivarri J, Tonatto J, Leggio Neme MF. 2006. Dinámica de la brotación potencial de caña semilla micropropagada y termotratada de tres cultivares de caña de azúcar. *Revista Industrial Agrícola de Tucumán* 83: 1–7.
- Ghosh MR. 1989. Concepts of Insect Control. New Age International, Limited, New Delhi, India.
- Ostengo S, Espinosa MA, Díaz JV, Chavanne ER, Costilla DD, Cuenya MI. 2014. Mejoramiento genético. Distribución de variedades comerciales de caña de azúcar en la provincia de Tucumán, R. Argentina. Relevamiento de la campaña 2013/2014. *Avance Agroindustrial* 35: 10–14.
- Pérez MLP, Del Río MG, Lanteri, AA. 2012a. Posición taxonómica de *Acrotomopus atropunctellus* (Coleoptera: Curculionidae) y descripción del daño producido en el cultivo de caña de azúcar en la Argentina. *Revista de la Sociedad Entomológica Argentina* 71: 265–273.
- Pérez MLP, Isas MG, Salvatore AR, Gastaminza GA, Trumper EV. 2012b. Periodo de emergencia y proporción de sexos de *Acrotomopus atropunctellus* (Boheman) (Coleoptera: Curculionidae) en el cultivo de la caña de azúcar, campaña 2011–2012. Resúmenes Zoología Agrícola. Jornadas Fitosanitarias Argentinas 14. 3, 4 y 5 de Octubre de 2012. Potrero de los Funes, San Luis, Argentina.
- Pérez MLP, Isas M, Salvatore AR, Trumper E, Pérez DO, Gastaminza G. 2016. Picudo perforador de la caña de azúcar. *Avance Agroindustrial* 37: 30–31.
- Pinheiro JC, Bates DM. 2000. Mixed-effects Models in S and S-PLUS. Statistics and Computing, Springer, New York, New York.
- Pinheiro J, Bates D, DebRoy S, Sarkar D. 2015. Linear and Nonlinear Mixed Effects Models. R package version 3.1-57.
- Romero, E. 2002. Dinámica de la brotación, emergencia y crecimiento inicial de la caña de azúcar. Efecto del genotipo, factores ambientales y manejo. Doctoral thesis. Facultad de Agronomía y Zootecnia, Universidad Nacional de Tucumán, Argentina.
- Romero ER, Scandaliaris J, Olea I, Tonatto J, Sotomayor L. 2005. Emergence and early growth of plant and ratoon cane crops under different temperature regimes, pp. 168–175 *In Proceedings of the International Society of Sugar Cane Technologists Congress* 25, Guatemala, Guatemala.
- Salvatore AR, Acosta M, Willink E. 2006. El picudo perforador de la caña de azúcar *Acrotomopus atropunctellus* (Coleoptera: Curculionidae) en caña de azúcar Tucumán, Argentina. Resumen presentado en XII Jornadas Fitosanitarias Argentina, Catamarca, Junio de 2006.