



Interaction and Distribution of Beetles (Insecta: Coleoptera) Associated with *Heliconia bihai* (Heliconiaceae) Inflorescences

Authors: Oliveira, Thais Ranielle Souza de, Moura, Shayne Rodrigues de, Cruz, Denise Dias da, Loges, Vivian, and Martins, Celso Feitosa

Source: Florida Entomologist, 101(2) : 160-165

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.101.0202>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Interaction and distribution of beetles (Insecta: Coleoptera) associated with *Heliconia bihai* (Heliconiaceae) inflorescences

Thais Ranielle Souza de Oliveira^{1,*}, Shayne Rodrigues de Moura², Denise Dias da Cruz³, Vivian Loges², and Celso Feitosa Martins³

Abstract

The insect fauna associated with the genus *Heliconia* (Heliconiaceae) is quite diversified and includes terrestrial and aquatic species. In plants with an upright inflorescence, the bracts may form phytotelmata. Insects are one of the main taxa that have adapted to phytotelmata environments, including species of Diptera, Coleoptera, Hemiptera, Dermaptera, Plecoptera, Trichoptera, and Odonata, among other orders. The Coleoptera fauna associated with phytotelmata is poorly studied, and the objective of this study was therefore to determine the distribution of beetles in *Heliconia bihai* (L.) (Zingiberales: Heliconiaceae) bracts in cultivated and uncultivated areas in northeastern Brazil. In addition, we describe their functional relationships. With respect to the coleopteran assemblage, 6 families were found in both areas and 3 were found only in the uncultivated *Heliconia*. The most abundant and frequently occurring family in both areas was Hydrophilidae, represented by a single species, *Pelosoma lafertei* (Mulsant) (Coleoptera: Hydrophilidae). Staphylinidae were the second most abundant and the second most frequently occurring family. The detritivores were most abundant (6 species), followed by herbivores (3 species), and predators (2 species). Abundance and richness were higher in the uncultivated area. Temperature and pH were the main factors affecting the coleopteran assemblages. A high acidity indicates an environment rich in organic matter, the main food source in phytotelmata environments.

Key Words: phytotelmata; aquatic insects; tropical flowers; bracts

Resumo

A fauna de insetos associada ao gênero *Heliconia* (Heliconiaceae) é bastante diversificada e inclui espécies terrestres e aquáticas. Em plantas com inflorescência vertical, as brácteas podem formar fitotelmas. Os insetos são um dos principais táxons adaptados aos ambientes fitotelmatas, incluindo espécies de Diptera, Coleoptera, Hemiptera, Dermaptera, Plecoptera, Trichoptera e Odonata, entre outros ordens. A fauna de Coleoptera associada ao ambiente fitotelmata é pouco estudada, e o objetivo deste estudo foi determinar a distribuição de besouros nas brácteas de *H. bihai* (L.) (Zingiberales: Heliconiaceae) em áreas cultivadas e não cultivadas no Nordeste do Brasil. Além disso, descrevemos suas relações funcionais. Em relação à ocorrência de coleópteros, 6 famílias foram encontradas em ambas as áreas e 3 foram encontradas apenas em brácteas de *Heliconia* não cultivada. A família mais abundante e frequente em ambas as áreas foi Hydrophilidae, representada por uma única espécie, *Pelosoma lafertei* (Mulsant) (Coleoptera: Hydrophilidae). Os Staphylinidae foram a segunda família mais abundante e mais frequente. Os detritívoros foram mais abundantes (6 espécies), seguidos de herbívoros (3 espécies), e predadores (2 espécies). A Abundância e a riqueza foram maiores na área não cultivada. A temperatura e o pH foram os principais fatores que afetaram a assembléia de coleópteros. Uma alta acidez indica um ambiente rico em matéria orgânica, o principal recurso alimentar do ambiente fitotelmata.

Palavras Chave: fitotelmata; insetos aquáticos; flores tropicais; brácteas

In plants with upright inflorescences, microhabitats called phytotelmata may form in the bracts. Each phytotelma consists of a small water reservoir formed by plant structures that have the ability to retain water (Maguire 1971; Machado-Alisson et al. 1983; Richardson et al. 2000). There is a great diversity of species associated with these small ecosystems. They occur primarily in plants found in the Neotropical region, where more than 1,500 plant species possess phytotelma, for example in the carnivorous plants, bromeliads, and axillary buds of many species that accumulate water. Phytotelma may contain a rich

fauna, mainly arthropods, forming simple to complex food webs (Fish 1983; Kitching 2001). In phytotelma environments, the combination of physical and chemical conditions of the water, such as volume, pH, temperature, electrical conductivity, and the amount of available resources (e.g., organic matter and debris), regulates the abundance of the species present (Kitching 2001).

The insect fauna associated with the genus *Heliconia* (Heliconiaceae) is quite diversified and includes both terrestrial and aquatic species. Insects are one of the principal groups that have adapted to

¹University Center Euro American, Department of Health Sciences, Brasília, Distrito Federal 70200-001, Brazil, E-mail: thaisranielle@gmail.com (T. R. S. O.)

²Universidade Federal Rural de Pernambuco, Departamento de Agronomia, Recife, Pernambuco 52171-900, Brazil, E-mail: shaynermoura@gmail.com (S. R. M.); vloges@yahoo.com (V. L.)

³Federal University of Paraíba, Department of Systematics and Ecology, João Pessoa, Paraíba 58051-900, Brazil, E-mail: denidacruz@dse.ufpb.br (D. D. C.); cmartins@dse.ufpb.br (C. F. M.)

*Corresponding author; E-mail: thaisranielle@gmail.com

phytotelma environments, including, among other orders, species of Diptera, Dermaptera, Coleoptera, Hemiptera, Hymenoptera, and Lepidoptera (Seifert & Seifert 1979; Seifert 1980; Frank & Barrera 2010; Oliveira et al. 2010; Jalinsky et al. 2014; Staines & Garcia-Robledo 2014). The order Diptera occurs widely in all types of phytotelmata; however, studies on other groups of insects in these environments are more limited (Campos 2010). In particular, information on coleopteran fauna associated with *Heliconia* phytotelmata is still limited. Most of the species inhabiting phytotelmata are terrestrial; however, a few families are semiaquatic (e.g., Chrysomelidae) or aquatic (e.g., Hydrophilidae) (Frank & Barrera 2010; Aristizábal et al. 2013).

In Brazil, publications addressing the phytotelmata fauna focus on bromeliad water reservoirs (called 'tanks') in the northern and southeastern regions of the country (Mestre et al. 2001; Torreias & Ferreira-Keppler 2011). Studies associating the Coleoptera order with *Heliconia* are scarce. However, the presence of beetles of the families Hydrophilidae, Staphylinidae, Nitidulidae, and Curculionidae in *Heliconia bihai* (L.) (Zingiberales: Heliconiaceae) was reported in northeastern Brazil by Oliveira et al. (2010).

One approach to understand the role of biodiversity is functional diversity, in which the species are classified based on their effects on ecosystem processes. There are different ways of quantifying functional diversity, one of which is to form groups of species that have similar functions in the ecosystem (Díaz & Cabido 2001). This allows us to discern the trophic relationships between these organisms and to better understand the ecosystem structure.

According to Benítez-Malvido et al. (2016), tropical forest fragmentation affects biotic interactions; however, information on how this process affects animal trophic guilds and their pattern of interactions with host plants is scarce. Those authors sought to assess changes in biotic interactions in continuous forests and fragments, assigning arthropods associated with *Heliconia aurantiaca* Verschaff. (Zingiberales: Heliconiaceae) into broad trophic guilds such as omnivores, herbivores, and predators. In this paper, we investigated coleopterans associated with *Heliconia*, with the objective to determine the distribution of beetles in *H. bihai* bracts, to categorize their functional relationships, and to determine if there is a specific relationship between the species and the phytotelma environment (wild and cultivated plants). We hypothesized that species richness, abundance, and diversity of Coleoptera in inflorescences of cultivated species would be lower than in inflorescences of naturally occurring species.

Materials and Methods

Inflorescences of *H. bihai* were harvested on a farm containing areas with cultivated and wild (uncultivated) *Heliconia*, located in Camaragibe-PE (8.0219°S, 34.9925°W), in northeastern Brazil. The vegetation in this area is characterized by trees, grass, and sugar cane. The distance between the cultivated area and the wild area is 700 m.

The cultivated area represented the collection of *Heliconia* germplasm of the Federal Rural University of Pernambuco, planted in Jan 2007, with partial shade provided by trees. Plants were cultivated with spacing of 3 m, and 4 m between rows, prepared in blocks with 4 replicates per genotype. Each plot was formed by a clump and covered a total area of 12 m², representing 6 m² of useful area. Irrigation was provided through a high sprinkler, and insecticides were not applied in the area.

The wild (uncultivated) area was represented by an Atlantic Forest fragment with a total area of 145.3 m², classified as Lowland Ombrophylous Forest (Veloso et al. 1992). It is an evergreen forest with a canopy height of 30 m, with emergent trees up to 40 m tall. The

area features clumps of *H. bihai* that are located in a steep portion between the edge of the forest and the watercourse, forming a shaded and moist area.

Inflorescences were collected monthly from Aug 2011 to Aug 2012; field collections were initiated at 7:00 AM and continued until 9:00 AM. In each area, all available inflorescences were collected and the bracts were counted. Abiotic parameters were measured directly on all bracts. The pH of the water in the bract was measured using a Hanna Instruments® portable potentiometer (Model number HI98100, Woonsocket, Rhode Island, USA).

To determine water temperature, an Incoterm® portable digital thermometer (Model number 6132, Porto Alegre, Rio Grande do Sul, Brazil), was used. Water volume in the tank was measured with a 5-mL graduated pipette. The liquid contained within the bracts was then pipetted into individual plastic pots. To collect the insects, the inflorescences were dissected and a brush and water jets were used to dislodge adults and larvae from inside the floral bracts. Subsequently, this liquid was examined with a stereoscopic microscope to collect the remaining insects. The insects were stored in bottles in 70% alcohol for further sorting and counting.

For each species, abundance, relative abundance, and frequency of occurrence were calculated. The abundance of each species was calculated by totaling the total number of individuals observed in that species. The relative abundance was the number of a species in relation to the total number of individuals of all species observed in each inflorescence. The frequency of occurrence corresponds to the proportion of inflorescences in which each species occurred relative to the total number of inflorescences sampled. Diversity and evenness of the beetle fauna were analyzed using the Shannon-Wiener (H') and Pielou (J') indices (Magurran 2004). The species were classified following Bouchard et al. (2011) and vouchers were deposited in the Entomological Collection of the Zoology Department of the Federal University of Paraná, Curitiba, Brazil. The functional categorization of the beetle fauna followed the classification of Kitching (2000) and Merritt & Cummins (1996).

The Shapiro Wilk Test was used to test the normality of the data. Levene's test was used to test the equality of the variances (homoscedasticity) of the abiotic and biotic parameters. The Mann-Whitney U Test was used to compare the diversity, richness, and abundance between the 2 collection areas. Kruskal-Wallis analysis was used to compare abundance by species, richness, and abundance among inflorescences with 2, 3, 4, and 5 bracts. All analyses were carried out using the software package Statistica 8.0 (Statsoft 2010).

To test the effect of environmental variables on the distribution patterns of the coleopteran phytotelmata fauna, we used canonical correspondence analysis. The canonical correspondence analysis was performed to verify the degree of relationship among abiotic variables (pH, temperature, and volume) and the most abundant species in each area (uncultivated and cultivated). The Monte Carlo test was performed with a standard of 999 permutations and a significance level of 5%. The tests were conducted in the software Canoco 4.5 (Ter Braak & Šmilauer 2002).

Results

A total of 2,331 beetles was collected from 293 *H. bihai* inflorescences, representing 9 families of Coleoptera (Table 1). Six families were found in both areas and 3 were found only in the uncultivated *Heliconia* (Table 1). Six families occurred in both the larva and adult stages, but most individuals were adults. The families with aquatic habits were Hydrophilidae and *Elmidae* (both Coleoptera), whereas the family with semiaquatic habits was Nitidulidae and those with terres-

Table 1. Relative abundance (RA), frequency of occurrence (F), presence of adult (A) and larval (L) stages, and functional category of Coleoptera (FC) collected from uncultivated and cultivated *Heliconia bihai* inflorescences in Camaragibe-PE. Y indicates presence of the stage; N indicates absence.

Family	Morphospecies	Uncultivated		Cultivated		FC
		RA (%)	F (%)	RA (%)	F (%)	
Hydrophilidae	<i>Pelosoma lafertei</i>	71.1	89.6	59.2	52.0	Detritivore
Staphylinidae	<i>Paederomimus</i> sp.	8.1	37.0	15.9	17.1	Predator
	<i>Tachyporus</i> sp.	12.3	39.0	9.3	13.6	Predator
Nitidulidae	<i>Colopterus vulneratus</i>	1.1	1.9	3.2	3.0	Detritivore
	Nitidulidae sp.	4.5	24.7	11.8	16.1	Detritivore
Histeridae	<i>Hololepta</i> sp.	0.3	1.9	0.1	0.5	Detritivore
Curculionidae	<i>Metamasius hemipterus</i>	0.3	2.6	0	0	Herbivore
	Curculionidae sp.	0.1	0.6	0.3	1.0	Herbivore
Elmidae	Elmidae sp.	1.3	8.4	0	0	Detritivore
Scarabaeidae	Scarabaeidae sp.	0.3	1.9	0	0	Detritivore
Not identified	Not identified sp.	0.1	0.6	0	0	—
Chrysomelidae	Chrysomelidae	0.5	3.9	0.2	0.5	Herbivore

trial habits were Chrysomelidae, Staphylinidae, Histeridae, Curculionidae, and Scarabaeidae (all Coleoptera). The number of families with terrestrial habits was higher than the number of those with aquatic or semiaquatic habits.

The most abundant and frequently occurring family in the 2 areas was Hydrophilidae, represented by a single species, *Pelosoma lafertei* (Mulsant) (Coleoptera: Hydrophilidae). Staphylinidae was the second most abundant and frequently occurring family, represented by the only 2 predator species observed in the bracts (Table 1). The majority of the species among the sampled Coleoptera were detritivores (6), followed by herbivores (3), and predators (2) (Table 1).

The total number of beetles collected from the uncultivated area was 1,574, whereas we collected 757 from the cultivated area. Twelve morphospecies were recorded in the uncultivated area and 8 in the cultivated area (Table 1). Nitidulidae, Staphylinidae, and Curculionidae were the most species-rich taxa, with 2 morphospecies each. The uncultivated area displayed greater species richness (uncultivated: 1.57 ± 0.9 ; cultivated: 0.79 ± 1.4) and abundance (uncultivated: 6.9 ± 9.0 ; cultivated: 2.4 ± 5.0) per inflorescence (richness: $U = 7,059$; $Z = -4.7$; $P < 0.001$; abundance: $U = 6,701$; $Z = -4.5$; $P < 0.001$).

Table 2. Mean occurrence of key species and richness and abundance of insects in the uncultivated area when inflorescence size (number of bracts) is compared using the Mann-Whitney test.

	Uncultivated				
	Number of Bracts		Mann-Whitney		
	2	4	U	Z	P
<i>Pelosoma lafertei</i>	1.1 ± 3.1	6.6 ± 7.8	196	-2.9	0.002
<i>Paederomimus</i> sp.	0.1 ± 0.4	0.5 ± 0.7	242	-2.5	0.009
Richness	0.6 ± 0.6	1.8 ± 3.3	160	-3.5	<0.0001
Abundance	1.8 ± 3.1	9.1 ± 7.1	139	-3.8	<0.0001
	2	5	U	Z	P
<i>Pelosoma lafertei</i>	1.1 ± 3.1	6.3 ± 10.1	137	-3.2	0.001
<i>Paederomimus</i> sp.	0.1 ± 0.4	1.1 ± 1.6	174	-2.9	0.003
Richness	0.6 ± 0.6	2.1 ± 1.6	145	-3.0	0.002
Abundance	1.8 ± 3.1	9.8 ± 12.4	155	-2.7	0.005
Overall abundance	3	4	U	Z	P
	5.1 ± 6.4	9.1 ± 7.1	382	-2.2	0.025

There was no difference in evenness between areas (uncultivated: 0.71; cultivated: 0.74).

We sampled approximately equal numbers of inflorescences (20–50), with 2 to 5 bracts per inflorescence, in each area. In the uncultivated area, when comparing the inflorescences composed of a smaller number of bracts (2) with those with larger numbers (4 and 5), the abundance of *P. lafertei* and *Paederomimus* sp. (Coleoptera), and the abundance and total richness of beetles were higher in inflorescences with more bracts (Table 2). Comparing inflorescences with 3 and 4 bracts, only the total abundance of beetles was significantly different (Table 2).

A similar pattern can be observed in the cultivated area, where the abundance of *P. lafertei* and *Paederomimus* sp., and the abundance and total richness of beetles also were higher in inflorescences with more bracts (4 and 5) when compared to those with fewer bracts (2 and 3) (Table 3). The abundance of *Tachyporus* sp. (Coleoptera) in inflorescences with 3 and 4 bracts also was higher in the inflorescences with more bracts (Table 3).

The abiotic factors showed a small range in all size inflorescences (2–5 bracts) (Table 4) in both cultivated and uncultivated areas. Inflorescences in cultivated area showed bracts with a more acidic environment, with higher temperatures and less volume (Table 4).

Canonical correspondence analyses suggest that abiotic variables may affect the species distribution in the inflorescence, with significant correlations in both cultivated (Trace = 0.248; F-ratio = 2.24; $P = 0.046$) and uncultivated (Trace = 0.158; F-ratio = 1.77; $P = 0.041$) areas. In the cultivated area, *Pelosoma lafertei* was positively correlated with pH and volume, whereas *Tachyporus* sp. and *Paederomimus* sp. were positively correlated with temperature. In the uncultivated area, *P. lafertei* and *Elmidae* sp. were positively correlated with volume in the bracts, and *Tachyporus* sp. and *Paederomimus* sp. were affected by the temperature (Fig. 1).

Discussion

The uncultivated area presented a greater diversity relative to the cultivated area. Inflorescences with higher numbers of bracts showed

Table 3. Mean occurrence of key species and richness and abundance of insects in the cultivated area when inflorescence size (number of bracts) is compared using the Mann-Whitney test.

	Cultivated				
	Average \pm SD		Mann-Whitney		
	2	5	U	Z	P
<i>Pelosoma lafertei</i>	0.3 \pm 0.5	3.5 \pm 6.4	439	-2.8	0.004
<i>Paederomimus</i> sp.	0.03 \pm 0.1	0.9 \pm 1.9	465.5	-3.2	0.001
Richness	0.6 \pm 0.7	1.2 \pm 1.0	472	-2.3	0.017
Abundance	0.8 \pm 1.2	5.0 \pm 8.3	313	-4.0	0.000
	3	4	U	Z	P
<i>Tachyporus</i> sp.	0.03 \pm 0.1	0.3 \pm 0.6	1098	-2.6	0.007
	3	5	U	Z	P
<i>Pelosoma lafertei</i>	0.7 \pm 2.4	3.5 \pm 6.4	843	-3.1	0.001
<i>Paederomimus</i> sp.	0.2 \pm 0.1	0.9 \pm 1.9	652	-3.8	<0.0001
Richness	0.5 \pm 0.7	1.2 \pm 1.0	598.5	-4.1	<0.0001
Abundance	1.4 \pm 2.8	5.2 \pm 8.3	598.5	-4.1	0.000
	4	5	U	Z	P
<i>Pelosoma lafertei</i>	1.1 \pm 3.2	3.5 \pm 6.4	707.5	-2.9	0.003
Abundance	2.2 \pm 3.8	5.2 \pm 8.3	666	-2.9	0.002

greater abundance and total richness of beetles. The abundance of *Paederomimus* sp. and *P. lafertei* also presented this pattern in both areas. As the number of bracts increased in the inflorescences, abundance and richness also tended to increase. This change in the colonizing fauna has been reported previously (Seifert 1982; Richardson & Hull 2000; Garcia-Robledo et al. 2005; Yee & Willig 2007) and recently, Benítez-Malvido et al. (2016) have shown the influence of the fragmentation of the tropical forest on the phytotelmata arthropod biodiversity in *Heliconia* bracts. The habitat affects species abundance, and the diversity and composition were higher in continuous forest (Benítez-Malvido et al. 2016).

The Coleoptera fauna was predominantly detritivorous. The species *P. lafertei*, similar to a large portion of the Hydrophilidae, feed on organic matter present in the environment; adult beetles are mainly saprobiotic, whereas the larvae are predators of several invertebrates (Hansen 1999; Short & Hebauer 2006; Fikáček et al. 2010). *Pelosoma lafertei* has been reported previously in Pernambuco associated with *H. bihai* (Oliveira et al. 2010). *Elmidae* presented a low level of abundance; however, this family is common in aquatic environments in South America (Manzo 2005). *Elmidae* are detritivores, and were associated with the phytotelmata of bamboo and of floral bracts of palm trees in Venezuela (Sanchez & Liria 2009). In Brazil, they were associated with the phytotelmata of bromeliads (Bromeliaceae) in the Amazon region. Larvae may have 5 to 8 instars, depending on the genus. They can be found on stones or between mosses and on the roots of plants

(Passos et al. 2007; Elliott 2008). There are many ecologically similar species that seemingly occupy the same niche.

Nitidulidae are usually found in organic substrates such as decaying fruits, animal carcasses, flowers, and fungi. Some species have predatory habits (Fernandes et al. 2012). In South America, nitidulids were reported in the bromeliad *Aechmea distichantha* Lem. (Bromeliaceae) in Argentina (Montero et al. 2010), and in inflorescences of *Xanthosoma* Schott (Araceae) in Costa Rica (Garcia-Robledo et al. 2005). In southeastern Brazil, nitidulids are associated with the bromeliad *Vriesea inflata* Wawra (Bromeliaceae) in fragments of the Atlantic Forest (Mestre et al. 2001).

The fact that the number of beetles with terrestrial habits was higher than those with aquatic habits may be associated with the fact that these species are visitors. They usually are found foraging on different parts of the inflorescence. There is a high diversity of terrestrial species in phytotelma environments in the neotropical region (Campos & Fernández 2011), and many studies have reported terrestrial beetles in bromeliads (Campos 2010; Torreias & Ferreira-Keppler 2011). Some studies have reported terrestrial families in *Heliconia* bracts (Frank & Barrera 2010; Oliveira et al. 2010; Aristizábal et al. 2013). The family Scarabaeidae has a diverse biology, including adults that feed on excrements, carcasses, fungi, foliage, pollen, fruits, or roots; some species are diurnal and can be found on leaves (Silva et al. 2011). Many species are pollinators, recycling plant material and excreta, and are considered beneficial (Schiestl & Dotterl 2012). There are few records associating this family with phytotelmata, but in *Heliconia* spp., they are more common in parts of the plant where water does not accumulate (Aristizábal et al. 2013). Species associated with the rhizome and the pseudostem of *Heliconia* inflorescences were reported in Colombia (Aristizábal et al. 2013).

Two predators of the family Staphylinidae, *Tachyporus* sp. and *Paederomimus* sp., were identified. In *H. bihai*, species of this family of beetles were recorded as top predators in the food chain formed inside the bracts, feeding on Diptera larvae (Frank & Barrera 2010). This family is particularly interesting from the functional perspective because they can be detritivorous, herbivorous, mycophagous, or predacious (Clough et al. 2007). These insects occupy different habitats, such as fallen parts of plants, tree holes, debris, vegetation, flowers, etc. (Honek & Kocian 2003; Greeney 2001; Sanabria et al. 2008; Brunke & Marshall 2011).

Histeridae also occupy different types of habitats such as manure, carrion, and decomposing plant material (Hawkeswood 2006). The adults also have been found in other Eryngium (Apiaceae) phytotelmata in the subtropical and temperate regions of Argentina (Campos & Fernández 2011). However, these species represent a small part of the species complex.

Among the herbivores, the family Curculionidae has the largest number of records of insects found in phytotelma environments (Campos & Fernández 2011); it was found in inflorescences of *Heliconia* spp. cultivated in Colombia (Aristizábal et al. 2013), as well as in bromeliad species in Argentina (Montero et al. 2010) and in Brazil (Mestre et

Table 4. Values (average \pm SD) of pH, volume (mL), and temperature ($^{\circ}$ C) in the inflorescence groups with 2, 3, 4, and 5 *Heliconia bihai* bracts from cultivated and uncultivated areas in Camaragibe-PE.

No. of bracts/ inflorescence	Cultivated			Uncultivated		
	pH	Temperature ($^{\circ}$ C)	Volume (mL)	pH	Temperature ($^{\circ}$ C)	Volume (mL)
2	6.5 \pm 1.3	27.3 \pm 2.9	2.0 \pm 1.6	6.5 \pm 1.1	25.9 \pm 1.5	5.8 \pm 1.1
3	6.8 \pm 1.4	28.4 \pm 2.3	1.5 \pm 0.9	6.6 \pm 1.0	26.3 \pm 1.8	4.7 \pm 0.6
4	7.1 \pm 1.3	28.9 \pm 2.4	1.5 \pm 1.1	5.8 \pm 1.4	25.2 \pm 2.8	4.7 \pm 0.5
5	7.2 \pm 1.7	28.7 \pm 2.1	1.6 \pm 1.2	6.6 \pm 0.8	27.0 \pm 1.6	4.5 \pm 0.7




Fig. 1. Ordination diagram produced by canonical correspondence analysis for the samples of *Heliconia bihai* inflorescences collected in the (A) cultivated and (B) uncultivated area. The species are represented by triangles with the abbreviations of their scientific names, and the environmental variables are represented by vectors. Pel = *Pelosoma lafertei*; Tac = *Tachyporus* sp.; Pae = *Paederomimus* sp.; Elm = *Elmidae* sp.

al. 2001). *Metamasius hemipterus* (L.) (Coleoptera: Dryophthoridae) occurred infrequently, and this species has been observed in *H. bihai* bracts in Pernambuco (Oliveira et al. 2010). Another species from this genus, *Metamasius callizona* (Champion) (Coleoptera: Dryophthoridae), is found in native bromeliads in Florida, endangering phytotelmata fauna due to the destruction of these plants (Frank & Fish 2008).

Chrysomelidae had the second highest number of species associated with phytotelma. Adults feed externally on plant tissue, while larvae feed internally and externally. Semiaquatic beetles of this family associated with *Heliconia* bracts are frequently found feeding on the bract, but located below the surface of the floral liquid (Naem 1990). Insects of this family are commonly found in phytotelma of Zingiberales (Poales) species. It seems that there is some phylogenetic influence in the choice of the host plant (Schmitt & Frank 2013).

Pelosoma lafertei is typical of the *Heliconia*-inhabiting species found in this study because it is affected by pH and water volume.

The environmental condition seems to guarantee the occurrence of *P. lafertei*. The acidity indicates that the environment is rich in organic matter, which is the main food source of phytotelmata fauna (Nishadh & Anoopdas 2014) supporting *P. lafertei*, which is a detritivorous species. The higher water volume in the uncultivated area could allow the occurrence of aquatic *Elmidae* species (Manzo 2005). Bracts with lower temperatures guarantee the occurrence of *Paederomimus* sp. in the uncultivated area. *Paederomimus* sp. and *Tachyporus* sp. are predators and consume Diptera larvae, which are extremely abundant in the bracts (Frank & Barrera 2010).

Our studies demonstrated a greater richness of species associated with the uncultivated area. Also, bract features such as greater volume, higher acidity, and lower temperature seem to influence the Coleoptera assemblage in *H. bihai* bracts. This is important because not only does the uncultivated area contribute to the maintenance of diversity, but these plants may also represent a source of species potentially harmful to crops.

Acknowledgments

The authors would like to thank the undergraduate work-study students from the Floriculture Laboratory of the Federal Rural University of Pernambuco (UFRPE), who assisted with data entry and support in the field; and we would like to thank Professor G. H. Rosado Neto (Universidade Federal do Paraná, Brazil) for species identification. Funding for this project was granted by the Coordination of Improvement of Higher Level Personnel (CAPES) and the Graduate Program in Biological Sciences-Zoology of the Federal University of Paraíba, João Pessoa, Paraíba, Brazil (UFPB).

References Cited

- Aristizábal LF, Ospina KA, Vallejo UA, Henao ER, Salgado M, Arthurs SP. 2013. Entomofauna associated with *Heliconia* spp. (Zingiberales: Heliconiaceae) grown in the central area of Colombia. *Florida Entomologist* 96: 112–119. <https://doi.org/10.1653/024.096.0114>
- Benítez-Malvido J, Dáttilo W, Martínez-Falcón AP, Durán-Barrón C, Valenzuela J, López S, Lombera R. 2016. The multiple impacts of tropical forest fragmentation on arthropod biodiversity and on their patterns of interactions with host plants. *PLoS One* 11: 1–15. <http://dx.doi.org/10.1371/journal.pone.0146461>
- Bouchard P, Bousquet Y, Davies AE, Alonso-Zarazaga MA, Lawrence JF, Lyal CH, Newton AF, Reid CAM, Schmitt M, Ślipiński SA, Smith ABT. 2011. Family-group names in Coleoptera (Insecta). *ZooKeys* 88: 1–972. <http://doi.org/10.3897/zookeys.88.807>
- Brunke AJ, Marshall SA. 2011. Contributions to the faunistics and bionomics of Staphylinidae (Coleoptera) in northeastern North America: discoveries made through study of the University of Guelph Insect Collection, Ontario, Canada. *ZooKeys* 75: 29–68. <http://doi.org/10.3897/zookeys.75.767>
- Campos RE. 2010. Eryngium (Apiaceae) phytotelmata and their macro-invertebrate communities, including a review and bibliography. *Hydrobiologia* 652: 311–328. <http://dx.doi.org/10.1007/s10750-010-0364-y>
- Campos RE, Fernández LA. 2011. Coleopterans associated with plants that form phytotelmata in subtropical and temperate Argentina, South America. *Journal of Insect Science* 11: 1–18. <https://doi.org/10.1673/031.011.14701>
- Clough Y, Kruess A, Tschardt T. 2007. Organic versus conventional arable farming systems: functional grouping helps understand staphylinid response. *Agriculture, Ecosystems and Environment* 118: 285–290. <https://doi.org/10.1016/j.agee.2006.05.028>
- Díaz S, Cabido M. 2001. Vive la différence: plant functional diversity matters to ecosystem processes. *Trends in Ecology and Evolution* 16: 646–655. [https://doi.org/10.1016/S0169-5347\(01\)02283-2](https://doi.org/10.1016/S0169-5347(01)02283-2)
- Elliott J. 2008. The ecology of riffle beetles (Coleoptera: Elmidae). *Freshwater Reviews* 1: 189–203.
- Fernandes DRR, Bená DC, Lara RIR, Ide S, Perioto NW. 2012. Nitidulidae (Coleoptera) asociados a frutos de café (*Coffea arabica* L.). *Coffee Science* 7: 135–138.

- Fikáček M, Gentili E, Short AEZ. 2010. Order Coleoptera, family Hydrophilidae. Arthropod Fauna of the UAE 3: 135–165.
- Fish D. 1983. Phytotelmata flora and fauna, pp. 161–190 In Frank JH, Lounibos LP [eds.], Phytotelmata: Terrestrial Plants as Hosts for Aquatic Insect Communities. Plexus, Medford, New Jersey, USA.
- Frank JH, Barrera R. 2010. Natural history of *Belonuchus* Nordmann spp. and allies (Coleoptera: Staphylinidae) in *Heliconia* L. (Zingiberales: Heliconiaceae) flower bracts. Insecta Mundi 110: 1–12.
- Frank JH, Fish D. 2008. Potential biodiversity loss in Florida bromeliad phytotelmata due to *Metamasius callizona* (Coleoptera: Dryophthoridae), an invasive species. Florida Entomologist 91:1–8. [https://doi.org/10.1653/0015-4040\(2008\)091\[0001:PBLIFB\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2008)091[0001:PBLIFB]2.0.CO;2)
- García-Robledo C, Quintero-Marin P, Mora-Kepfer F. 2005. Geographic variation and succession of arthropod communities in inflorescences and infructescences of *Xanthosoma* (Araceae). Biotropica 37: 650–656. <https://doi.org/10.1111/j.1744-7429.2005.00082.x>
- Greeney HF. 2001. The insects of plant-held waters: a review and bibliography. Journal of Tropical Ecology 17: 241–260.
- Hansen M. 1999. World Catalogue of Insects 2. Hydrophiloidea (s.str.)(Coleoptera). Apollo Books Publisher. Devon, United Kingdom.
- Hawkeswood TJ. 2006. Some notes and a review on *Hololepta lissopyga* Marsen, 1853 (Coleoptera : Histeridae). Calodema 6: 1–3.
- Honek A, Kocian M. 2003. Importance of woody and grassy areas as refugia for field Carabidae and Staphylinidae (Coleoptera). Acta Societatis Zoologicae Bohemicae 67: 71–81.
- Jalinsky J, Chaboo CS, Radocy TA, Wertenberger R. 2014. Insect diversity in phytotelmata habitats of two host plants, *Heliconia stricta* Huber (Heliconiaceae) and *Calathea lutea* Schult (Marantaceae) in the south-east Amazon of Peru. Journal of the Kansas Entomological Society 87: 299–311. <https://doi.org/10.2317/JKES130816.1>
- Kitching RL. 2000. Food webs and container habitats: the history and ecology phytotelmata. Journal of Ecology 89: 904–904. <http://dx.doi.org/10.1046/j.0022-0477.2001.00593-4.x>
- Kitching RL. 2001. Food webs in phytotelmata: “bottom-up” and “top-down” explanations for community structure. Annual Review of Entomology 46: 729–760. <https://doi.org/10.1146/annurev.ento.46.1.729>
- Machado-Allison C, Rodríguez D, Barrera R. 1983. The insect community associated with inflorescences of *Heliconia caribaea* Lamarck in Venezuela, pp. 247–270 In Frank JH, Lounibos LP [eds.], Phytotelmata: Terrestrial Plants as Hosts for Aquatic Insect Communities. Plexus Publishing, Medford, New Jersey, USA.
- Maguire B. 1971. Phytotelmata: biota and community structure determination in plant-held waters. Annual Review of Ecology and Systematics 2: 439–464. <https://doi.org/10.1146/annurev.es.02.110171.002255>
- Magurran AE. 2004. Measuring Biological Diversity. Blackwell Publishing, Oxford, United Kingdom.
- Manzo V. 2005. Key to the South America genera of *Elmidae* (Insecta: Coleoptera) with distributional data. Studies on Neotropical Fauna and Environment 40: 201–208.
- Merritt RW, Cummins KW. 1996. An Introduction to the Aquatic Insects of North America. Kendall and Hunt Publishing, Dubuque, Iowa, USA.
- Mestre LAM, Aranha JMR, Esper MLP. 2001. Macroinvertebrate fauna associated to the bromeliad *Vriesea inflata* of the Atlantic Forest (Paraná State, southern Brazil). Brazilian Archives of Biology and Technology 44: 89–94. <http://dx.doi.org/10.1590/S1516-89132001000100012>
- Montero G, Feruglio C, Barberis IM. 2010. The phytotelmata and foliage macrofauna assemblages of a bromeliad species in different habitats and seasons. Insect Conservation and Diversity 3: 92–102. <http://dx.doi.org/10.1111/j.1752-4598.2009.00077.x>
- Naeem S. 1990. Resource heterogeneity and community structure: a case study in *Heliconia imbricata* phytotelmata. Oecologia 84: 29–38. <http://dx.doi.org/10.1007/BF00665591>
- Nishadh KAR, Anoopdas KS. 2014. Tree-hole aquatic habitats: inhabitants, processes and experiments. A review. International Journal of Conservation Science 5: 253–268.
- Oliveira TRS, Sena DCA, Loges, V, Camara CAG, Reis AC. 2010. Insetos (Arthropoda, Insecta) em inflorescências de *Heliconia bihai* (L.) L. (Heliconiaceae). Revista Brasileira Horticultura Ornamental 16: 174–178. <http://dx.doi.org/10.14295/rbho.v16i2.560>
- Passos IM, Nessimian JL, Junior NF. 2007. Chaves para identificação dos gêneros de *Elmidae* (Coleoptera) ocorrentes no Estado do Rio de Janeiro, Brasil. Revista Brasileira de Entomologia 51: 42–53. <http://dx.doi.org/10.1590/S0085-56262007000100008>
- Richardson BA, Hull GA. 2000. Insect colonisation sequences in bracts of *Heliconia caribaea* in Puerto Rico. Ecological Entomology 25: 460–466. <http://dx.doi.org/10.1046/j.1365-2311.2000.00269.x>
- Richardson BA, Rogers C, Richardson MJ. 2000. Nutrients, diversity, and community structure of two phytotelm systems in a lower montane forest, Puerto Rico. Ecological Entomology 25: 348–356. <http://dx.doi.org/10.1046/j.1365-2311.2000.00255.x>
- Sanabria C, Armbrecht I, Gutiérrez-Chacón C. 2008. Diversidad de estafilínidos (Coleoptera: Staphylinidae) en cinco sistemas productivos de los Andes Colombianos. Revista Colombiana de Entomología 34: 217–223.
- Sanchez E, Liria J. 2009. Relative abundance and temporal variation of macroinvertebrates in a Venezuelan cloud forest habitat. International Journal of Tropical Insect Science 29: 3–10. <https://doi.org/10.1017/S1742758409377553>
- Seifert RP. 1980. Mosquito fauna of *Heliconia aurea*. Journal of Animal Ecology 49: 687–697. <https://doi.org/10.2307/422113>
- Seifert RP. 1982. Neotropical *Heliconia* insect communities. The Quarterly Review of Biology 57: 1–128.
- Seifert RP, Seifert FH. 1979. A *Heliconia* insect community in a Venezuelan cloud forest. Ecology 60: 462–467. <http://dx.doi.org/10.2307/1936064>
- Schiestl FP, Dötterl S. 2012. The evolution of floral scent and olfactory preferences in pollinators: coevolution or pre-existing bias? Evolution 66: 2042–2055.
- Schmitt M, Frank M. 2013. Notes on the ecology of rolled-leaf hispinines (Chrysomelidae, Cassidinae) at La Gamba (Costa Rica). ZooKeys 332: 55–69. <https://doi.org/10.3897/zookeys.332.5215>
- Short AE, Hebauer F. 2006. World catalogue of Hydrophiloidae additions and corrections, (1999–2005) (Coleoptera). Koleopterologische Rundschau 76: 315–359.
- Silva, PD, Vaz-de-Mello FZ, Di Mare RA. 2011. Guia de identificação das espécies de Scarabaeinae (Coleoptera: Scarabaeidae) do município de Santa Maria, Rio Grande do Sul, Brasil. Biota Neotropica 11: 329–345.
- Statsoft. 2010. Statistica data analysis software system (version 8.0). <http://www.statsoft.com> (last accessed 12 Jun 2014)
- Staines CL, García-Robledo C. 2014. The genus *Cephaloleia* Chevrolat, 1836 (Coleoptera, Chrysomelidae, Cassidinae). ZooKeys 436: 1–355. <https://doi.org/10.3897/zookeys.436.5766>
- Ter Braak CJ, Šmilauer PC. 2002. Canoco reference manual and CanoDraw for Windows user's guide: software for canonical community ordination (version 4.5). <http://www.canoco.com> (last accessed 12 Jun 2014)
- Torreias S, Ferreira-Keppler RL. 2011. Macroinvertebrates inhabiting the tank leaf terrestrial and epiphyte bromeliads at Reserva Adolpho Ducke, Manaus, Amazonas. Brazilian Archives of Biology and Technology 54: 1193–1202.
- Veloso HP, Rangel Filho ALR, Lima JCA. 1992. “Classificação da vegetação brasileira” adaptada a um sistema universal [online]. In Ministério da Economia, Fazenda e Planejamento, Fundação Instituto Brasileiro de Geografia e Estatística, Diretoria de Geociências, Departamento de Recursos Naturais e Estudos Ambientais [eds.]. Brasília, Brazil. <http://biblioteca.ibge.gov.br/visualizacao/livros/liv63011.pdf> (last accessed 2 Jun 2014)
- Yee DA, Willig MR. 2007. Colonisation of *Heliconia caribaea* by aquatic invertebrates: resource and microsite characteristics. Ecological Entomology 32: 603–612. <http://dx.doi.org/10.1111/j.1365-2311.2007.00918.x>