



First Report of the Eastern Ant Cricket, *Myrmecophilus pergandei* Bruner, (Orthoptera: Myrmecophilidae), Collected from an Imported Fire Ant Colony, *Solenopsis invicta* × *richteri* (Hymenoptera: Formicidae)

Author: Hill, JoVonn G.

Source: Journal of Orthoptera Research, 18(1) : 57-58

Published By: Orthopterists' Society

URL: <https://doi.org/10.1665/034.018.0103>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First report of the eastern ant cricket, *Myrmecophilus pergandei* Bruner, (Orthoptera: Mymecophilidae), collected from an imported fire ant colony, *Solenopsis invicta* × *richteri* (Hymenoptera: Formicidae)

Accepted April 11, 2009

JOVONN G. HILL

Mississippi Entomological Museum, Department of Entomology and Plant Pathology, Box 9775, Mississippi State University 39762.
Email: jgh4@entomology.msstate.edu

Key words

Myrmecophilus pergandei, Mymecophilidae, imported, fire ants, *Solenopsis*, trophallaxis, inquiline

Species in the family Myrmecophilidae (Orthoptera) are small, apterous, inquiline inhabitants of ant nests, and a number of species are known to strigilate and engage in trophallaxis with their ant hosts (Wheeler 1900, Schimmer 1909, Carter 1939, Henderson & Akre 1986). Several studies indicate that the ant hosts are typically aggressive toward the crickets, but that the crickets mimicked the mutual grooming and trophallactic behavior of their hosts thereby getting close to them, and usually avoided attack by using their superb jumping ability (Wheeler 1900, Henderson & Akre 1986).

Four species are known from the United States; however, only the eastern ant cricket (*Myrmecophilus pergandei* Bruner), occurs in the southeastern United States (Blatchley 1920, Hebard, 1920, Capinera *et al.* 2004, MacGown & Hill 2006). *Myrmecophilus pergandei* has been recorded as an inquiline from numerous species of native ants including *Aphaenogaster treatae* Forel, *Camponotus castaneus* (Latreille), *C. chromaiodes* Bolton, *C. herculeanus* (L.), *C. pennsylvanicus* (DeGeer), *C. novaeboracensis* (Fitch), *Crematogaster lineolata* (Say), *Formica fusca* L., *F. obscuriventris* Mayr., *F. pallidefulva* Latreille, *F. subsericea* Say, *F. truncorum* Fab., *Lasius umbratus* (Nylander), and *Paratrechina parvula* (Mayr) (Hebard 1920, Carter 1939). It has also been successfully kept with captive ant colonies, including *Crematogaster lineolata*, *C. missouriensis* Emery, *Neivamyrmex nigrescens* (Cresson) and *Pheidole dentata* Mayr (Carter 1939).

Though *M. pergandei* has been reported to inhabit colonies of *Solenopsis geminata*, a purported exotic ant species in North America, it has not been recorded from colonies of *Solenopsis xyloni* McCook, another purported exotic species, or colonies of the imported fire ants, *Solenopsis invicta* Buren, *S. richteri* Forel, or their hybrid *S. invicta* × *richteri* (Travis 1941). Carter (1939) introduced several *M. pergandei* into a laboratory colony of *S. xyloni* and found that the crickets were met with aggressive behavior and were unable to approach the ants for feeding. In most cases the *Solenopsis* captured and devoured the crickets within 24 h of their introduction into the nest.

However, on 9 September 2008, several individuals of *M. pergandei* were observed within a colony of the hybrid imported fire ant, *Solenopsis invicta* × *richteri*. This colony was not the typical mound in an open area for which imported fire ants are well known, but instead the colony was nesting in a rotting log in an open bottomland hardwood forest in the Noxubee National Wildlife Refuge (lat 33° 16' 17"N, long 88° 47' 17"W). Also found in the colony was

an individual of *Myrmecosaurus ferrugineus* Bruch (Staphylinidae), a beetle that is native to South America and associated with imported fire ant colonies. Voucher specimens of *M. pergandei* and *S. invicta* × *richteri* are both pinned and stored in alcohol. The specimen of *M. ferrugineus* is pinned. All specimens are deposited in the Mississippi Entomological Museum.

This finding represents a new host record for *M. pergandei*, and it is the first report of it occurring with an ant truly known to be exotic. It is interesting that *M. pergandei* has not yet been documented as occurring in colonies of the imported fire ant, given the amount of research conducted on these exotics since their introduction to North America in the early part of the 20th century. However, it is not overly surprising that *M. pergandei* finds imported fire ants a suitable host, as a related species, *Myrmecophila nebrascensis* Bruner, has been reported to occur in the colonies of *Solenopsis xyloni* McCook and *S. invicta* Buren (Neece & Barteli 1981, 1982).

Acknowledgements

This research was supported by the Mississippi Forestry Experiment Station Project MIS-311080, USDA-ARS Areawide Management of Imported Fire Ant Project, R. L. Brown (PI), and a grant from Noxubee National Wildlife Refuge, R. L. Brown, J. A. MacGown, and J. G. Hill (Co PIs). This is publication J-11529 of the Mississippi Agricultural and Forestry Experiment Station.

References

- Blatchley W.S. 1920. Orthoptera of Northeastern America. Nature Publishing Company, Indianapolis.
- Capinera J.L., Scott R.D., Walker T.J. 2004. Field Guide to Grasshoppers, Katydid, and Crickets of the United States. Cornell University Press, Ithaca, NY.
- Carter W.M. 1939. Some Observations of *Myrmecophila* in Mississippi. Unpub. M.Sc. Thesis, Mississippi State University.
- Hebard M. 1920. A revision of the North American species of the genus *Myrmecophila*. Transactions American Entomological Society 46: 91-111.
- Henderson G., Akre R.D. 1986. Biology of the myrmecophilous cricket, *Myrmecophila manni*, (Orthoptera: Gryllidae). Journal Kansas Entomological Society 59: 454-467.
- MacGown J.A., Hill J.G. 2006. The eastern ant cricket, *Myrmecophilus pergandei* Bruner (Orthoptera: Myrmecophilidae), reported from Mississippi, U.S.A. Journal Mississippi Academy of Science 51: 180-182.
- Neece K.C., Barteli D.P. 1981. Insects associated with *Solenopsis* spp. in southeastern Texas. Southwestern Entomologist 6: 307-311.
- Neece K.C., Barteli D.P. 1982. A faunistic survey of the organisms associated with ants of western Texas. Graduate Studies. Texas Tech. University Press. 36 pp.

- Schimmer F. 1909. Beitrag zu einer Monographie der Gryllodeengattung *Myrmecophila* Latr. Zeitschrift Wissenschaftliche Zoologie. 93: 409-534.
- Travis B.V. 1941. Notes on the biology of the fire ant *Solenopsis geminata* (F.) in Florida and Georgia. Florida Entomologist 24: 15-22.
- Wheeler W.M. 1900. The habits of *Myrmecophila nebracensis* Bruner. Psyche 9: 111-115.