

Laboratory Evaluation of Five Chitin Synthesis Inhibitors Against the Colorado Potato Beetle, *Leptinotarsa decemlineata*

Authors: Karimzadeh, R., Hejazi, M. J., Rahimzadeh Khoei, F., and Moghaddam, M.

Source: Journal of Insect Science, 7(50) : 1-6

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.007.5001>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Laboratory evaluation of five chitin synthesis inhibitors against the Colorado potato beetle, *Leptinotarsa decemlineata*

R. Karimzadeh¹, M. J. Hejazi^{1,a}, F. Rahimzadeh Khoei² and M. Moghaddam²

¹ Department of Plant Protection, College of Agriculture, University of Tabriz, Tabriz, 51666- 14888, Iran

² Department of Agronomy and Plant Breeding, College of Agriculture, University of Tabriz, Tabriz, 51666-14776, Iran

Abstract

Results of laboratory experiments are reported that tested the effects of five chitin synthesis inhibitors, diflubenzuron, cyromazine, lufenuron, hexaflumuron and triflumuron, on second instars of the Colorado potato beetle, *Leptinotarsa decemlineata* (Say) (Coleoptera: Crysomelidae), originally collected from potato fields of Bostanabaad, a town 66 km southeast of Tabriz, Iran. In bioassays, the larvae were fed potato leaves dipped in aqueous solutions containing chitin synthesis inhibitors. The mortalities and abnormalities of the treated larvae were recorded 72 hours after treatments. LC₅₀ values were 58.6, 69.6, 27.3, 0.79 and 81.4 mg ai/L for diflubenzuron, cyromazine, lufenuron, hexaflumuron and triflumuron, respectively. Compared with phosalone, which is one of the common insecticides used for controlling this pest in Iran, lufenuron and hexaflumuron seem to be much more potent, and if they perform equally well in the field, they would be suitable candidates to be considered as reduced risk insecticides in management programs for *L. decemlineata* due to much wider margin of safety for mammals and considerably fewer undesirable environmental side effects.

Keywords: cyromazine, benzoylphenyl ureas, diflubenzuron, hexaflumuron, lufenuron, triflumuron

Correspondence: ^amjhejazi@tabrizu.ac.ir

Received: 6 February 2006 | **Accepted:** 23 January 2007 | **Published:** 27 September 2007

Copyright: This is an open access paper. We use the Creative Commons Attribution 2.5 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Volume 7, Number 50

Cite this paper as:

Karimzadeh R, Hejazi MJ, Rahimzadeh Khoei F, Moghaddam M. 2007. Laboratory evaluation of five chitin synthesis inhibitors against the Colorado potato beetle, *Leptinotarsa decemlineata*. 6pp. *Journal of Insect Science* 7:50, available online: insectscience.org/7.50

Introduction

Leptinotarsa decemlineata (Say) (Coleoptera: Chrysomelidae), the Colorado potato beetle is the most devastating defoliator of potato plants worldwide. If populations of this pest are not controlled, they can cause a total loss of yield by defoliating potato plants prior to tuber formation. Tuber formation and filling are the most susceptible stages of potato plants with regard to damage by *L. decemlineata*. Therefore, pest management guidelines are designed to limit total defoliation to 10–25% at these stages (Hare 1990).

Widespread resistance of *L. decemlineata* to most insecticide chemical groups has been reported in many parts of the world (Harris and Svec 1981; French *et al.* 1992). In some parts of the United States of America this insect has even developed resistance to the newly commercialized neonicotinoids (Mota-Sanchez *et al.* 2006). Effective chemical control of this pest requires new insecticides with novel mechanisms of action. The effects of chitin synthesis inhibitors on *L. decemlineata* have been studied by several researchers (Cutler *et al.* 2005a; Furlong and Groden 2001; Sirota and Grafius 1994). Cutler *et al.* 2007 reported that novaluron at 50 g ai/ha provided excellent and prolonged effect against *L. decemlineata* and could be a valuable tool in future *L. decemlineata* management programs. These insecticides with specific mode of action are relatively new to Iran. These compounds interfere with formation of chitin and control immature stages of many pests with relatively low harm to beneficial arthropods (Consoli *et al.* 2001; Wakgari and Giliomee 2003; Catangui *et al.* 1996). Based on the study conducted by Lucas *et al.* 2004 cyromazine was not lethal to the first and third instars of the lady beetle, *Coleomegilla maculata* (a predator of *L. decemlineata*) and seems to be a suitable compound for protection of *C. maculata* populations.

Consoli *et al.* 2001 stated that lufenuron and triflumuron did not affect the parasitization capacity of the parasitoid, *Trichogramma galloi*, and caused almost 100% mortality of larvae of the sugar cane borer, *Diatraea saccharalis*, when used to treat eggs prior to parasitization. Laboratory bioassays indicated that triflumuron did not cause significant mortality in *Coccidoxenoides peregrinus*, a parasitoid of the mealybug, *Pseudococcus longispinus* (Wakgari and Giliomee 2003). Most of these compounds

have low soil persistence and high residual activity on foliage (Cutler *et al.* 2005b; Malinowski and Pawinska 1992; Tuttle and Ferro 1988). Thus, the use of these insecticides may allow build up of the population of at least some natural enemies compared with chemical insecticides of broader spectrum action.

The aim of this study was to assess the effects of some chitin synthesis inhibitors against *L. decemlineata* population from an important potato-growing region of Iran by estimating their LC₅₀ values.

Materials and Methods

Insecticides

The chitin synthesis inhibiting insecticides used were: diflubenzuron (25 WP, Hebei vian biochemical, www.chinanusa.net), hexaflumuron (Consult 10 EC, Dow AgroSciences, www.dowagro.com); triflumuron (Starycide 48 SC, www.bayer.com), lufenuron (Match 50 EC, www.syngenta.com). These compounds belong to benzoylphenyl ureas structurally. The other insecticide tested was cyromazine (Trigard 75 WP, Novartis, www.novartis.com) which belongs to a different chemical group, the triazines.

Insects

The *L. decemlineata* colony was established using adults and eggs collected from potato fields in Bostanabaad and maintained in greenhouse conditions (26 ± 3° C; RH = 50 ± 15%; photoperiod of 16:8 L:D). The 2nd instars of the colony reared over three generations were used to test the insecticides. To have uniformly aged larvae in the experiments, the 1st instars that were ready to molt were separated 24 hours prior to the bioassays and newly (up to 24 hours) molted 2nd instars were used.

Bioassays

Based on preliminary experiments, the ranges of concentrations tested were 62.5–168, 22.5–450, 25–50, 0.5–1.6 and 50–75 mg ai/L for triflumuron, cyromazine, lufenuron, hexaflumuron and diflubenzuron, respectively. Each treatment consisted of five concentrations and a control. Potato leaves were dipped in aqueous solutions with different concentrations of the chitin synthesis inhibitors. Tween 80 was used at a concentration of 500 ppm as a surfactant to ensure complete wetting of the leaves. After drying, the petiole of each leaf was




Figure 1. Two 2nd instars of *Leptinotarsa decemlineata* 72 hours after treatment with Cyromazine. A blister filled with hemolymph protruding from prothorax (A) and the hindgut everted from the anus (B).

inserted in a 1.5 ml micro tube filled with tap water through a hole made in the micro tube cap. Each micro tube was then put in a 14 x 7 x 4 cm transparent plastic box. Fifteen 2nd instars (up to 24 hours old) of *L. decemlineata* were put on each treated leaf. One screened hole (4 cm in diameter), on the removable lid of each plastic box provided ventilation. The boxes were kept in insectarium at $25 \pm 1^\circ \text{C}$, $\text{RH} = 50 \pm 10\%$ and a photoperiod of 16:8 L:D. The mortalities and abnormalities of the treated larvae were recorded 72 hours after treatment. This was done because the larvae in the controls molted to 3rd instar after this period of time. The treatments were replicated four times. Each replication was done at a different day and the solutions used for each treatment were freshly prepared each time.

The probit option of the Statistical Package for Social Sciences was used for data analysis (SPSS 1999). The toxicities of the insecticides tested were evaluated based on 95% confidence limits of LD_{50} ratios. If the 95% confidence interval included 1, then the difference between LC_{50} s were considered insignificant (Robertson and Preisler 1992).

Results and Discussion

Symptoms

Chitin synthesis inhibitors act by interfering with chitin synthesis. Hence, most of the larvae treated with these compounds showed symptoms at molting. Partial molting was seen in some larvae treated with benzoylphenyl ureas. Inability in casting the old head capsule occurred in others. This was followed by death of the larvae.

The toxicity symptoms in larvae treated with cyromazine were most dramatic and seen as elongation and increase in turgor. The intersegmental membranes were stretched. In some treated larvae hindgut protruded from the anus or a blister filled with brown fluid (probably hemolymph) was seen on the first tergum (Figure 1). Hughes *et al.* 1989 also reported these symptoms. These researchers suggest that cyromazine probably decreases chitin synthesis in endocuticle of the hindgut and weakens the junction of the midgut and hindgut. The poisoned larvae became dark red in color and were lethargic, while untreated larvae had a shiny bright red appearance. The blister-like lesions and lethargy shown by the 2nd instars of *L.*




Figure 2. Concentration- response relationship between chitin synthesis inhibitors and the 2nd instars of *Leptinotarsa decemlineata*, 72 hours after treatment.

decemlineata in this study were similar to those reported by Sirota and Grafius (1994). The symptoms exhibited by the treated *L. decemlineata* larvae were consistent with symptoms reported for some other species of insects such as *Lucilia cuprina*, *Manduca sexta* and *Lymantria dispar* treated with chitin synthesis inhibitors (Abdel-Monem *et al.* 1980; Kotze *et al.* 1992; Root and Dauterman 1996). The similarities in symptoms seen in different orders of insects probably indicate a common mode of action (Sirota and Grafius 1994).

LC₅₀ values

LC₅₀ values for the 2nd instars of the *L. decemlineata* are shown in Table 1. Figure 2 presents the relationship between the probit of percentage mortalities and the logarithm of the concentrations of the chitin synthesis inhibitors tested. Hexaflumuron and lufenuron were more effective than the other insecticides. Based on the estimated LC₉₀, the toxicities of the chitin synthesis inhibitors tested can be rated in the following order: hexaflumuron > lufenuron > diflubenzuron > triflumuron > cyromazine.

Triflumuron was the least toxic to the 2nd instars compared with the other benzoylphenyl ureas tested. Malinowski and Pawinska (1992) studied the effect of chlorfluazuron, teflubenzuron, hexaflumuron, triflumuron and GR 572 on *L. decemlineata*. They also reported that triflumuron was less effective than other benzoylphenyl ureas. Based on the 95% confidence intervals of the LC₅₀ ratios, the different insecticides were compared with each other. Toxicities of the CSIs tested were significantly different except that the difference between diflubenzuron and triflumuron was not significant. The LC₅₀ value for triflumuron reported by Cooper *et al.* (1983) was 75 mg ai/L for 2nd instars of *L. decemlineata*, which compares favorably with our estimate of 81.4 mg ai/L. This may be due to the relative similarities in sensitivities of the populations tested and the experimental procedures used. Furlong and Groden (2001) reported 123 mg ai/L as the LC₅₀ of cyromazine for 2nd instars of *L. decemlineata*, which was larger than 69.6 mg ai/L in our study. In their study the larvae fed on treated leaf disks for 24 hours, while in ours the feeding period on treated leaves was 72 hours. The LC₅₀ of

Table 1. LC₅₀ values of CSIs on 2nd instars of *Leptinotarsa decemlineata* 72 hours after using the leaf dip method.

Insecticides	LC ₅₀ (mg ai/L) (95% CI)	Slope ± SE	LC ₉₀ (mg ai/L) (95% CI)	χ ²
Diflubenzuron	58.60 (55.75–61.00)	8.10 ± 1.12	84.50 (78.14–96.17)	1.43 ^{ns}
Cyromazine	69.60 (51.93–89.73)	1.34 ± 0.17	628.00 (403.27–1264.50)	2.80 ^{ns}
Lufenuron	27.30 (23.60–29.94)	4.30 ± 0.74	54.90 (43.40–70.83)	0.96 ^{ns}
Hexaflumuron	0.79 (0.64–0.91)	5.76 ± 0.50	1.30 (1.07–1.96)	7.78 ^{ns}
Triflumuron	81.40 (73.82–88.00)	4.03 ± 0.48	168.50 (148.59–203.63)	1.67 ^{ns}

ns = non-significant

diflubenzuron in our study (58.6 mg ai/L) is comparable to 50 mg ai/L reported by Hegazy *et al.* (1989).

The slopes of the dose-response lines of the chitin synthesis inhibitors tested (except for cyromazine) were quite steep and the differences between the highest and lowest concentrations were low. That is, the population is phenotypically homogeneous, and with a fairly small increase in insecticide concentration, the mortality would increase considerably. This necessitates more careful use of these chitin synthesis inhibitors in the field to prevent exerting a high selection pressure that could eliminate the susceptible individuals and lead to selection of resistant individuals (Robertson and Preisler 1992). Cyromazine is currently proposed for control of Colorado potato beetle and leafminers on potato (Anonymous 1999). This insecticide provided good control of early instars of *L. decemlineata* in field trials as reported by Bishop *et al.* (1990) and Sirota and Grafius (1994).

Our results indicated that among the chitin synthesis inhibitors tested, hexaflumuron and lufenuron (with LC₅₀ values of 0.79 and 27.3 mg ai/L respectively) were the most effective at low concentrations against the *L. decemlineata*. In comparison, phosalone, which is one of the most commonly used insecticides for controlling this pest in Iran, had an LC₅₀ range of 48.72 – 64.12 mg ai/L for five Iranian populations of *L. decemlineata* tested (Pourmirza 2005). Hence, these chitin synthesis inhibitors seem to be more potent and, if they perform equally well in the field, they would be suitable candidates to be considered as reduced risk insecticides for controlling *L. decemlineata* in Iran due to their relatively much wider margin of safety (Catangui *et al.* 1996; Griffith *et al.* 2000; Tunaz and Uygun 2004). Since these compounds do not belong to the groups of insecticides conventionally used for *L. decemlineata* control in Iran, they can be used in rotation with other insecticides. This would conform to the effective *L. decemlineata* management strategy of not applying the same class of compound more than once within a growing season.

Acknowledgements

We thank the Board of Directors of Graduate Studies of the University of Tabriz for financial support of this research. We are also thankful to Syngenta-Iran for kindly providing cyromazine and lufenuron used in this study.

References

- Abdel-Monem AH, Cameron EA, Mumma RO. 1980. Toxicological studies on the molt inhibiting insecticide (EL- 494) against the gypsy moth and effect on chitin biosynthesis. *Journal of Economic Entomology* 73: 22-25.
- Anonymous . 1999. Cyromazine; pesticide tolerance. *Federal register* 64: 50043-50050.
- Bishop B, Grafius E, Henry P, Roragen K, Maier R, Rattlingourd P. 1990. Colorado potato beetle control, foliar insecticides, 1988. *Insecticide & Acaricide Tests* 15: 127-128.
- Catangui MA, Fuller BW, Walz AW. 1996. Impact of Dimilin® on nontarget arthropods and its efficacy against rangeland grasshoppers. In: U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 1996. *Grasshopper Integrated Pest Management User Handbook*. Technical Bulletin No. 1809. Section VII. Washington, DC.
- Consoli FL, Botelho PSM, Parra JRP. 2001. Selectivity of insecticides to the egg parasitoid *Trichogramma galloi* Zucchi 1988, (Hym., Trichogrammatidae). *Journal of Applied Entomology* 125: 37- 43.
- Cooper RM, Lindquist RK, Simonet DE. 1983. Timing applications of SIR 8514 for control of the Colorado potato beetle (Coleoptera:Chrysomelidae) on potatoes. *Journal of Economic Entomology* 76: 563-566.
- Cutler GC, Scott-Dupree CD, Tolman JH, Harris CR. 2007. Field efficacy of novaluron for control of Colorado potato beetle (Coleoptera:Chrysomelidae) on potato. *Crop Protection* 26:760-767.
- Cutler GC, Scott-Dupree CD, Tolman JH, Harris CR. 2005a. Acute and sublethal toxicity of novaluron, a novel chitin synthesis inhibitor, to *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae). *Pest Management Science* 61: 1060-1068.
- Cutler GC, Tolman JH, Scot-Dupree CD, Harris CR. 2005b. Resistance potential of Colorado potato beetle (Coleoptera: Chrysomelidae) to novaluron. *Journal of Economic Entomology* 98: 1685-1693.
- French NM, Heim DC, Kennedy GG. 1992. Insecticide resistance patterns among Colorado potato beetle, *Leptinotarsa decemlineata* (Say) (Coleoptera: Chrysomelidae), populations in North Carolina. *Pesticide Science* 36: 95-100.
- Furlong MJ, Groden E. 2001. Evaluation of synergistic interactions between the Colorado potato beetle (Coleoptera: Chrysomelidae) pathogen *Beauveria bassiana* and the insecticides, imidacloprid and cyromazine. *Journal of Economic Entomology* 94: 344-356.
- Griffith MB, Barrows EM, Perry SA. 2000. Effect of diflubenzuron on flight of adult aquatic insects (Plecoptera, Trichoptera) following emergence during the second year after aerial application. *Journal of Economic Entomology* 93: 1695-1700.

- Hare JD. 1990. Ecology and management of the Colorado potato beetle. *Annual Review of Entomology* 35: 81-100.
- Harris CR, Svec HJ. 1981. Colorado potato beetle resistance to carbofuran and several other insecticides in Quebec. *Journal of Economic Entomology* 74: 421-424.
- Hegazy G, Cock A, Auda M, Degheele D, Decock A. 1989. Diflubenzuron toxicity, effect on the cuticle ultrastructure, chitin and protein content of the Colorado potato beetle *Leptinotarsa decemlineata* (Say) (Coleoptera: Chrysomelidae). *Mededelingen van de Faculteit Landbouwwetenschappen*, 51: 89-101.
- Hughes PB, Dauterman WC, Motoyama N. 1989. Inhibition of growth and development of tobacco hornworm (Lepidoptera: Sphingidae) larvae by cyromazine. *Journal of Economic Entomology* 82: 45-51.
- Kotze AC. 1992. Effects of cyromazine on reproduction and offspring development in *Lucilia cuprina* (Diptera: Calliphoridae). *Journal of Economic Entomology* 85: 1614-1617.
- Lucas E, Giroux S, Demougeost S, Duchesne RM, Coderre D. 2004. Compatibility of a natural enemy, *Coleomegilla maculata lengi* (Col., Coccinellidae) and four insecticides used against the Colorado potato beetle (Col., Chrysomelidae). *Journal of Applied Entomology* 128: 233-239.
- Malinowski H, Pawinska M. 1992. Comparative evaluation of some chitin synthesis inhibitors as insecticides against Colorado potato beetle *Leptinotarsa decemlineata* (Say). *Pesticide Science* 35: 349-353.
- Mota-Sanchez D, Hollingworth RM, Grafius EJ, Moyer DD. 2006. Resistance and cross-resistance to neonicotinoid insecticides and spinosad in the Colorado potato beetle, *Leptinotarsa decemlineata* (Say) (Coleoptera: Chrysomelidae). *Pest management science* 62: 30-37.
- Pourmirza AA. 2005. Local variation in susceptibility of Colorado potato beetle (Coleoptera: Chrysomelidae) to insecticide. *Journal of Economic Entomology* 98: 2176-2180.
- Robertson JL, Preisler HK. 1992. *Pesticide bioassays with arthropods*. CRC Press.
- Root DS, Dauterman WC. 1996. Cyromazine toxicity in different laboratory strains of the tobacco hornworm (Lepidoptera: Sphingidae). *Journal of Economic Entomology* 89: 1074-1079.
- Sirota JM, Grafius E. 1994. Effects of cyromazine on larval survival, pupation and adult emergence of Colorado potato beetle (Coleoptera: Chrysomelidae). *Journal of Economic Entomology* 87: 577-582.
- SPSS. 1999. *SPSS user's manual*. Release 10.0. SPSS, Chicago, IL.
- Tunaz H, Uygun N. 2004. Insect Growth Regulators for Insect Pest Control. *Turkish Journal of Agriculture and Forestry* 28: 377-387.
- Tuttle AF, Ferro DN. 1988. Laboratory evaluation of the insect growth regulator CME-13406 on Colorado potato beetle (Coleoptera: Chrysomelidae). *Journal of Economic Entomology* 81: 654-657.
- Wakgari WM, Giliomee JH. 2003. Natural enemies of three mealybug species (Hemiptera: Pseudococcidae) found on citrus and effects of some insecticides on the mealybug parasitoid *Coccidoxenoides peregrinus* (Hymenoptera: Encyrtidae) in South Africa. *Bulletin of Entomological Research* 93: 243-254.