



Successful nesting by 2 endangered Hawaiian waterbird species in a restored Indigenous wetland agroecosystem

Authors: Harmon, Kristen C., Opie, Eryn N. P., Miner, Ali'i, Paty-Miner, Iokepa, Kukea-Shultz, Jonathan K., et al.

Source: The Wilson Journal of Ornithology, 134(1) : 102-105

Published By: The Wilson Ornithological Society

URL: <https://doi.org/10.1676/20-00064>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

The Wilson Journal of Ornithology 134(1):102–105, 2022

DOI: 10.1676/20-00064

Successful nesting by 2 endangered Hawaiian waterbird species in a restored Indigenous wetland agroecosystem

Kristen C. Harmon,^{1*} Eryn N. P. Opie,¹ Ali'i Miner,^{2,3,4} Iokepa Paty-Miner,^{2,3,4}
Jonathan K. Kukea-Shultz,² Kawika B. Winter,^{1,5} and Melissa R. Price¹

ABSTRACT—The Hawaiian Stilt (*Himantopus mexicanus knudseni*) and Hawaiian Gallinule (*Gallinula galeata sandvicensis*) are federally endangered waterbirds endemic to the Hawaiian Islands. Both species are conservation-reliant; their population persistence is dependent on invasive predator control and removal of invasive plants that degrade habitat. We present observations of successful nesting by one Hawaiian Stilt pair and one Hawaiian Gallinule pair at a

site managed within an adaptive Indigenous agroecological framework on the island of O'ahu, Hawai'i. The Hawaiian Stilt nest, found in February 2019, contained 4 eggs and produced 3 hatchlings, 2 of which were banded and monitored after hatching. The Hawaiian Gallinule nest, found in February 2020, contained 6 eggs and produced 5 hatchlings. Although no individuals were banded from this nest, 2 adults and 2 hatchlings were continuously observed in the nesting area after the eggs hatched. *Lo'i kalo* Hawaiian wetland agroecosystems centered around the cultivation of *kalo* (taro; *Colocasia esculenta*), have the potential to expand Hawaiian waterbird habitat beyond state and federal protected areas. We are aware of unpublished accounts of Hawaiian waterbirds nesting in commercially farmed *lo'i kalo*, but until now, there have been no previously published accounts of native waterbirds breeding in *lo'i kalo* managed as Indigenous agroecosystems. Received 28 May 2020. Accepted 7 December 2021.

¹ Department of Natural Resources and Environmental Management, College of Tropical Agriculture and Human Resources, University of Hawai'i at Mānoa, Honolulu, HI, USA

² Kāko'o 'Ōiwi, He'eia, HI, USA

³ Kuhialoko, Waiawa Kai, HI, USA

⁴ Hawai'i Coral Reef Initiative, University of Hawai'i at Mānoa, Honolulu, HI, USA

⁵ Hawai'i Institute of Marine Biology, University of Hawai'i at Mānoa, He'eia, HI, USA

* Corresponding author: kmcorey@hawaii.edu

Key words: conservation, *Gallinula galeata sandvicensis*, Hawai'i, *Himantopus mexicanus knudseni*, *lo'i*, wetland.

Anidación exitosa de 2 especies de aves acuáticas hawaianas en peligro en un agroecosistema indígena restaurado

RESUMEN (Spanish)—La monjita *Himantopus mexicanus knudseni* y la gallareta *Gallinula galeata sandvicensis* son aves acuáticas federalmente en peligro endémicas a las islas hawaianas. Ambas especies dependen de acciones de conservación; la persistencia de sus poblaciones depende del control de depredadores invasores y de la remoción de plantas invasoras que degradan sus hábitats. Presentamos observaciones de anidación exitosa de una pareja de monjita y una de gallareta en un sitio manejado mediante prácticas indígenas agroecológicas en la isla de O‘ahu, Hawái. El nido de la monjita, encontrado en febrero de 2019, contenía 4 huevos y produjo 3 polluelos, 2 de los cuales fueron anillados y monitoreados después de eclosionar. Aunque ningún individuo de este nido fue anillado, 2 adultos y 2 polluelos fueron continuamente observados en el área del nido después de que los huevos eclosionaron. Los agroecosistemas de humedales hawaianos (l‘oi) tienen el potencial de expandir el hábitat de las aves acuáticas hawaianas más allá de las áreas protegidas estatales y federales. Sabemos que hay reportes de aves acuáticas hawaianas sin publicar en l‘oi comercialmente cultivados, pero hasta ahora no había reportes previos publicados de aves acuáticas nativas en l‘oi bajo manejo como agroecosistemas indígenas.

Palabras clave: conservación, *Gallinula galeata sandvicensis*, Hawái, *Himantopus mexicanus knudseni*, humedal, l‘oi.

The Hawaiian Stilt (*Ae‘o*; *Himantopus mexicanus knudseni*) and Hawaiian Gallinule (*‘Alae ‘ula*; *Gallinula galeata sandvicensis*) are federally endangered waterbirds endemic to the Hawaiian Islands. Both species benefit from conservation actions, as their populations rely on control of invasive plants and predators to remain viable (Reed et al. 2012). Archeological evidence (Burney et al. 2001, Burney and Kikuchi 2006) suggests that *lo‘i kalo* Hawaiian wetland agroecosystems, centered on the cultivation of *kalo* (taro; *Colocasia esculenta*), greatly expanded waterbird habitat in the pre-contact era (Winter et al. 2018b). However, in the 19th century, productive and biologically diverse wetland agroecosystems were transitioned into low-biodiversity monocultures for plantation agriculture, which ultimately contributed to a regime shift in social-ecological systems in Hawai‘i (Winter et al. 2018a, 2018b).

A decline in *lo‘i kalo*, along with a decline in Hawaiian social-ecological systems, an increase in introduced species, and an increase in hunting of waterfowl associated with the introduction of guns to the islands, led to a collapse in native waterbird populations (Kame‘eleihiwa 1992). Today the Hawaiian Stilt and Hawaiian Gallinule are both

habitat-limited species (Reed et al. 1998; van Rees and Reed 2018, 2020), and sea-level rise has the potential to further exacerbate the recovery of their populations (Kane et al. 2014, van Rees and Reed 2018, Harmon et al. 2021). However, restoration of *lo‘i kalo* systems has the potential to preserve Hawaiian cultural practices and provide sustainable food production, while simultaneously increasing suitable habitat for Hawai‘i’s endangered waterbirds (Harmon et al. 2021).

Research conducted in commercial taro plantations on a U.S. Fish and Wildlife Service refuge (Greer 2005, Gee 2007) suggests that *lo‘i kalo* provide foraging and nesting habitat for endangered Hawaiian waterbirds. However, *lo‘i kalo* management practices used in single-objective high-intensity agribusiness models (Evans 2008) include substantial use of fertilizers and hydrological practices that facilitate avian botulism outbreaks in waterbirds. Such approaches to taro cultivation potentially create population sinks that attract waterbirds to breed and forage but result in increased mortality and decreased reproductive success (Battin 2004, Robertson and Hutto 2006, Buderman et al. 2020). In contrast, multi-objective, Indigenous wetland agroecosystems (Winter et al. 2018a) consist of a patchwork of fields in different phases of cultivation (Winter et al. 2018b), secondary crops along field borders (Kurashima and Kirch 2011), and *‘auwai* (irrigation ditches and canals facilitating hydrological regimes that minimize the likelihood of botulism outbreaks), creating habitat mosaics that support a diversity of wetland organisms (Winter et al. 2020). As there are no previously published accounts of native waterbirds breeding in *lo‘i kalo* managed as Indigenous agroecosystems rather than as agribusiness plantations, we present 2 observations of successful nesting by Hawaiian Stilts and Hawaiian Gallinules within 2 *lo‘i kalo* basins at a site managed using Indigenous agroecology practices on the island of O‘ahu, Hawai‘i.

The He‘eia National Estuarine Research Reserve (NERR) on O‘ahu, Hawai‘i (21.4339°N, 157.8112°W) contains an invasive-dominated wetland that is currently being restored for multiple Indigenous wetland agroecology systems including *lo‘i kalo* and aquaculture production. Management of *lo‘i kalo* at the site includes practices that aim to establish self-sustaining waterbird populations with limited human impact. In these restored



Figure 1. A Hawaiian Stilt nest with 4 eggs inside a wet, fallow *lo'i kalo* basin managed by Kāko'o 'Ōiwi within the He'eia National Estuarine Research Reserve on O'ahu. Three of 4 eggs hatched, and all 3 chicks survived at least 1 year, remaining with the adult pair in the immediate area.

lo'i kalo, one Hawaiian Stilt and one Hawaiian Gallinule nest were discovered in 2019 and 2020, respectively, during a waterbird survey. The Hawaiian Stilt nest with 4 eggs was found inside a wet, fallow *lo'i kalo* basin within the He'eia NERR on 22 May 2019. The nest had been constructed on a small mudflat within the basin, surrounded by sparse vegetation (Fig. 1). The nest was monitored with a Bushnell No-Glow Aggressor HD Trophy Camera, consistent with U.S. Fish and Wildlife Service Threatened and Endangered Species Permit #TE-25955C-1. The camera was programmed to take 2 images back-to-back immediately upon infrared motion activation with a 5 s delay between each successive activation. One control photo was taken every hour using field scan mode. The camera was checked weekly for battery life and data card retrieval and was removed immediately after the eggs hatched. The Hawaiian Gallinule nest with 6 eggs was found on 27 February 2020 inside a flooded *lo'i kalo* basin surrounded by 'ohā (*kalo* offshoots) also within the He'eia NERR (Fig. 2). The nest was monitored weekly during observational waterbird surveys.

Nest camera images of the Hawaiian Stilt nest confirmed that 3 of the 4 eggs hatched from 10 to 12 June 2019. Two of the chicks were captured and fitted with U.S. Geological Survey bands, as well as PVC color bands for individual identification, at ca. 4 weeks of age (pre-fledge). The 2 breeding adults and 3 offspring survived past the original nesting observation and continued to be observed foraging and resting within components of the agroecosystem, including basins at various *kalo* development stages, berms, and nearby



Figure 2. A Hawaiian Gallinule nest with 6 eggs inside a flooded *lo'i kalo* basin managed by Kāko'o 'Ōiwi within the He'eia National Estuarine Research Reserve on O'ahu. Five of the 6 eggs hatched, and 2 chicks survived a minimum of 3 months, remaining in the immediate area.

wetlands and intertidal mud flats. On 13 March 2020 the Hawaiian Gallinule nest was found empty, and 5 gallinule chicks were observed near the nest site. Subsequent field surveys in March 2020 confirmed that the 2 adults and at least 2 chicks were still onsite and regularly utilizing *lo'i kalo* basins, nearby streams, and nearby wetlands. These individuals are not banded, but we believe our observations are of the same family, as gallinules are highly territorial (Chang 1990) and no other gallinule nests were observed during this time.

As sea-level rise is projected to negatively impact both the Hawaiian Stilts (Harmon et al. 2021) and Hawaiian Gallinules (van Rees and Reed 2018), the acquisition or creation of new waterbird habitat will be critical to achieve recovery goals for endangered endemic Hawaiian waterbird populations into the future. This study supports reported observations of waterbirds nesting and successfully fledging since 2010 within *lo'i kalo* basins at He'eia (JKK, pers. obs.; USFWS 2019), and further supports the notion that restoring Hawaiian wetland agroecosystems, under an Indigenous resource management framework, as suggested by Harmon et al (2021), does indeed have the potential to increase food production while expanding endangered waterbird habitat (Harmon et al. 2021). Moreover, contemporary Indigenous management of *lo'i kalo* often includes actions that are analogous with endangered species management in protected

areas, such as control of invasive plants and predators (Greer 2005), management of hydrological conditions to meet waterbird life history needs (USDA 2009), and management of human impacts, all of which are critical for Hawaiian waterbird conservation (VanderWerf 2012, Underwood et al. 2014). Thus, Indigenous management of Hawaiian agroecosystems, coupled with predator control, offers a sustainable landscape-level solution for expanding conservation of these species beyond state and federal managed protected areas.

Acknowledgments

We thank Kāko'o 'Ōiwi staff, J.L. Idle, and L.C. Zavas for their comments and support. Funding for this publication was made possible by the He'eia National Estuarine Research Reserve (award #NAI9NOS4200078), the State of Hawai'i Department of Land and Natural Resources (award #011068-00001), and Sustainable Resources Group International Inc. (award #011328-00001).

Literature cited

- Battin J. 2004. When good animals love bad habitats: Ecological traps and the conservation of animal populations. *Conservation Biology*. 18:1482–1491.
- Buderman FE, Devries JH, Koons DN. 2020. Changes in climate and land use interact to create an ecological trap in a migratory species. *Journal of Animal Ecology*. 89:1961–1977.
- Burney DA, James HF, Burney LP, Olson SL, Kikuchi W, et al. 2001. Fossil evidence from a diverse biota from Kauai and its transformation since human arrival. *Ecological Monographs*. 71:615–641.
- Burney DA, Kikuchi WA. 2006. Millennium of human activity at Makauwahi Cave, Mahaulepu, Kauai. *Human Ecology*. 34:219–247.
- Chang PR. 1990. Strategies for managing endangered waterbirds on Hawaiian national wildlife refuges [master's thesis]. Amherst (MA): University of Massachusetts.
- Evans D. 2008. Taro mauka to makai: A taro production and business guide for Hawai'i growers. Honolulu (HI): University of Hawaii Press.
- Geer HK. 2007. Habitat characteristics of refuge wetlands and taro lo'i used by endangered waterbirds at Hanalei National Wildlife Refuge, Hawai'i [master's thesis]. Vermillion (SD): South Dakota State University.
- Greer NM. 2005. Ethnoecology of taro farmers and their management of Hawaiian wetlands and endangered waterbirds in taro agroecosystems [dissertation]. Seattle (WA): University of Washington.
- Harmon K, Winter KB, Kurashima N, Fletcher CH, Kane HH, Price MR. 2021. The role of Indigenous practices in expanding waterbird habitat in the face of rising seas. *Anthropocene*. 34:100293.
- Kame'elehiwa L. 1992. Native land and foreign desires. Honolulu (HI): Bishop Museum Press.
- Kane H, Fletcher CH, Frazer LN, Barbee MM. 2014. Critical elevation levels for flooding due to sea-level rise in Hawaii. *Regional Environmental Change*. 15:1679–1687.
- Kurashima N, Kirch PV. 2011. Geospatial modeling of pre-contact Hawaiian production systems on Molokai Island, Hawaiian Islands. *Journal of Archaeological Science*. 38:3662–3674.
- Reed JM, DesRochers DW, VanderWerf EA, Scott MJ. 2012. Long-term persistence of Hawaii's endangered avifauna through conservation-reliant management. *BioScience*. 62:881–892.
- Reed JM, Elphick CS, Oring LW. 1998. Life-history and viability analysis of the endangered Hawaiian Stilt. *Biological Conservation*. 84:35–45.
- Robertson B, Hutto R. 2006. A framework for understanding ecological traps and an evaluation of existing evidence. *Ecology*. 87:1075–1085.
- Underwood JG, Silbernagle M, Nishimoto M, Uyehara K. 2014. Non-native mammalian predator control to benefit endangered Hawaiian waterbirds. *Proceedings of the Vertebrate Pest Conference 26*. Davis (CA): University of California Davis; p. 32–39.
- [USDA] U.S. Department of Agriculture. 2009. Practices to enhance native wildlife habitat on wetland taro farms. Washington (DC): USDA Natural Resources Conservation Service, Pacific Islands Area.
- [USFWS] U.S. Fish and Wildlife Service. 2019. Biological Opinion for the NOAA He'eia National Estuarine Research Reserve, O'ahu, Hawai'i. Honolulu (HI): U.S. Fish and Wildlife Service.
- VanderWerf EA. 2012. Hawaiian bird conservation action plan. Honolulu (HI): Pacific Rim Conservation.
- van Rees CB, Reed JM. 2018. Predicted effects of landscape change, sea level rise, and habitat management on the extirpation risk of the Hawaiian Common Gallinule (*Gallinula galeata sandvicensis*) on the island of O'ahu. *PeerJ*. 6:e4990.
- van Rees CB, Reed JM. 2020. Multiple sources of evidence for density dependence in the endangered Hawaiian Stilt (*Himantopus mexicanus knudseni*). *Population Ecology*. 62:207–219.
- Winter KB, Beamer K, Vaughan M, Friedlander AM, Kido MH, et al. 2018a. The Moku System: Managing biocultural resources for abundance within social-ecological regions in Hawai'i. *Sustainability* (Special issue on Biocultural Restoration in Hawai'i). 10:3554.
- Winter KB, Lincoln NK, Berkes F. 2018b. The social-ecological keystone concept: A metaphor for understanding the structure and function of a biocultural system. *Sustainability* (Special issue on Biocultural Restoration in Hawai'i). 10:3294.
- Winter KB, Lincoln NK, Berkes F, Alegado RA, Kurashima N, et al. 2020. Ecomimicry in Indigenous resource management: Optimizing ecosystem services to achieve resource abundance with examples from Hawai'i. *Ecology and Society*. 25(2):26.