

First Record of *Cydia tonosticha* (Meyrick) (Tortricidae) from Chile and a New Host Plant

Author: Bobadilla, Dante

Source: The Journal of the Lepidopterists' Society, 69(4) : 331-333

Published By: The Lepidopterists' Society

URL: <https://doi.org/10.18473/lepi.69i4.a1>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

FIRST RECORD OF *CYDIA TONOSTICHA* (MEYRICK) (TORTRICIDAE) FROM CHILE AND A NEW HOST PLANT

Additional key words: *Cydia largo*, Fabaceae, Neotropical, *Prosopis alba*

Cydia Hübner, 1825 is a cosmopolitan and highly diverse genus of Tortricidae (Gilligan et al. 2014) that is currently represented in Chile by two species (Vargas & Parra 2006, Razowski & Pelz 2010): *C. pomonella* (Linnaeus, 1758) and *C. largo* Heppner, 1981. *Cydia pomonella*, commonly known as the codling moth, is a widely distributed pest of apple, pear, and walnut (Gilligan & Epstein 2012); in Chile it is a primary pest of apple (Razowski & Pelz 2010). *Cydia largo*, whose larvae are associated with Fabaceae, was described from southern Florida and was also recorded from Cuba (Heppner 1981); later it was reported from the coastal valleys of the Atacama Desert of Chile (Vargas & Parra 2006), where its larvae are florivorous on *Acacia macracantha* (Fabaceae) (Vargas & Parra 2006, 2009).

As part of a survey of Lepidoptera associated with native plants in the coastal valleys of the Atacama Desert of northern Chile, some seed-feeding larvae were detected in pods of *Prosopis alba* (Fabaceae) in October 2014 in the Lluta Valley, Arica Province. Infested pods were collected and brought to the laboratory in plastic vials and were kept at room temperature to obtain adults. Five adults were reared, and they were identified as *Cydia tonosticha* (Meyrick, 1922) based on morphology of the male and female genitalia (Lima 1952, Heppner et al. 2009, Razowski 2011) (Fig. 1–3).

Cydia tonosticha was described from Amazonas, Brazil, and apparently it is widely distributed in the Neotropics; it has been reported from Panama, Peru, and Venezuela (Heppner et al. 2009, Razowski 2011). Its synonym (*Laspeyresia cassiana* Lima, 1952) also has a Brazilian type locality in Rio de Janeiro State. Larvae of *C. tonosticha* have been recorded as seed-feeders in pods of four species of Fabaceae belonging to the genera *Cassia* and *Stryphnodendron*, based on sampling performed in Brazil and Panama (Lima 1952, Becker 1971, Nomura et al. 1976, Pentead-Dias et al. 2008, Razowski 2011). In the collection of the National Museum of Natural History, Washington, DC, USA (USNM), there are specimens of *C. tonosticha* reared from *Acacia farnesiana* (Panama), *Cassia fistula* (Brazil), *Cassia moschata* (Panama), *Cassia grandis* (Panama), *Senna bacillaris* (Brazil), and *Senna rugosa* (Brazil). Surprisingly, *Punica granatum* (Punicaceae) was recently recorded as a host for *C. tonosticha* in Peru (Heppner et al. 2009). In addition, the wasp *Pseudophanerotoma* (*Pseudophanerotoma*) *alvarengai* Zettel, 1990

(Hymenoptera: Braconidae: Cheloninae) was recorded parasitizing larvae of *C. tonosticha* in Brazil (Pentead-Dias et al. 2008).

This is the first record of *C. tonosticha* from Chile, adding one more country to the distribution range of this widespread Neotropical species. The southern limit previously reported in Peru is in Omate, Departamento de Moquegua, about 200 km north of the Lluta Valley (Heppner et al. 2009). It remains unknown whether the presence of *C. tonosticha* in Chile is a result of a recent range expansion or the species is native to the coastal valleys of the Atacama Desert. Although the original description of *C. tonosticha* was based on Brazilian specimens, its wide distribution raises the question about its evolutionary origin, as the type locality does not necessarily represent the geographic origin of the species (Gonçalves et al. 2015); molecular studies at the population level are required in order to explore this scenario (Valade et al. 2009).

This is the first record of a species of *Prosopis* as a host plant for larvae of *C. tonosticha*, adding one more genus to the family most commonly recorded for this species (Lima 1952, Becker 1971, Pentead-Dias et al. 2008, Razowski 2011). Apparently, Fabaceae is an important host plant family for Tortricidae in northern Chile (Clarke 1987, Vargas & Parra 2006, Vargas 2011).

Finally, the discovery of *C. tonosticha* in northern Chile, together with other recent additions to the Chilean Tortricidae (Vargas & Parra 2006, Vargas 2011, 2012,




FIG. 1–3. *Cydia tonosticha* (Meyrick, 1922). 1. Female adult in dorsal view; scale bar = 1 mm. 2. Male genitalia in ventral view; scale bar = 0.1 mm. 3. Female genitalia in ventral view; scale bar = 0.1 mm.

Vargas et al. 2015), highlight the importance of surveys to understand the diversity of this family throughout these arid landscapes.

Material examined. Chile, Arica. Three males, two females: Lluta, Arica, Chile, November 2014, D. Bobadilla coll., seed-feeder larvae in pods of *Prosopis alba*, October 2014 (IDEA). Vouchers will be deposited in the Colección Entomológica de la Universidad de Tarapacá (IDEA), Arica, Chile.

ACKNOWLEDGMENTS

The authors thank John W. Brown (USNM) for important comments and suggestions on a preliminary version and for kindly provide the data on host plants of *C. tonosticha* in USNM; Clorinda Vergara-Cobian (UNALM) for sending literature; Patricia Jiménez Guarda (SAG) for encouragement to publish this record; Ricardo Mendoza Mamani (UTA) for field assistance; Marcelo Vargas-Ortiz (UTA) for editing the figures; and Lafayette Eaton for checking the English. This study was supported by project DGI 9710-14, Dirección General de Investigación, Universidad de Tarapacá.

LITERATURE CITED

- BECKER, V. O. 1971. Microlepidópteros que vivem nas plantas cultivadas no Brasil. II. O nome correto da lagarta das favas da *Cassia fistula* L. (Lepidoptera, Tortricidae). Bol. Univ. Fed. Paraná Zool. 4: 45–46.
- CLARKE, J.F.G. 1987. Two new *Cryptophlebia* Walsingham from Chile (Lepidoptera: Tortricidae). Acta Entomol. Chilena 14: 7–12.
- GILLIGAN, T. M. & M. E. EPSTEIN. 2012. TortAI, Tortricids of Agricultural Importance to the United States (Lepidoptera: Tortricidae). Identification Technology Program (ITP), USDA/APHIS/PPQ/CPHST, Fort Collins, CO. <http://idtools.org/id/leps/tortai/> (Accessed 23 January 2015)
- GONÇALVES, G. L., G. R. P. MOREIRA, R. BRITO & H. A. VARGAS. 2015. Stranger in a known land: Bayesian analysis confirms the presence of an Australian leaf miner in the Chilean Atacama Desert. Bioinvasions Rec. 4: 67–73.
- HEPPNER, J. B. 1981. A new *Cydia* (Lepidoptera: Tortricidae) from Florida and Cuba. J. Lepid. Soc. 35: 278–280.
- HEPPNER, J. B., C. E. VERGARA-COBIAN & M. NARREA-CONGO. 2009. *Cydia tonosticha* in Peru (Lepidoptera: Tortricidae). Lepid. Novae 2: 45–47.
- NOMURA, H., A. B. GUSMAN, D. L. CORRÊA & L. NEMOTO. 1976. Desenvolvimento de *Laspeyresia* sp. (Lepidoptera, Grapholitidae), nova praga de sementes do barbatimão, *Stryphnodendron barbatimann* (Velloso) (Leguminosae, Mimosoideae). Científica 4: 6–13.
- PENTEADO-DIAS, A. M., A. R. NASCIMENTO & M. M. DIAS. 2008. The description of the male and the first host data of *Pseudophanerotoma* (*Pseudophanerotoma*) *alvarengai* Zettel, 1990 (Hymenoptera: Braconidae: Cheloniinae). Zool. Med. Leiden 82: 401–405.
- RAZOWSKI, J. 2011. New species, new genera, and new combinations of Grapholitini (Lepidoptera: Tortricidae) from the Neotropical Region. Acta Zool. Cracov. 53A: 37–101.
- RAZOWSKI, J. & V. PELZ. 2010. Tortricidae from Chile (Lepidoptera: Tortricidae). SHILAP Revta. Lepid. 38: 5–55.
- VALADE, R., M. KENIS, A. HERNANDEZ-LOPEZ, S. AUGUSTIN, N. MARI MENA, E. MAGNOUX, R. ROUGERIE, F. LAKATOS, A. ROQUES & C. LOPEZ-VAAMONDE. 2009. Mitochondrial and microsatellite DNA markers reveal a Balkan origin for the highly invasive horse-chestnut leaf miner *Cameraria ohridella* (Lepidoptera, Gracillariidae). Mol. Ecol. 18: 3458–3470.
- VARGAS, H. A. 2011. A new species of *Eccopsis* Zeller (Lepidoptera, Tortricidae) from the coastal valleys of northern Chile, with the first continental record of *E. galapagana* Razowski & Landry. Rev. Bras. Entomol. 55: 216–218.
- VARGAS, H. A. 2012. *Strepsicrates smithiana* Walsingham (Lepidoptera, Tortricidae): first record from Chile and a newly documented host plant. Rev. Bras. Entomol. 56: 381–382.
- VARGAS, H. A. & L. E. PARRA. 2006. Nuevos registros de distribución y notas biológicas de *Cydia largo* Heppner (Lepidoptera: Tortricidae). Gayana 70: 293–294.
- VARGAS, H. A. & L. E. PARRA. 2009. Prospección de lepidópteros anatófagos asociados a *Acacia macracantha* Willd. (Fabaceae) en el norte de Chile. Rev. Bras. Entomol. 53: 291–293.
- VARGAS, H. A., P. POLLO, D. S. BASILIO, G. L. GONÇALVES & G. R. P. MOREIRA. 2015. A new cecidogenous species of *Eugnosta* Hübner (Lepidoptera: Tortricidae) associated with *Baccharis salicifolia* (Asteraceae) in the northern Chilean Atacama Desert: life-history description and phylogenetic inferences. Zootaxa 3920: 265–280.

DANTE BOBADILLA, *Departamento de Recursos Ambientales, Facultad de Ciencias Agronómicas, Universidad de Tarapacá, Casilla 6-D, Arica, Chile.*
 HÉCTOR A. VARGAS (*corresponding author*), *Departamento de Recursos Ambientales, Facultad de Ciencias Agronómicas, Universidad de Tarapacá, Casilla 6-D, Arica, Chile; email: havargas@uta.cl; lepvargas@gmail.com*

Submitted for publication 31 March 2015; revised and accepted 22 April 2015.

