

The Rediscovery Of Joiceya Praeclarus Talbot 1928 (Lepidoptera: Riodinidae), More Than 80 Years after Its Description

Authors: Greve, Roberto R., Callaghan, Curtis, Kaminski, Lucas A., and Freitas, André V. L.

Source: The Journal of the Lepidopterists' Society, 67(1) : 56-57

Published By: The Lepidopterists' Society

URL: <https://doi.org/10.18473/lepi.v67i1.a7>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

THE REDISCOVERY OF *JOICEYA PRAECLARUS* TALBOT 1928 (LEPIDOPTERA: RIODINIDAE), MORE THAN 80 YEARS AFTER ITS DESCRIPTION

Additional key words: Atlantic Forest, Brazil, cerrado, endangered species, Nymphidiini, Parque Nacional do Iguaçu

The family Riodinidae is the third most speciose butterfly group in the Neotropics, with more than 1,300 described species in this region (Callaghan & Lamas 2004), and shows an extraordinary diversity of morphological, ecological and life history traits (reviewed in Brown 1993a; DeVries 1997). Although in the past 30 years there have been advances in riodinid taxonomy (see comments in Hall 2005: 3–5), there have been few studies on the ecology and natural history of most riodinid species. Our poor understanding of riodinid biology is partly a result of the apparent rarity of many species, which may be a consequence of them occurring in small, localized populations and/or that some inhabit the forest canopy, are rarely attracted to baits, making them difficult to sample (Callaghan 1978; Brown 1993a; DeVries et al. 1994; Hall & Willmott 2010). Some highly endemic species may therefore be vulnerable to habitat destruction, but the difficulty in observing or sampling them complicates assessing their natural population densities. Consequently, any biological observations of rare riodinid species are potentially important.

Joiceya praeclarus Talbot, 1928 (Fig. 1) is a good example of the above situation. This monotypic genus was described from two males collected in two cerrado areas in Tombador and Cuiabá, in Mato Grosso, central Brazil (Talbot 1928), and despite intensive efforts, it has not been observed for more than 80 years following its description, and its biology and natural history remain unknown (Brown 1993b). Because of its apparent rarity and restricted geographical distribution, *J. praeclarus* was the first Riodinidae to appear on a red list of endangered species (Bernardes et al. 1989), and has been evaluated as “Endangered” by the IUCN (IUCN 2011).

On 9 September 2011, at 11:20 hs, a single male of *J. praeclarus* was observed and collected at Foz do Iguaçu, Paraná, Brazil (25°33'S 54°31'W, 205 m a.s.l.), about 1,200 km from its type locality in Mato Grosso. The site consisted of a small patch of secondary riparian forest along the Köhlenberger stream, inside a matrix of old abandoned pastures and country houses, about 8.4 km from Iguaçu National Park. The male was flying about 1 m high, and landing upside down beneath leaves with wings closed over the body. The individual has been collected for photographs (Fig. 1) and was deposited at the Museu de Zoologia “Adão José Cardoso” (ZUEC),




FIG. 1. Adult male of *Joiceya praeclarus*. **A**, perched above leaves; **B**, about to take off—note the dorsal blue metallic coloration. Scale bars = 0.5 mm.

Universidade Estadual de Campinas, Campinas, São Paulo, Brazil. Despite intensive subsequent searches, no other individuals were observed on following days at the site. The observation of *J. praeclarus* in Foz do Iguaçu is of obvious importance for its conservation assessment, given the paucity of data previously available for the species. Since the conservation status of the species is based on the assumption of its occurrence at a single locality, this status should be revised from “Endangered” to “Data deficient”. It now seems possible that *J. praeclarus* may occur in similar habitats from Mato Grosso to Paraná, and obviously more observations are needed to assess its habitat requirements and true distribution. The fact that the individual *J. praeclarus* was observed in a small, secondary forest fragment suggests it may be a much more tolerant species than formerly assumed, with populations potentially

persisting in a variety of different habitats in the region of Foz do Iguaçu. In this case the rarity of this species might be explained by its persistence in small localized populations, making adults seldom observed (additionally, adults can fly in restricted periods of the day or of the year, as known for several riodinids), or it is simply a canopy species which is very rarely collected in the understory, as is the case with several rare riodinids which usually fly very high in the forest (DeVries 1997; Hall & Willmott 2010).

ACKNOWLEDGEMENTS

We thank Keith Willmott and Phil DeVries for critical reading of the manuscript. RRG thanks to his wife Márcia dos Anjos, who helped in field work, and also to the volunteers of CEAEC (Center for Higher Studies of Conscientiology), who allowed field work in its property. LAK thanks Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP 10/51340-8). AVLF thanks FAPESP (grant 04/05269-9) and CNPq (fellowship 302585/2011-7). This publication is part of the RedeLep “Rede Nacional de Pesquisa e Conservação de Lepidópteros” SIS-BIOTA-Brasil/CNPq (563332/2010-7) and BIOTA-FAPESP program (11/50225-3).

LITERATURE CITED

- BERNARDES, A.T., A.B.M. MACHADO, & A.B. RYLANDS. 1990. Fauna Brasileira Ameaçada de Extinção. Fundação Biodiversitas / IBAMA, Belo Horizonte, Brazil. 65 pp.
- BROWN JR. K.S. 1993A. Neotropical Lycaenidae—an overview, pp. 45–61. *In*: New, T.R. (ed.). Conservation biology of Lycaenidae (Butterflies). IUCN, Gland, Switzerland.
- . 1993B. Selected Neotropical species, pp. 146–149. *In*: New, T.R. (ed.). Conservation biology of Lycaenidae (Butterflies). IUCN, Gland, Switzerland.
- CALLAGHAN, C.J. 1978. Studies on restinga butterflies. II. Notes on the population structure of *Menander felsina* (Riodininae). *J. Lepid. Soc.* 32: 87–94.
- CALLAGHAN, C.J. & G. LAMAS. 2004. Riodinidae, pp. 141–170. *In*: Lamas, G. (ed.), Checklist: Part 4A. Hesperioidea—Papilionoidea. *In*: Heppner, J.B. (ed.), Atlas of Neotropical Lepidoptera. Scientific Publishers, Gainesville, FL.
- DEVRIES, P.J. 1997. The butterflies of Costa Rica, Vol. 2, Riodinidae. Princeton University Press, Princeton, NJ, 288 pp.
- DEVRIES, P.J., I.A. CHACON & D. MURRAY. 1994. Toward a better understanding of host use biodiversity in riodinid butterflies (Lepidoptera). *J. Res. Lepid.* 31: 103–126.
- IUCN 2011. IUCN Red List of Threatened Species. Version 2011.2. <www.iucnredlist.org>. Downloaded on 21 February 2012.
- HALL, J.P.W. 2005. A phylogenetic revision of the Napaena (Lepidoptera: Riodinidae: Mesosemiini). The Entomological Society of Washington, Washington, DC. vi + 235 pp.
- HALL, J.P.W. & K.R. WILLMOTT. 2010. Description of a new *Lucillella* species Riodinidae: Symmachiini discovered in the eastern Andes of Ecuador using the single rope canopy access technique. *J. Lepid. Soc.* 64: 139–146.
- TALBOT, G. 1928. List of Rhopalocera collected by Mr. C.L. Collette in Matto Grosso, Brazil. *Bull. Hill Mus.* 2: 192–220.
- ROBERTO R. GREVE, *Rua da Cosmoética, 1847 casa 32 - Bairro Cognópolis Foz do Iguaçu, CEP 85853-755, Paraná, Brazil*, CURTIS CALLAGHAN, *Casa Pica Pau, Floresta de la Sabana, Cra. 7 137-04, Bogotá, Colombia*, LUCAS A. KAMINSKI & ANDRÉ V. L. FREITAS, *Departamento de Biologia Animal e Museu de Zoologia, Instituto de Biologia, Universidade Estadual de Campinas, C.P. 6109, CEP 13083-970, Campinas, São Paulo, Brazil; email: baku@unicamp.br*

Received for publication 15 March 2012; revised and accepted 11 April 2012.