

Frederickia nom. n., a new replacement name for the moth genus Rindgea Ferguson, 2008 (Lepidoptera: Geometridae)

Authors: Rajaei, Hossein, Hausmann, Axel, and Sihvonen, Pasi

Source: Integrative Systematics: Stuttgart Contributions to Natural History, 5(2) : 203-204

Published By: Stuttgart State Museum of Natural History

URL: <https://doi.org/10.18476/2022.310840>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SHORT COMMUNICATION

Frederickia nom. n., a new replacement name for the moth genus *Rindgea* Ferguson, 2008 (Lepidoptera: Geometridae)HOSSEIN RAJAEI¹, AXEL HAUSMANN² & PASI SIHVONEN³

Abstract

A new replacement name, *Frederickia* nom. n., is proposed for the preoccupied genus name *Rindgea* Ferguson, 2008 (Lepidoptera: Geometridae), and the 14 species previously in that genus are combined with the new name, as follows: *Frederickia s-signata* (Packard, 1873), **comb. n.**; *Frederickia hypaethrata* (Grote, 1881), **comb. n.**; *Frederickia cyda* (Druce, 1893), **comb. n.**; *Frederickia maricopa* (Hulst, 1898), **comb. n.**; *Frederickia nigricomma* (Warren, 1904), **comb. n.**; *Frederickia parcata* (Grossbeck, 1908), **comb. n.**; *Frederickia subterminata* (Barnes & McDunnough, 1913), **comb. n.**; *Frederickia stipularia* (Barnes & McDunnough, 1913), **comb. n.**; *Frederickia flaviterminata* (Barnes & McDunnough, 1913), **comb. n.**; *Frederickia ballandrata* (Wright, 1923), **comb. n.**; *Frederickia indeterminata* (McDunnough, 1939), **comb. n.**; *Frederickia piccoloi* (Rindge, 1976), **comb. n.**; *Frederickia disparcata* (Ferguson, 2008), **comb. n.**; and *Frederickia prolificata* (Ferguson, 2008), **comb. n.**

Key words: Ennominae, *Eubolia cyda*, Macariini, moth, nomenclature.

Zusammenfassung

Ein Ersatzname, *Frederickia* nom. n., wird für den bereits besetzten Gattungsnamen *Rindgea* Ferguson, 2008 (Lepidoptera: Geometridae) vorgeschlagen. Ebenso werden die 14 bisher in dieser Gattung enthaltenen Arten wie folgt mit dem neuen Namen kombiniert: *Frederickia s-signata* (Packard, 1873), **comb. n.**; *Frederickia hypaethrata* (Grote, 1881), **comb. n.**; *Frederickia cyda* (Druce, 1893), **comb. n.**; *Frederickia maricopa* (Hulst, 1898), **comb. n.**; *Frederickia nigricomma* (Warren, 1904), **comb. n.**; *Frederickia parcata* (Grossbeck, 1908), **comb. n.**; *Frederickia subterminata* (Barnes & McDunnough, 1913), **comb. n.**; *Frederickia stipularia* (Barnes & McDunnough, 1913), **comb. n.**; *Frederickia flaviterminata* (Barnes & McDunnough, 1913), **comb. n.**; *Frederickia ballandrata* (Wright, 1923), **comb. n.**; *Frederickia indeterminata* (McDunnough, 1939), **comb. n.**; *Frederickia piccoloi* (Rindge, 1976), **comb. n.**; *Frederickia disparcata* (Ferguson, 2008), **comb. n.**; und *Frederickia prolificata* (Ferguson, 2008), **comb. n.**

FERGUSON (2008: 322), in his revision of the North American Geometridae moth fauna, described a new genus, *Rindgea* Ferguson, 2008, and fixed as its type species *Eubolia cyda* Druce, 1893 (Fig. 1), by original designation. He included 14 species in the new genus and classified it in the subfamily Ennominae and tribe Macariini. The species of this genus feed on desert Fabaceae and occur mainly in the south-western USA and Mexico (FERGUSON 2008). The genus was named to honour American lepidopterist FREDERICK H. RINDGE (1921–1996).

During the preparation of the “Online taxonomic facility of geometrid moths” (see RAJAEI et al. 2022), it was noted that *Rindgea* Ferguson, 2008 is a junior homonym of *Rindgea* Johnson, 1993, a New World genus in the butterfly family Lycaenidae. *Rindgea* Johnson, 1993 is currently considered as a junior synonym of *Nicolaea* Johnson, 1993 (LAMAS 2004).

As requested by the International Code of Zoological Nomenclature (ICZN 1999, Article 60), we propose *Frederickia* nom. n. as a new replacement name for *Rindgea* Ferguson, 2008, respecting the original intent behind the naming of this genus by FERGUSON (2008).

In consequence of the new replacement name for *Rindgea*, the following new combinations are established (original genus given in parentheses):

- Frederickia s-signata* (Packard, 1873) (*Macaria*), **comb. n.**
- Frederickia hypaethrata* (Grote, 1881) (*Phasiane*), **comb. n.**
- Frederickia cyda* (Druce, 1893) (*Eubolia*), **comb. n.**
- Frederickia maricopa* (Hulst, 1898) (*Diastictis*), **comb. n.**
- Frederickia nigricomma* (Warren, 1904) (*Semiothisa*), **comb. n.**
- Frederickia parcata* (Grossbeck, 1908) (*Sciagraphia*), **comb. n.**
- Frederickia subterminata* (Barnes & McDunnough, 1913) (*Macaria*), **comb. n.**
- Frederickia stipularia* (Barnes & McDunnough, 1913) (*Macaria*), **comb. n.**
- Frederickia flaviterminata* (Barnes & McDunnough, 1913) (*Macaria*), **comb. n.**
- Frederickia ballandrata* (Wright, 1923) (*Phasiane*), **comb. n.**
- Frederickia indeterminata* (McDunnough, 1939) (*Semiothisa*), **comb. n.**
- Frederickia piccoloi* (Rindge, 1976) (*Semiothisa*), **comb. n.**
- Frederickia disparcata* (Ferguson, 2008) (*Rindgea*), **comb. n.**
- Frederickia prolificata* (Ferguson, 2008) (*Rindgea*), **comb. n.**




Fig. 1. *Frederickia cyda* (Druce, 1893), which FERGUSON (2008) designated as the type species of the genus *Rindgea* Ferguson, 2008. (Photo by ANDY REAGO and CHRISSY McCLARREN, taken in USA, Texas, Estero Llano Grande State Park on 1 March 2015; photo licensed under Creative Commons Attribution 2.0 Generic license and available from Wikipedia at https://en.wikipedia.org/wiki/Rindgea_cyda)


These new combinations will be applied in the “Online taxonomic facility of geometrid moths” (see RAJAEI et al. 2022).

Acknowledgements

We appreciate VALERIE LEVESQUE-BEAUDIN (University of Guelph, Canada) for correction of related data in the BOLD

Authors' addresses:

¹State Museum of Natural History Stuttgart, Rosenstein 1, D-70191 Stuttgart, Germany; e-mail (corresponding author): hossein.rajaei@smns-bw.de;  <https://orcid.org/0000-0002-3940-3734>

²SNSB-Zoologische Staatssammlung München, Münchhausenstr. 21, D-81247 Munich, Germany;  <https://orcid.org/0000-0002-0358-9928>

³Finnish Museum of Natural History, P. O. Box 17, Pohjoinen Rautatiekatu 13, FI-00014 University of Helsinki, Finland;  <https://orcid.org/0000-0003-2237-9325>

ZooBank registration: <https://zoobank.org/References/EDB6ED6E-DE3F-42E1-ADC8-1062BDADC627>

Manuscript received: 09.XII.2022; accepted: 11.XII.2022.

database. We are thankful to reviewers ALBERTO ZILLI (Natural History Museum, London) and DOMINIC WANKE (Staatliches Museum für Naturkunde Stuttgart, Stuttgart) for their critical review of the submitted version of this paper and their constructive comments.

References

- FERGUSON, D. C. (2008): Geometroidea, Geometridae (part): Ennominae (part): Abraxini, Cassymini, Macariini. – In: HODGES, R. W. et al. (eds.): *The Moths of North America. Fascicle 17.2*, 431 pp.; Bakersfield, California (The Wedge Entomological Research Foundation).
- INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE [ICZN] (1999) *International Code of Zoological Nomenclature*. Fourth edition, xxix + 306 pp.; London (International Trust for Zoological Nomenclature). [Available with relevant declarations and amendments from: <https://www.iczn.org/the-code/the-code-online/>]
- JOHNSON, K. (1993): A new Neotropical genus of branded Angulopine-like hairstreak butterflies (Lepidoptera, Lycaenidae, Theclinae). – *Reports of the Museum of Natural History, University of Wisconsin-Stevens Point* **40**: 1–7.
- LAMAS, G. (2004): Checklist: Part 4A. Hesperioidea – Papiilionoidea. – In: LAMAS, G. (ed.): *Atlas of Neotropical Lepidoptera*, 439 pp.; Gainesville, Florida (Scientific Publishers).
- RAJAEI, H., HAUSMANN, A., SCOBLE, M., WANKE, D., PLOTKIN, D., BREHM, G., MURILLO-RAMOS, L. & SIHVONEN, P. (2022): An online taxonomic facility of Geometridae (Lepidoptera), with an overview of global species richness and systematics. – *Integrative Systematics* **5** (2): 145–192. <https://doi.org/10.18476/2022.577933>