



Diurnal Primates in Sri Lanka and People's Perception of Them

Authors: Nahallage, Charmalie A. D., Huffman, Michael A., Kuruppu, Nadeera, and Weerasingha, Tharindi

Source: Primate Conservation, 23(1) : 81-87

Published By: Conservation International

URL: <https://doi.org/10.1896/052.023.0109>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Diurnal Primates in Sri Lanka and People's Perception of Them

Charmalie A. D. Nahallage^{1,2}, Michael A. Huffman², Nadeera Kuruppu³ and Tharindi Weerasingha⁴

¹Department of Sociology and Anthropology, University of Sri Jayawardenepura, Colombo, Sri Lanka

²Department of Social Systems Evolution, Primate Research Institute, Kyoto University, Japan

³Department of Wildlife Conservation, Colombo, Sri Lanka

⁴Department of Social Welfare, Divisional Secretariat Department, Matara, Sri Lanka

Abstract: Five species of non-human primates occur in Sri Lanka—the toque macaque (*Macaca sinica*), purple-faced langur (*Trachypithecus vetulus*), slender lorises (*Loris tardigradus* and *Loris lydekkerianus*) and the gray langur (*Semnopithecus priam thersites*). The primates of Sri Lanka are endemic and considered to be Critically Endangered or Endangered. Here we report on some observations, information from interviews with local people regarding primate-human interactions, and also morphological differences in the subspecies we observed during field visits in 2004, 2005, and 2007. When asked, most people stated that they believed that primate populations had increased over the years, and many consider them to be agricultural pests due to the damage they inflict on crops. Due to religious beliefs, hunting and killing of primates were reported in low frequency, but some eat the meat of purple-faced langur for medicinal purposes. The most common methods people use to prevent monkeys from damaging crops are throwing stones, and the use of firecrackers or any other way of producing loud noises. The major threat that these primates face is the destruction of their habitat due to deforestation, human population growth and the expansion of various rural development projects. Public awareness programs for schools and through the media are needed to encourage the protection of these animals and their habitats.

Key words: Toque macaque, purple-faced leaf langur, gray langur, human-primate conflict

Introduction

Sri Lanka is an island, situated between 79°39' and 81°53'E, and 05°54' and 09°52'N, in the Indian Ocean, off the southeastern tip of India. Although small (65,000 km²), Sri Lanka has many endemic species in its fauna and flora (Gunethilleke and Gunethilleke 1983; Erdelen 1988), including five primate species (Table 1) all of which are threatened (Dela 2007; Rudran 2007). The western purple-faced langur (*Trachypithecus vetulus nestor*) and the Horton Plains slender loris (*Loris tardigradus nycticeboides*) are Critically Endangered (IUCN 2008) and have been listed amongst the World's 25 Most Endangered Primates (Dela and Rowe 2006; Nekaris 2006). The toque macaques (three subspecies), the gray-handed crested langur (*Semnopithecus priam thersites*), the remaining three subspecies of the purple-faced langur, and the Sri Lankan subspecies of the lorises, *L. tardigradus* and *L. lydekkerianus*, are all Endangered (IUCN 2008).

The forest cover of Sri Lanka has been declining at a steady rate over the past few decades due to agricultural and irrigational developmental projects and to human settlements

(Erdelen 1988; Wickramagamage 1998). Recent developments in the industrial and agricultural sectors and the growing human population have been particularly damaging to the country's remaining forests, and a greater awareness of their plight and measures for their protection are paramount for the survival of these primates. The total closed-canopy forest cover decreased from about 84% of the total land area in 1881 to about 30% in 2005. Rudran (2007) estimated that 81% of the habitat of the Critically Endangered western purple-faced langur is in deforested areas with dense human populations, and only two natural forest patches of about 21 km² remaining around two reservoirs. Owing to the expansion of the plantation industry, forest cover in the hill country has been reduced to isolated patches on hill tops and a handful of reserves above the 1,524 m contour (Wickramagamage 1998), posing a serious threat for the survival of the Endangered slender loris, *L. t. nycticeboides* (Mittermeier *et al.* 2007).

Most of the forests in wet zone and dry zone areas have been fragmented. As a result, primates tend to frequent farms and agricultural plots in search of food. This inevitably creates conflict, as has been reported in many parts of Sri Lanka

Table 1. Primate species and subspecies in Sri Lanka.

	Common name ¹	Reference
<i>Macaca sinica sinica</i> (Linnaeus, 1771)	Common toque macaque	Dittus (1975)
<i>Macaca sinica aurifrons</i> Pocock, 1931	Pale-fronted toque macaque	Dittus (1975)
<i>Macaca sinica opisthomelas</i> Hill, 1942	Hill zone toque macaque	Dittus (1975)
<i>Trachypithecus vetulus vetulus</i> (Erxleben, 1777) ²	Southern purple-faced langur	Molur <i>et al.</i> (2003)
<i>Trachypithecus vetulus monticola</i> (Kelaart, 1850) ²	Highland purple-faced langur	Molur <i>et al.</i> (2003)
<i>Trachypithecus vetulus philbricki</i> (Phillips, 1927) ²	Northern purple-faced langur	Molur <i>et al.</i> (2003)
<i>Trachypithecus vetulus nestor</i> Bennett, 1833 ²	Western purple-faced langur	Molur <i>et al.</i> (2003)
<i>Semnopithecus priam thersites</i> (Blyth, 1847) ³	Grey-handed crested langur	Molur <i>et al.</i> (2003)
<i>Loris tardigradus tardigradus</i> (Linnaeus, 1758)	Red slender loris	Nekaris & Jayawardene (2003, 2004)
<i>Loris tardigradus nycticeboides</i> Hill, 1942	Horton Plains slender loris	Nekaris & Jayawardene (2003, 2004)
<i>Loris lydekkerianus grandis</i> Hill & Phillips, 1932	Highland slender loris	Nekaris & Jayawardene (2003, 2004)
<i>Loris lydekkerianus nordicus</i> Hill, 1933	Northern Ceylonese slender loris	Nekaris & Jayawardene (2003, 2004)

¹Common names follow Brandon-Jones *et al.* (2004).

²Brandon-Jones *et al.* (2004) and Dela (2007) consider that the purple-faced langur is a member of the genus *Semnopithecus*. Groves (2001, 2005), Molur *et al.* (2003), and Rudran (2007), on the other hand, place it in the genus *Trachypithecus*.

³Groves (2005) considers *Semnopithecus priam thersites* to be a junior synonym of *S. p. priam* (Blyth, 1844).

(McDougal 1987; Sukumar 1989; Nowell and Jackson 1996; Katugaha *et al.* 1999; Santiapillai and Jayawardene 2004). Many of the Sri Lankan primates are found near Buddhist and Hindu temples. The priests are more tolerant, and the constant supply of food received from large numbers of pilgrims (for example, Kataragama, Sellakataragama and Vadasitikanda) keep them around the temple grounds. Primates are otherwise generally restricted to certain National Parks, sanctuaries and remaining forest patches, and only the toque macaque is widespread.

In order to implement effective conservation strategies, it is important to know the current distributions of these primates and their remaining forest habitats in each province and district, as well as the threats they face. Our research aims to document the primate species, their group numbers and sizes, in the national parks and temples. Important too is an understanding of the perceptions that the local communities have of the primates, in order to better assess the context for conservation initiatives.

Methods

To date we have carried out three field trips in Sri Lanka; 2004, 2005 and 2007. Even though subspecies differences among these primates have been clearly described (Dittus 1975; Molur *et al.* 2003), there are no published photographs which effectively illustrate them. We collected photographic records of the toque macaques, purple-faced langurs and gray langurs for this purpose.

In 2004, we spent one week in the Anuradhapura and the Kandy-Udawattakele forest area; in 2005, two weeks traveling to Sinharaja and Polonnaruwa; and in 2007, the Wildlife Department provided permits to survey the primate populations of Bundala, Yala, Udawalawe and Horton Plains national parks. Our surveys, 5–20 February (16 days), consisted of walking the existing trails, accompanied by a guide, and recording all groups seen and their sizes. On our way to

these national parks we also collected information on primates in Rumassala, around Kataragama, Badulla, Dambulla and Kandy (for further details of places visited see Table 2). We have also observed primates living in the suburbs of Colombo (Wijerama—around the University of Sri Jayawardenepura, Boralasgamuwa, Navinna).

In 2007, we interviewed people during the field trip using a questionnaire, accompanied by images of each primate species with their common names in Sinhalese, Tamil and English. The questionnaire included 28 questions on such topics as the primates that could be seen in the area, the approximate number of groups and their group size, whether they damage crops, whether measures are taken to prevent crop damage, about the extent of hunting and eating primates, land use, and about peoples' opinions of primates.

Results

External morphological subspecies traits

Characteristic external morphological differences of some of the different subspecies observed are shown in Figures 1–3 (see Table 1 for subspecies listings). The toque macaque of the wet zone (*Macaca sinica aurifrons*) has the darkest pelage color of the three subspecies. The dry zone subspecies (*M. s. sinica*) has the lightest pelage and is the smallest. It has the shortest crown hair length, while the highland subspecies (*M. s. opisthomelas*) has the longest, and *M. s. aurifrons* is intermediate (Fig. 1).

Among the four purple-faced langurs (*Trachypithecus vetulus*), the southern subspecies (*T. v. vetulus*) has the darkest pelage color and their white rump patch is more apparent than in the western (*T. v. nestor*) and northern (*T. v. philbricki*) subspecies. The montane *T. v. monticola* lacks a rump patch, is the largest of the four, and has the longest cheek hairs (Fig. 2). During our surveys, we found that gray langurs (*Semnopithecus priam*) in the southern dry zone are generally lighter in pelage color than in the north (Fig. 3).




Figure 1. The three macaque subspecies found in Sri Lanka; **a.** *Macaca sinica sinica*–(Sellakataragama); **b.** *Macaca sinica aurifrons*–(Kandy: Peradeniya Gardens) and **c.** *Macaca sinica opisthomelas*–(Ohiya). Photographs by Charmalie Nahallage.




Figure 2. The four purple-faced langur subspecies found in Sri Lanka; **a.** *Trachypithecus vetulus nestor*–(Wijerama: Colombo); **b.** *Trachypithecus vetulus monticola*–(Pattipola); **c.** *Trachypithecus vetulus vetulus*–(**c1.** Rumas-sala, **c2.** Sinharaja); and **d.** *Trachypithecus vetulus philbricki*–(Polonnaruwa). Photographs by Charmalie Nahallage.




Figure 3. The gray langur *Semnopithecus priam thersites* found in Sri Lanka; **a.** north central dry zone (Anuradhapura) (photograph by Michael Huffman); and **b.** southern dry zone (Bundala National Park) (photograph by Charmalie Nahallage).

Distributions of primate species in National Parks and surrounding areas

We observed only two primate species in the three southern national parks of Bundala, Yala and Udawalawe: the toque macaque and the gray langur. The gray langur was the most commonly seen. We saw more groups of gray langurs than macaques, and they were larger (Table 2). The gray langurs in the parks were more habituated to humans and easier to observe. The situation was similar outside the parks. In the Kataragama area (Southern province), grey langurs were seen mostly in Buddhist and Hindu temples, where they were partially provisioned by devotees and priests, and were quite tame. Few macaque groups were observed around these temples, and their groups were smaller (Table 2).

In Horton Plains National Park we recorded four purple-faced langur groups and one macaque group. The langur groups were relatively small and very shy. The macaque group we saw was quite tame, commonly begging for food from people along the road. Outside the park, in the areas of Ohiya and Pattipola, we observed one macaque group and one langur troop (Table 2). The langurs were again quite shy compared to the macaques.

The macaque groups we encountered in the Kandy-Peradeniya Botanical Gardens and in Dambulla were small, except for one we saw in Udawattakale (Table 2). They were quite habituated and commonly begged or stole food from local residents and tourists.

Results of the Questionnaires

We interviewed 39 people in 2007: 14 women and 25 men, 20 to 85 years old. Twelve were small-scale fruit or vegetable vendors, whose livelihoods were affected by primate activities. Housewives and government officers each accounted for six. Wildlife officers and guides working in the park accounted for four each, retired government officers, teachers, traditional doctors, a justice of the peace, fisherman,

Table 2. Number of places visited and primate species observed.

Province District	Location	Species	<i>T. vetulus</i> No. of groups (size)	<i>S. priam</i> No. of groups (size)	<i>M. sinica</i> No. of groups (size)
Western					
Colombo	Wijerama	<i>T. v. nestor</i>	3 (6±1.5)	—	—
	Navinna		1 (5)		
	Boralasgamuwa		1 (7)		
Sabaragamuwa					
Ratnapura	Sinharaja Forest	<i>T. v. vetulus</i>	1 (8)	—	1(5)
		<i>M. s. aurifrons</i>			
Ratnapura	Udawalawa National Park	<i>S. p. thersites</i>		3 (22±27)	1 (20–25)
		<i>M. s. sinica</i>			
Southern					
Galle	Rumassala	<i>T. v. vetulus</i>	2 (8±0.7)		
		<i>M. s. aurifrons</i>			
Hambantota	Bundala National Park	<i>S. p. thersites</i>		3 (22±17)	2 (13±10)
		<i>M. s. sinica</i>			
Hambantota	Vadasitikanda	<i>S. p. thersites</i>		3 (8±6)	2 (8±3)
	Kataragama	<i>M. s. sinica</i>		3 (27±12)	1 (9)
	Sellakataragama			2 (5±0.7)	1 (8)
	Katagamuwa			1 (7)	1 (10)
	Sithulpahuwa			1 (31)	1 (26)
	Yala National Park			7 (13±9)	
Uva					
Moneragala	Wellawaya	<i>M. s. aurifrons</i>			1 (6)
Badulla	Rawana Ella	<i>T. v. monticola</i>			1 (15)
	Ohiya	<i>M. s. opisthomelas</i>	1 (20)		1 (10)
Central					
Nuwaraeliya	Horton Plains National Park	<i>T. v. monticola</i>	4 (7±3)		1 (15)
	Pattipola	<i>M. s. opisthomelas</i>	1(6)		
Kandy	Peradeniya Gardens	<i>M. s. aurifrons</i>			
	Udawattakele				1*
	Victoria Reservoir				1*
Matale	Dambulla	<i>M. s. sinica</i>			3 (8±5)
North Central					
Anuradhapura	Anuradhapura	<i>M. s. sinica</i>		*	*
Polonnaruwa	Polonnaruwa	<i>M. s. sinica</i>	*	*	*
North Western					
Kurunegala	Kurunegala	<i>M. s. sinica?</i>			*

* Observed but could not determine the number of groups nor their size.

priests and a postmaster accounted for one each. In 2007, we spent the most time in the Southern and Uva provinces, so the results of the questionnaire strongly reflect conditions there rather than in the other provinces we visited. Sixty-one percent of the questionnaires were from the Southern Province, 21% from Uva Province, 10% from the Central Province and 8% were from the Western Province.

When asked about primate numbers, 82% of the people believed that numbers had increased over the years, and 95% informed us that the primates raid crops. All primates except for the lorises were considered pests in all the provinces we visited. Of the people we interviewed, 67% considered them to be pests, 13% did not, and 20% failed to comment. Fifty-one percent said that people do not hunt or kill monkeys, 38% said that some people in their area were known to kill monkeys (all from the Southern and Uva provinces), and 10% made no comment or were unsure.

The most common cause of primate deaths was reported as predation by dogs (26%), leopards (18%), crocodiles (8%) and pythons (8%). Twenty-one percent of the interviewees indicated that there were no natural predators of monkeys in their areas. Other factors reported to be responsible for deaths were electrocution from power lines (13%), hunting (8%) and road kill (3%).

People use a number of methods to prevent monkeys from raiding their crops and gardens. The most common was to use firecrackers (37%) to chase them away. Other frequently used methods were throwing stones (16%) or making loud noises (8%). Others reported using catapults/sling shots (5%) and dogs (3%). Less frequently used were electric barriers, covering fruit trees with nets, hanging red flags or mirrors on crop plants or in the vicinity, and scarecrows. Mostly, these methods were effective and harmless. Only in a few areas did people shoot them (6%) or use traps (2%).

Discussion

The majority of people we interviewed told us that monkey populations and the incidence of crop-raiding had increased over the years. Macaques (*Macaca*), baboons (*Papio*) and guenons (*Cercopithecus*) are the principal crop-raiding monkeys in Asia and Africa (Else 1991; Hill 1997; Naughton-Treves 2001; Osborn and Hill 2005; Riley 2007). This is partly due to their complex social organization, adaptable and intimidating behavior, ability to travel on the ground and in the trees, and their reliance on unspecialized and omnivorous diets (Frothman-Quick 1986; Else 1991; Hill 2000; Webber *et al.* 2007). We have no information on primate population sizes in the past, and so it is difficult to determine whether the perceived increase in numbers is real or due to changes in behavior or forest loss (increased, and forced, proximity to humans). We believe that the last of the three possibilities is the most likely.

Threats to each of the primate species differ in different parts of the country and depend in part on the presence of other primates and the socio-economic status of the area. For example, the purple-faced langurs are the most common primate in the Western Province, and there they are considered pests, damaging houses and raiding garden crops (Dela 2007; Rudran 2007). In Nuweraeliya district (Central Province) they co-exist with macaques, which people consider to be the more troublesome, being more present in human settlements, and stealing food from houses and raiding crops more often than langurs, which are more shy and tend to avoid human habitations. The more terrestrial and omnivorous lifestyle of macaques, compared to the arboreal leaf-eating langurs, brings them closer to humans, and they are considered as pests in many areas. In a similar study around Kibale National Park in Uganda, Hill (2000) found that red-tailed guenons (*Cercopithecus ascanius*) and L'Hoest's guenons (*C. lhoesti*) were more frequent visitors to farms than olive baboons (*Papio anubis*), but the baboons caused more damage to the crops. In the Western Province, langurs can be mostly found in gardens and they eat fruits grown for household consumption but not crops grown for commercial use. In these parts, primates were considered to be more of a nuisance than crop raiding pests, and people were quite tolerant of them. In the Southern and Central provinces, however, macaques and langurs raid commercial crops with more serious consequences. People have a harsher opinion of them as pests. Macaques are considered pests in the districts of Anuradhapura, Polonnaruwa, Kandy and Matale. Gray langurs were considered to be more serious pests than macaques in the areas of Bundala, Kataragama, Sellakataragama, Vadesitikanda and Sithulpahuwa due to their large group sizes and their habit of stealing food from shops as well as damaging crops. More systematic long-term data collection is planned to verify these trends.

In Sri Lanka, people are generally tolerant toward animals due to religious beliefs, and the major threat to their survival is more likely to be the loss of their forest habitats. The few reports we received of killings were all secret

operations and on a small-scale since primates are protected by law. Hunting primates for food was not a common practice in most of these areas since most of the people are Buddhists and against the killing of any animal. In some areas, however, especially in the Southern Province, people believe that monkey meat can cure certain illnesses and some use especially the purple-faced langur for medicinal purposes. In the Southern Province it is a common belief that the meat of the purple-faced langur is good for asthma or that it can cure sight defects. Of course, none of this has been medically proven. Another interesting belief of people in the south is that monkey organs (heart and lungs) are being used in the cities for organ transplants in humans due to their close similarity to humans. Some believe that the monkey's right leg contains human flesh, and even if they eat the meat they usually avoid eating this part. When we inquired about the macaques, many reported that they would not eat them because they are smaller and have too little meat—only the meat of the larger langurs was eaten. Consumption of monkey meat can be fatal if it is not prepared properly. Lamabadusuriya (1992) reported an outbreak of salmonella following consumption of monkey meat in the Southern Province (the species eaten was not reported). The author believed that the meat was probably contaminated because the monkey was already dead when the people found it.

Although most farmers believe that primates cause more damage to their crops than other species, some researchers have systematically quantified the crop damage caused by primates and other animals and shown that the damage is far less than the farmers believe (Siex and Struhsaker 1999; Riley 2007). The larger size and large group sizes of primates can give people the wrong impression as to the extent of crop damage. In Sri Lanka too, it is necessary to systematically quantify the damage caused by primates and inform farmers of these results. Together with their help, it will be possible to implement methods to control crop damage, benefiting both primates and farmers alike. Public awareness programs for schools and through the media are needed to encourage the protection of these animals and their habitats.

Acknowledgments

Nahallage and Huffman thank the HOPE Project, a Core-to-core program sponsored by the Japan Society for the Promotion of Science (JSPS), for travel funds to carry out the 2006 and 2007 field visits. We also thank the Wildlife Department of Sri Lanka for giving permission to conduct research in Bundala, Yala, Udawalawa and Horton Plains national parks. Thanks are due to Prof. Singha Kumara, Department of Forestry and Environmental Science, University of Sri Jayawardenepura, for arranging the logistics of our field trip to his forest ecology study site in Sinharaja in 2006. We are also grateful to Nelum Yapa and Sisira Siriwardene for their participation and help in our trip to Sinharaja. Our sincere thanks to Wolfgang Dittus for his invitation and hospitality during our stay at his field site in Polonnaruwa, and to his

field assistants for their help. Thanks are also due to all the people we met in the national parks during our field visits.

Literature Cited

- Brandon-Jones, D., A. A. Eudey, T. Geissmann, C. P. Groves, D. J. Melnick, J. C. Morales, M. Shekelle and C.-B. Stewart. 2004. Asian primate classification. *Int. J. Primatol.* 25(1): 97–164.
- Dela, J. 2007. Seasonal food use strategies of *Semnopithecus vetulus nestor*, at Panadura and Piliyandala, Sri Lanka. *Int. J. Primatol.* 28: 607–626.
- Dela, J. and N. Rowe. 2006. Western purple faced langur *Semnopithecus vetulus nestor* Bennett, 1833. In: R. A. Mittermeier, C. Valladares-Padua, A. B. Rylands, A. A. Eudey, T. M. Butynski, J. U. Ganzhorn, R. Kormos, J. M. Aguiar and S. Walker. 2006. Primates in peril: The world's 25 most endangered primates, 2004–2006, pp.12–13, 24. *Primate Conserv.* (20): 1–28.
- Dittus, W. P. J. 1975. Population dynamics of the toque monkey, *Macaca sinica*. In: *Socioecology and Psychology of Primates*, R. H. Tuttle (ed.), pp.125–151. Mouton Publishers, The Hague.
- Erdelen, W. 1988. Forest ecosystems and nature conservation in Sri Lanka. *Biol. Conserv.* 43: 115–135.
- Else, J. G. 1991. Non-human primates as pests. In: *Primate Responses to Environmental Change*, H. O. Box (ed.), pp.155–165. Chapman and Hall, London.
- Frothman-Quick, D. L. 1986. Activity budgets and the consumption of human foods in two troops of baboons (*Papio anubis*) at Gilgil Kenya. In: *Primate Ecology and Conservation*, J. G. Else and P. C. Lee (eds.), pp.221–228. Cambridge University Press, Cambridge.
- Groves, C. P. 2001. *Primate Taxonomy*. Smithsonian Institution Press, Washington, DC.
- Groves, C. P. 2005. Order Primates. In: *Mammal Species of the World: A Taxonomic and Geographic Reference*, 3rd Edition, Volume 1, D. E. Wilson and D. M. Reeder (eds.), pp.111–184. Johns Hopkins University Press, Baltimore.
- Gunathilleke, C. V. S. and I. A. U. N. Gunathilleke. 1983. A forestry case study of the Sinharaja rainforest in Sri Lanka. In: *Forest and Freshwater Development and Conservation in Asia and the Pacific*, L. S. Hamilton (ed.), pp.289–357. Westview Press, Colorado.
- Hill, C. M. 2001. Conflict of interest between people and baboons: Crop raiding in Uganda. *Int. J. Primatol.* 21: 299–315.
- Hill, C. M. 1997. Crop-raiding by wild vertebrates: the farmer's perspective in an agricultural community in western Uganda. *Int. J. Pest Mgmt.* 43: 77–84.
- IUCN. 2008. *2008 IUCN Red List of Threatened Species*. International Union for Conservation of Nature and Natural Resources (IUCN), Species Survival Commission (SSC), Gland, Switzerland, and Cambridge, UK. Available at: <<http://www.iucnredlist.org>>. Accessed on 15 October 2008.
- Katugaha, H. I. E., M. de Silva and C. Santiapillai. 1999. A long-term study on the dynamics of the elephant (*Elephas maximus*) population in Ruhuna National Park, Sri Lanka. *Biol. Conserv.* 89: 51–59.
- Lamabadusuriya, S. P., C. Perera, I. V. Devasiri, U. K. Jayantha and N. Chandrasiri. 1992. An outbreak of salmonellosis following consumption of monkey meat. *J. Trop. Med. Hyg.* 95: 292–295.
- McDougal, C. 1987. The man-eating tiger in geographic and historical perspective. In: *Tigers of the World: The Biology, Biopolitics, Management and Conservation of Endangered Species*, R. L. Tilson and U. S. Seal (eds.), pp.435–448. Noyes Publications, Park Ridge, NJ.
- Mittermeier, R. A., J. Ratsimbazafy, A. B. Rylands, L. Williamson, J. F. Oates, D. Mborá, J. U. Ganzhorn, E. Rodríguez-Luna, E. Palacios, E. W. Heymann, M. C. M. Kierulff, L. Yongcheng, J. Supriatna, C. Roos, S. Walker and J. M. Aguiar. 2007. Primates in Peril: The World's 25 Most Endangered Primates, 2006–2008. *Primate Conserv.* (22): 1–40.
- Molur, S., D. Brandon-Jones, W. Dittus, A. Eudey, A. Kumar, M. Singh, M. M. Feeroz, M. Chalise, P. Priya and S. Walker. 2003. *Status of South Asian Primates: Conservation Assessment and Management Plan (C.A.M.P.) Workshop Report, 2003*. Zoo Outreach Organization (ZOO) and CBSG–South Asia, Coimbatore, India.
- Naughton-Treves, L. 2001. Farmers, wildlife and the forest fringe. In: *African Rain Forest Ecology and Conservation: An Interdisciplinary Perspective*, A. D. Weber, L. J. T. White, A. Vedder and L. Naughton-Treves (eds.), pp.369–384. Yale University Press, New Haven.
- Nekaris, K. A. I. 2006. Horton Plains slender loris, Ceylon mountain slender loris *Loris tardigradus nycticeboides* (Hill, 1942). In: R. A. Mittermeier, C. Valladares-Padua, A. B. Rylands, A. A. Eudey, T. M. Butynski, J. U. Ganzhorn, R. Kormos, J. M. Aguiar and S. Walker. 2006. Primates in Peril: The World's 25 Most Endangered Primates, 2004–2006, pp.10–11, 23. *Primate Conserv.* (20): 1–28.
- Nekaris, K. A. I. and J. Jayawardene. 2003. Pilot study and conservation status of the slender loris (*Loris tardigradus* and *L. lydekkerianus*) in Sri Lanka. *Primate Conserv.* (19): 83–90.
- Nekaris, K. A. I. and J. Jayawardene. 2004. Survey of the slender loris (Primates, Lorisidae Gray, 1821: *Loris tardigradus* Linnaeus, 1758 and *Loris lydekkerianus* Cabrera, 1908) in Sri Lanka. *J. Zool. Soc., Lond.* 262: 327–338.
- Nowell, K. and P. Jackson (eds.). 1996. *Wild Cats: Status Survey and Action Plan*. IUCN, Gland, Switzerland.
- Osborn, F. V. and C. M. Hill. 2005. Techniques to reduce crop loss: Human and technical dimensions in Africa. In: *People and Wildlife: Conflict or Coexistence?* R. Woodroffe, S. Thirgood and A. Rabinowitz (eds.), pp.72–85. Cambridge University Press, Cambridge, UK.
- Patterson, J. D. and J. Wallis. 2005. *Commensalism and Conflict: The Primate-Human Interface*. American Society of Primatologists, Norman, Oklahoma.

- Riley, E. P. 2007. The human-macaque interface: Conservation implications of current and future overlap and conflict in Lindu National Park, Sulawesi, Indonesia. *Am. Anthropol.* 109: 473–484.
- Rudran, R. 2007. A survey of Sri Lanka's endangered and endemic western purple-faced langur (*Trachypithecus vetulus nestor*). *Primate Conserv.* (22): 139–144.
- Santiapillai, C. and R. Jayawardene. 2004. Conservation of the leopard and other carnivores in Sri Lanka. *Current Sci.* 86: 1063–1064.
- Siex, K. S. and T. T. Struhsaker. 1999. Colobus monkeys and coconuts: A study of perceived human-wildlife conflicts. *J. Appl. Ecol.* 36: 1009–1020.
- Sukumar, R. 1989. *The Asian Elephant: Ecology and Management*. Cambridge University Press, UK.
- Webber, A. D., C. M. Hill and V. Reynolds. 2007. Assessing the failure of a community-based human-wildlife conflict mitigation project in Budongo Forest Reserve, Uganda. *Oryx* 41: 177–184.
- Wickramagamage, P. 1998. Large-scale deforestation for plantation agriculture in the Hill country of Sri Lanka and its impacts. *Hydrol. Process* 12: 2015–2028.

Authors' addresses:

Charmalie A. D. Nahallage, Department of Sociology and Anthropology, University of Sri Jayawardenepura, Colombo, Sri Lanka. *Address for correspondence:* Department of Sociology and Anthropology, University of Sri Jayawardenepura, Gangodawila, Nugegoda, Sri Lanka, e-mail: <charmalie2@hotmail.com>.

Michael A. Huffman, Department of Social Systems Evolution, Primate Research Institute, Kyoto University, Japan.

Nadeera Kuruppu, Department of Wildlife Conservation, Colombo, Sri Lanka.

Tharindi Weerasingha, Department of Social Welfare, Divisional Secretariat Department, Matara, Sri Lanka.

Received for publication: October 2007

Revised: June 2008