



Expression and Function of Xmsx-2B in Dorso-Ventral Axis Formation in Gastrula Embryos

Authors: Onitsuka, Izumi, Takeda, Masatoshi, and Maéno, Mitsugu

Source: Zoological Science, 17(8) : 1107-1113

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.17.1107>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Expression and Function of *Xmsx-2B* in Dorso-Ventral Axis Formation in Gastrula Embryos

Izumi Onitsuka, Masatoshi Takeda and Mitsugu Maéno*

Department of Biology, Niigata University, Niigata 950-2181, Japan

ABSTRACT—*Msx* is a homeodomain-containing transcriptional factor that plays an essential role in pattern formation in vertebrata and invertebrata embryos. In *Xenopus laevis*, two *msx* genes have been identified (*Xmsx-1* and *Xmsx-2*). In the present study, we attempted to elucidate the expression and function of *Xmsx-2B* (formerly designated as *Xhox7.1'*) in early embryogenesis. Whole mount *in situ* hybridization analyses showed that the expression pattern of *Xmsx-2B* at gastrula and neurula stages was very similar to that of *Xmsx-1*: the transcript of *Xmsx-2B* was observed in ventral and lateral sides of the embryo. At the tailbud stage, however, the expression pattern of *Xmsx-2B* in neural tissues was distinct from that of *Xmsx-1*. An RNA injection experiment revealed that, like *BMP-4*, *Xmsx-2B* has a strong ventralizing activity. In the *Xmsx-2B*-injected embryos, differentiation of axial structures such as the notochord, muscle, and neural tissue was completely suppressed, whereas α -globin mRNA, a blood cell marker, was highly expressed. Simultaneous injection of *Xmsx-1* and *Xmsx-2B* RNAs showed that they function in an additive manner. It was also shown that coinjection of *Xmsx-2B* with a dominant-negative *BMP-4 receptor* (*tBR*), which can induce formation of secondary axis when injected alone in ventral blastomeres, suppressed secondary axis formation. Furthermore, *Xmsx-2B* also suppressed secondary axis formation, which was induced by a dominant-negative form of *Xmsx-1* (*VP16/msx-1*). Therefore, like *Xmsx-1*, *Xmsx-2B* is a downstream nuclear factor of the BMP-4-derived ventralizing signal, and these two factors probably share the same target molecules. In conclusion, *Xmsx-1* and *Xmsx-2B* function in dorso-ventral axis formation in early *Xenopus laevis* development.

INTRODUCTION

A number of studies have indicated that BMP-4 (bone morphogenetic protein-4), a member of the TGF- β superfamily, is a key regulator in body planning, especially in dorso-ventral patterning of mesodermal tissues and in neural tissue differentiation, in the *Xenopus laevis* embryo (Dale *et al.*, 1992; Jones *et al.*, 1992; Fainsod *et al.*, 1994; Graff *et al.*, 1994; Maéno *et al.*, 1994; Suzuki *et al.*, 1994; Schmidt *et al.*, 1995). Recently, homeodomain-containing transcriptional factors that mediate the BMP-4 signal have been isolated and analyzed: *Xmsx-1* (Maeda *et al.*, 1997; Suzuki *et al.*, 1997), *Xvent-1* (PV. 1) (Gawantka *et al.*, 1995; Ault *et al.*, 1996), *Xvent-2* (*Xbr-1*, *Vox*, *Xom*) (Onichtchouk *et al.*, 1996; Papalopulu and Kintner 1996; Ladher *et al.*, 1996) and *Xvex-1* (Shapira *et al.*, 1999). We and other investigators have revealed that *Xmsx-1* is expressed in the prospective ventral ectoderm and mesoderm at mid to late gastrula stages and that it has a ventralizing activity if its RNA is injected into the prospective dorsal region (Maeda *et al.*, 1997; Suzuki *et al.*, 1997). In ectodermal cells, *Xmsx-1* functions as an inhibitor of neurogenesis, and it has

been shown that *Xmsx-1* is a direct target gene of the BMP-4 signal (Suzuki *et al.*, 1997). More recently, we have demonstrated that *Xmsx-1* is an upstream factor of *Xwnt-8* in the ventralizing signal cascade and a possible direct repressor of organizer genes such as *gooseoid* (Takeda *et al.*, 2000).

In comparison to the numerous studies on *Xmsx-1*, quite a few studies have been carried out on the function of *Xmsx-2* (Su *et al.*, 1991; Gong and Kiba, 1999). In other species, *msx-2* has been studied extensively in various developing tissues such as bone, tooth, and limb bud (Monahan *et al.*, 1991; Coelho *et al.*, 1991; Mackenzie *et al.*, 1992; Jowett *et al.*, 1993). Most likely, the function of *msx-2* overlaps with that of *msx-1* (Semenza *et al.*, 1995), but in some cases *msx-2* may have distinct expression patterns and functions. For instance, expression domains of *msx-1* and *-2* overlap in some regions but are discrete along with anterior-posterior identity in chicken and mouse limb buds (Davidson *et al.*, 1991; Coelho *et al.*, 1991; Muneoka and Sassoon, 1992; Nohno *et al.*, 1992). The BMP-4/*msx-2* signal is involved in elimination of cranial neural crest cells in rhombomeres 3 and 5 (Graham *et al.*, 1993; Graham *et al.*, 1994; Takahashi *et al.*, 1998; Farlie *et al.*, 1999) and the interdigit regions of developing limbs (Zou and Niswander 1996; Ganam *et al.*, 1996; Ferrari *et al.*, 1998) by mediating apoptosis. Furthermore, the mechanisms of expressional control in mouse mammary development are dif-

* Corresponding author: Tel. +81-25-262-6183;
FAX. +81-25-262-6116.
E-mail maenobio@sc.niigata-u.ac.jp

ferent between *msx-1* and *-2* (Friedmann *et al.*, 1996; Phippard *et al.*, 1996).

These observations suggest that *msx-2* could have a distinct expression pattern and function in the early development of *Xenopus laevis*. In spite of the importance of the issue, expressional and functional natures of *msx-2* in *Xenopus* embryogenesis are poorly understood. *Xenopus msx-2* (*Xhox7.1'*) was first isolated by Su *et al.* (1991). In their report, however, major attention was paid to the expression of *Xmsx-1*. Afterward, no report has described the expression and function of *Xmsx-2* except for a very recent one in which a related but distinct gene from *Xhox7.1'* was described (Gong and Kiba, 1999). In this study, we attempted to elucidate the expression and the function of *Xhox7.1'*. We propose designating *Xhox7.1'* as *Xmsx-2B* in order to distinguish it from *Xmsx-2*, which was isolated by Gong and Kiba (1999). Our results showed that there is a strong correlation between the expression patterns of *Xmsx-1* and *Xmsx-2B* at gastrula to neurula stages and that those two genes play a role in the establishment of the dorso-ventral axis in developing *Xenopus* embryos.

MATERIALS AND METHODS

Cloning and cDNA construction

Xmsx-2B cDNA fragments of two different sizes were amplified by polymerase chain reaction (PCR) from *Xenopus* stage10 cDNA. A short fragment including 402 N-terminal base pairs was used as a probe for *in situ* hybridization, and another fragment including the whole protein-coding region (815 bp) was used for an RNA injection assay. The primer sequences for the PCR were as follows: 5'-TC T-GCG-CCC-CTT-TAC-TCA-CCG-CA-3' and 5'-TCA-AAG-TGC-AAG-AGG-ATG-GGC-TCA-3' for the 402-bp fragment; and 5'-TCT-GCG-CCC-CTT-TAC-TCA-CCG-CA-3' and 5'-CTA-TGT-CAT-GTC-ACC-CTC-CTC-AGA-TA-3' for the 815-bp fragment (Su *et al.*, 1991). Each product was ligated into pCRTM II (*In vitro* gene) for subcloning, and the DNA sequence of the inserts was determined using an Auto Read Sequencing kit (ALF DNA Sequencer, Pharmacia Biotech). The obtained sequence was completely identical to that of *Xhox7.1'*, which was previously reported (Su *et al.*, 1991). For injection of sense RNA of *Xmsx-2*, an 815-bp fragment including an open reading frame was excised with Not I, blunt ended, and ligated into a blunt-ended pCS2+ expression vector.

Embryos, mRNA micro-injection, and DAI judgement

Xenopus laevis embryos were obtained by artificial insemination after induction of females with 250 i.u. of human chorionic gonadotropin. Fertilized embryos were dejellied with 2.5% thioglycolic acid (pH8.3) and washed several times in 5% MMR. After the embryos were transferred into 3% Ficoll/Steinberg's solution, equatorial regions of either two ventral or dorsal blastomeres at the 4-cell stage were injected with mRNAs using a Nanoject microinjector (3.7-6 nl/embryo). pCS2+ plasmids containing *Xmsx-1* whole cDNA (1.8 kb) or the 815-bp *Xmsx-2B* fragment were linearized with Xho I, and capped mRNA was synthesized using SP6 polymerase (Ambion). *VP16/msx-1*, a dominant-negative form of *Xmsx-1*, has been described previously (Takeda *et al.*, 2000). Injected embryos were incubated in 50% Steinberg's solution at 23°C until the sibling embryos reached stage 33/34 for determining DAI (dorso-anterior index) as described by Kao and Elinson (1988). Developmental stages were designated according to Nieuwkoop and Faber (1967).

Whole-mount *in situ* hybridization

Whole-mount *in situ* hybridization was performed as described previously (Shain *et al.*, 1996). Digoxigenin-labeled antisense mRNA probes were synthesized from the following plasmids: *Xmsx-1*, an N-terminal fragment (EcoRI/HindIII, 440 bp) in pSP72; and *Xmsx-2B*, an N-terminal fragment (402 bp) in pCRTMII. The former plasmid was linearized with EcoRI, and RNA was synthesized by SP6 polymerase. The latter plasmid was linearized with BamHI, and RNA was synthesized by T7 polymerase. The positive signals were visualized using BM purple (Moors *et al.*, 1995). For neural marker probes, fragments of *slug* and *krox-20* were amplified by the PCR method from the cDNAs of stage18 whole embryo and of stage 27 head tissue, respectively. The primer sequences for the PCR were as follows: 5'-GAA-TCT-GGT-TGC-TGT-GTA-G-3' and 5'-CGT-TGC-TGG-ATT-GTC-TAG-GC-3' for *slug*; and 5'-CCA-ACC-GCC-CCA-GTA-AGA-CC-3' and 5'-GTG-TCA-GCC-TGT-CCT-GTT-AG-3' for *krox-20*. Both fragments were subcloned into a pCR2.1 vector (*In vitro* gene), and their DNA sequences were determined. For making an antisense RNA probe, the plasmids were cut with BamHI, and RNAs were synthesized by T7 polymerase.

RESULTS

Expression pattern of *Xmsx-2B* in the early embryo

To investigate the expression pattern of *Xmsx-2B* during embryogenesis and to compare it with that of *Xmsx-1*, whole-mount *in situ* hybridization analysis was performed using various stages of albino embryos. As shown in Fig.1, during gastrula and early neurula stages, the *Xmsx-2B* transcript was observed in the ventral and lateral marginal zones and animal pole area (Fig.1 B, D and F), as was also found in the case of *Xmsx-1* (Maeda *et al.*, 1997; Suzuki *et al.*, 1997; Fig.1A, C and E). At the neurula stage (stages 16–17), two parallel lines of transcript were observed at the dorsal side along the border between the prospective neural tissue and epidermis (arrowheads in Fig. 1G and H), and also strong spot signals were detected on the medial and lateral sides of the prospective neural crest (arrows in Fig. 1 G and H), with reference to the expression of *slug*, a neural crest marker (Mayor *et al.*, 1995; Fig. 1 J). We also compared the site of expression with that of *krox-20*, a marker of rhombomeres 3 and 5 (Bradley *et al.*, 1992; Fig.1 I and L), and we found that strong spot signals were located at the level of the hind brain (rhombomeres 3-5) (Fig. 1 H and K). At the early tailbud stage (stage 23), expression patterns became different between *Xmsx-1* and *Xmsx-2B* in the neural tissue (Fig.1 M, N, P and Q). *Xmsx-1* was highly expressed in the broad areas including fore-, mid- and hind-brains, according to the expression of *krox-20* (Fig.1 O and R), and spinal cord, while *Xmsx-2B* was expressed in the medial region of neural tissues. These observations suggest that *Xmsx-1* and *Xmsx-2B* have the same or similar functions in gastrula stages but have different functions after the neurula stage.

Xmsx-2B has a ventralizing activity

It has been shown in previous studies that *Xmsx-1* has ventralizing and anti-neuralizing activities in early embryogenesis (Maeda *et al.*, 1997; Suzuki *et al.*, 1997; Ishimura *et al.*, 2000; Takeda *et al.*, 2000). To examine whether *Xmsx-2B*








Fig. 1. Whole-mount *in situ* hybridization showing the expressions of *Xmsx-1* (A, C, E, G, M, P), *Xmsx-2B* (B, D, F, H, K, N, Q), *slug* (a neural crest marker) (J), and *krox-20* (a hindbrain marker) (I, L, O, R). At the mid gastrula stage (st.12), *Xmsx-1* and *Xmsx-2B* mRNAs are expressed in the ventral and lateral marginal zone (A–D) (A and B, vegetal view and dorsal side is toward the top; C and D, lateral view and dorsal side is to the right). At the late gastrula stage (st.13), transcripts are accumulated in the lateral regions adjacent to the presumptive neural fold (E, F). (E and F, dorsal view and anterior is toward the top). At the neurula stage (st.16–17), transcripts are detected along the border lines of prospective neural and non-neural tissues (arrowheads in G and H), and high levels of expressions are observed on the medial and lateral sides of the prospective neural crest cells (arrows in G and H), as shown by *slug* expression (J). These regions correspond to the level of the prospective hindbrain (rhombomeres 3–5), with reference to the expression of *krox-20* (I and L) (G–J, dorsal view and anterior is toward the top; K and L, anterior view and dorsal side is toward the top). At the tailbud stage (st.23), the expression patterns of *Xmsx-1* and *Xmsx-2B* in neural tissue become clearly distinct (M, N, P and Q). *Xmsx-1* is expressed in broad areas, including the fore-, mid- and hindbrains and the spinal cord, while *Xmsx-2B* is expressed in a narrower region of neural tissues. (M–O, dorsal view and anterior is toward the top; P–R, anterior view and dorsal side is toward the top). Bars in A, M and P indicate 0.5 mm.



has the same or similar activity, we performed mRNA injection experiments. Two dorsal blastomeres of 4-cell stage embryos were injected with various concentrations of mRNAs encoding for the full-length *Xmsx-2B* protein, and the injected embryos were incubated for 2 days until the sibling embryos reached the tailbud stage. The embryos injected with *Xmsx-2B* mRNA showed ventralized phenotypes and the average DAI was reduced in dose-dependent manner (Fig. 2 D, E, G, J and M). This activity of *Xmsx-2B* seemed very strong among the known ventralizing agents, and as little as 10 pg of mRNA was enough to exert a ventralizing effect (average DAI=4.3, n=154; data not shown). Injection of 1ng RNA led to a hyper-ventralized phenotype completely lacking the dorso-ventral axis (average DAI=0.3, n=98; Fig. 2 J, M). The phenotype induced by *Xmsx-2B* overexpression was very similar to that obtained by the injection of *BMP-4* mRNA (average DAI=0.1, n=92; Fig. 2 I, L). In these embryos, ectodermal, mesodermal and endodermal layers were clearly distinguished, but differentiation of dorsal tissues such as the notochord, neural tube or muscle was not observed (Fig.2 O and P). Furthermore, *in situ* hybridization analysis indicated that in these hyper-ventralized embryos, α -globin mRNA was strongly expressed in the anterior region (Fig. 2 I, J) but α -actin was not (Fig. 2 L, M).

Since the phenotypes of the embryos injected with *Xmsx-2B* and *BMP-4* mRNA were similar, we examined the blood cell-inducing ability of *Xmsx-2B* in the dorsal marginal zone (DMZ) explants. After injection of several concentrations of *Xmsx-2B* mRNAs (200–800 pg/embryo) into the marginal zone

Fig. 2. Ventralizing activities of *Xmsx-1* and *Xmsx-2B*. A-F. Typical ventralizing phenotypes obtained in the mRNA injection experiments. Two dorsal blastomeres of 4-cell stage embryos were injected with designated amounts of RNAs, and the embryos were allowed to develop until stage 33/34 to determine the DAI (dorso-anterior index; Kao and Elinson, 1988). Injection of 400 pg of *Xmsx-1* (B) and 25 pg of *Xmsx-2B* (E) led to a moderate ventralizing phenotype (average DAIs being 3.1 and 3.5, respectively). Injection of 800 pg of *Xmsx-1* (C) and 50 pg of *Xmsx-2B* (E), which are twice the amounts of B and E, resulted in a severely ventralized phenotype (average DAIs being 2.2 and 2.3, respectively). The embryos coinjected with 400 pg of *Xmsx-1* and 25 pg of *Xmsx-2B* (F) showed a similar degree of ventralization to that of C and E. G. Average DAI and SD of RNA-injected embryos are represented by a bar graph. The numbers of embryos injected in each groups were as follows: n=157 (uninjecte), n=185 (*Xmsx-1* 400 pg), n=77(*Xmsx-1* 800 pg), n=122 (*Xmsx-2B* 25 pg), n=152 (*Xmsx-2B* 50 pg), and n=140 (*Xmsx-1* 400 pg+ *Xmsx-2B* 25 pg). Note that both *Xmsx-1* and *Xmsx-2B* had ventralizing activities and that their effects were additive. H-P. Hyperventralized phenotype (DAI=0.3, n=98) induced by a high dose of *Xmsx-2B* RNA. Embryos were injected with 1ng of *Xmsx-2B* RNA (J, M) or 1ng of *BMP-4* RNA as a control (I, L), and they were allowed to develop until the sibling embryos reached st.33/34 (H, K). Whole-mount *in situ* hybridization was performed to show the expressions of α -globin (H, I and J) and α -actin (K, L and M). The expression patterns of these markers in the embryos, which had been injected with *BMP-4* and *Xmsx-2B*, were quite similar. N, O, and P represent histological views of embryos shown in H, I and J. Note that no axial structure was observed in *BMP-4* or *Xmsx-2B* mRNA-injected embryos, and, instead, α -globin-positive cells were dramatically increased in comparison with those in un.injected embryos. Bars in A, H and N indicate 1 mm, 1 mm, and 0.5 mm respectively.

Table 1. Incidence of secondary axis formation in injected embryos

mRNA	Number of injected embryos	Secondary axis (%)
<i>tBR</i>	127	56
<i>tBR+Xmsx-1</i>	112	0
<i>tBR+ Xmsx-2B</i>	117	0
<i>VP16/msx-1</i>	95	65
<i>VP16/msx-1+Xmsx-1</i>	81	0
<i>VP16/msx-1+ Xmsx-2B</i>	118	0

The experimental protocol is described in the legend of Fig. 3. Note that *Xmsx-1* and *Xmsx-2B* completely suppressed formation of the secondary axis induced by *tBR* (a dominant-negative form of the BMP-4 receptor) and *VP16/msx-1* (a dominant-negative form of *Xmsx-1*).




Fig. 3. A-D. Injection of *Xmsx-1* or *Xmsx-2B* mRNA suppressed the secondary axis formation induced by *tBR*, a dominant-negative form of the BMP-4 receptor. Two ventral blastomeres of 4-cell stage embryos were injected with *tBR* (500 pg) RNA (B), *tBR* (500 pg) with *Xmsx-1* (1000 pg) RNA (C), and *tBR* (500 pg) with *Xmsx-2B* (200 pg) RNA (D). The embryos were allowed to develop until st.33/34, and the existence of secondary axis was determined morphologically. E-H. Injection of *Xmsx-1* or *Xmsx-2B* mRNA suppressed the secondary axis formation induced by *VP16/msx-1*, a dominant-negative form of *Xmsx-1*. The embryos were injected with *VP16/msx-1* (600 pg) RNA (F), *VP16/msx-1* (600 pg) with *Xmsx-1* (200 pg) (G), or *VP16/msx-1* (600 pg) with *Xmsx-2B* (200 pg) (H). Bar in A indicates 1mm.

of dorsal blastomeres, DMZs were excised at stage 10.5 and cultured until stage 39. Northern blot analysis showed that α -globin mRNA was not induced in the explants even though the highest amount of mRNA was injected (data not shown). Thus, we concluded that among the target genes of the BMP-4 signal, *Xmsx-1* and *Xmsx-2B* are not sufficient for blood cell differentiation.

To investigate whether *Xmsx-1* and *Xmsx-2B* act coop-

eratively, these RNAs were injected alone or simultaneously into two dorsal blastomeres (Fig. 2 B-G). The average DAI obtained by 25 pg of *Xmsx-2B* mRNA was 3.1 ± 1.4 ($n=122$), whereas that of embryos that received 400 pg of *Xmsx-1* mRNA was 3.5 ± 1.1 ($n=185$). When these mRNAs were coinjected, the DAI fell to 2.3 ± 1.0 ($n=140$). Since similar values of DAI were obtained in the embryos injected with 50 pg *Xmsx-2B* (DAI= 2.3 ± 1.4 ; $n=152$) or 800 pg *Xmsx-1* (DAI= 2.2 ± 1.2 ; $n=77$), we concluded that *Xmsx-1* and *Xmsx-2B* function in ventralization in an additive manner.

Xmsx-2B mediates BMP-4 signaling

It has been shown that the expression of *Xmsx-1* is regulated by the BMP-4 signal (Maeda *et al.*, 1997; Suzuki *et al.*, 1997). Taken together with these expression patterns of *Xmsx-2B* in the early embryo (Fig. 1), it seems that *Xmsx-2B*, like *Xmsx-1*, is regulated by the BMP-4 signal. To confirm this hypothesis, we examined whether *Xmsx-2B* could disturb the formation of the secondary axis, which was induced by *tBR* mRNA injection. Two ventral blastomeres of 4-cell stage embryos were injected with 500 pg of *tBR* mRNA alone or coinjected with 1ng of *Xmsx-1* or 200 pg of *Xmsx-2B*. As shown in Table 1 and Fig. 3 B, 56 % ($n=127$) of *tBR*-injected embryos formed the secondary axis. In contrast, the formation of the secondary axis was completely suppressed if either *Xmsx-1* ($n=112$) or *Xmsx-2B* ($n=117$) was coinjected with *tBR* mRNA.

We also examined whether the ventralizing activity of *Xmsx-1* can be replaced by that of *Xmsx-2B*. For this purpose, we used a fusion construct of *Xmsx-1* and VP16, a dominant-negative form of *Xmsx-1*, which was shown to induce formation of the secondary axis (Takeda *et al.*, 2000). The embryos injected ventrally with *VP16/msx-1* mRNA formed an incomplete secondary axis without a head structure (65%; $n=95$, Fig. 3 F). Simultaneous injection either of *Xmsx-1* (Fig. 3 G; $n=81$) or *Xmsx-2B* (Fig. 3 H; $n=118$) completely suppressed the formation of the secondary axis. These results provide us with evidence that the activity of *Xmsx-1* can be replaced by that of *Xmsx-2B* and, therefore, these two transcriptional factors share the same target molecules in the ventralizing signal.

DISCUSSION

The expression of *Xmsx-2B* overlaps that of *Xmsx-1* in gastrula and neurula stages

Since Su *et al.* (1991) first reported the isolation of an *Xhox7.1'* (*Xmsx-2B*) cDNA clone in *Xenopus laevis*, there have been no reports describing the expression and function of *Xenopus msx-2*, despite the general view of importance in various vertebrate embryogenesis. Very recently, however, Gong and Kiba (1999) isolated a *msx-2* cDNA clone and emphasized its function in antero-posterior patterning of mesoderm formation. From the viewpoints of sequence similarity, expression pattern, and biological activity, *Xmsx-2B* and the cDNA isolated by Gong and Kiba (1999) are subtypes of *Xenopus msx-2*.

As expected, the expression pattern of *Xmsx-2B* was very similar to that of *Xmsx-1* (Maeda *et al.*, 1997; Suzuki *et al.*, 1997) during gastrula and neurula stages. Concomitant expressions of *msx-1* and *-2* have been reported in various tissues in mammalian species (Mackenzie *et al.*, 1992; Jowett *et al.*, 1993; Wang and Sassoon, 1995; Friedmann *et al.*, 1996; Phippard *et al.*, 1996) and in birds (Coelho *et al.*, 1991; Ganan *et al.*, 1996; Ganan *et al.*, 1998). At the gastrulation stage, the transcript was detected in the lateral and ventral marginal zone (Fig. 1), suggesting that *Xmsx-2B* and *Xmsx-1* have the same or similar functions at this stage. At the tailbud stage, however, the expression pattern in neural tissues was clearly distinct. We suggest that the mechanism of expressional control in the two *msx* genes is conserved at least in early embryogenesis.

Xmsx-2B acts as a ventralizing factor

As suggested by the expression pattern of *Xmsx-2B* in gastrulating embryos, the forced expression of *Xmsx-2B* mRNA in dorsal blastomeres induced a strong ventralized phenotype (Fig. 2 J, M), and α -globin mRNA was strongly expressed in the embryos (Fig. 2 J). By analyzing the ventralizing activity of *Xmsx-1*, we noticed that, unlike BMP-4, *Xmsx-1* alone lacks activity to upregulate the expression of ventral genes (Maeda *et al.*, 1997). On the other hand, *Xmsx-2B*-injected embryos completely lost the body axis, while the same dose of *Xmsx-1* had a weaker ventralizing effect. We predicted, therefore, that *Xmsx-2B* may be a cofactor of *Xmsx-1* in blood cell-inducing activity. However, Northern blot analysis revealed that *Xmsx-2B* could not induce α -globin expression in the dorsal marginal zone explants, while BMP-4 could. Although there was no difference in the translational activities of *Xmsx-1* and *Xmsx-2B* RNAs in the reticulocyte lysate (data not shown), the stabilities of proteins in the cells were not determined. As the *Xmsx-2B* cDNA used in this study lacks the 3' UTR, there is a possibility that this 3' UTR might regulate the stability of *Xmsx-2B* mRNA. From these observations, we concluded that even though *Xmsx-2B* has a strong activity, *Xmsx-1* and *Xmsx-2B* ventralize the embryo by suppressing the expression of organizer genes (Takeda *et al.*, 2000), and in the prospective ventral cells in which the BMP-4 signal is activated, another unknown factor is required to trigger the blood cell differentiation.

Xmsx-1 and Xmsx-2B function in the dorso-ventral patterning of mesoderm

In the present study, we found that *Xmsx-1* and *Xmsx-2B* function in the ventralizing cascade in an additive manner (Fig. 2 A-G). In addition, the ability of *Xmsx-2B* to suppress the secondary axis formation induced by *VP16/msx-1* RNA (a dominant negative form of *msx-1* protein) supports our conclusion that the ventralizing activity of *msx-1* can be replaced by that of *Xmsx-2B* and that their target gene(s) might be common. So far, the molecular nature of how these *msx* proteins interact with each other is unclear, but previous reports have suggested that *msx* and *dlx* proteins can form homo- and

heterodimers in various combinations (Zhang *et al.*, 1997). Since *msx* proteins are transcriptional repressors (Catron *et al.*, 1995) and *dlx* proteins are transcriptional activators (Zhang *et al.*, 1997), the biological activities in transcriptional regulation are mutually exclusive (Zhang *et al.*, 1997). Other candidates of *msx*'s partner are *Xvent-1* (PV. 1) (Gawantka *et al.*, 1995; Ault *et al.*, 1996) and *Xvent-2* (*Xbr-1*, *Vox*, *Xom*) (Onichtchouk *et al.*, 1996; Papalopulu and Kintner, 1996; Ladher *et al.*, 1996). These factors are also homeodomain-containing proteins and possessing ventralizing activity. The similarity of expression patterns and functions between *msx* and *vent* suggests that these proteins might interact with each other, directly or indirectly, in prospective ventral cells. To elucidate the molecular mechanisms of *msx*'s function in the ventralizing activity and the following specification of tissue distribution, isolation and analysis of the interacting molecules will be required in the future.

ACKNOWLEDGEMENTS

This work was partly supported by grants from The Saneyoshi Foundation, The Sumitomo Foundation and The Ministry of Education, Science and Culture of Japan (No. 09680721).

REFERENCES

- Ault KT, Dirksen ML, Jamrich M (1996) A novel homeobox gene *PV.1* mediates induction of ventral mesoderm in *Xenopus* embryos. *Proc Natl Acad Sci USA* 93: 6415–6420
- Bradley LC, Snape A, Bhatt S, Wilkinson DG (1992) The structure and expression of the *Xenopus Krox-20* gene: conserved and divergent of expression in rhombomeres and neural crest. *Mech Dev* 40: 73–84
- Catron KM, Zhang H, Marshall SC, Inostroza JA, Wilson JM, Abate C (1995) Transcriptional repression by *Msx-1* does not require homeodomain DNA-binding sites. *Mol Cell Biol* 15: 861–871
- Coelho CND, Sumoy L, Rodgers BJ, Davidson DR, Hill RE, Upholt WB, Koshier RA (1991) Expression of the chicken homeobox-containing gene *Ghox-8* during embryonic chick limb development. *Mech Dev* 34: 143–154
- Dale L, Howes G, Price BMJ, Smith JC (1992) Bone morphogenetic protein 4: a ventralizing factor in *Xenopus* development. *Development* 115: 573–585
- Davidson DR, Crawley A, Hill RE, Tickle C (1991) Position-dependent expression of two related homeobox genes in developing vertebrate limbs. *Nature* 352: 429–431
- Fainsod A, Steinbeisser H, De Robertis EM (1994) On the function of BMP-4 in patterning the marginal zone of the *Xenopus* embryo. *EMBO J* 13: 5015–5025
- Farlie PG, Kerr R, Thomas P, Symes T, Minichiello J, Hearn CJ, Newgreen D (1999) A paraxial excision zone creates patterned cranial neural crest cell outgrowth adjacent to rhombomeres 3 and 5. *Development* 126: 70–84
- Ferrari D, Lichtler AC, Pan ZZ, Dealy CN, Upholt WB, Koshier RA (1998) Ectopic expression of *Xhox-7.1* in posterior limb bud mesoderm impairs limb morphogenesis while inducing *BMP-4* expression, inhibiting cell proliferation, and promoting apoptosis. *Dev Biol* 197: 12–24
- Friedmann Y and Daniel CW (1996) Regulated expression of homeobox gene *Msx-1* and *Msx-2* in mouse mammary gland development suggests a role in hormone action and epithelial-stromal interactions. *Dev Biol* 177: 347–355
- Ganan Y, Macias D, Duterque-Coquillaud M, Ros MA, Hurlé JM (1996)

- Role of TGF β s and BMPs as signals controlling the position of the digits and the areas of interdigital cell death in the developing chick limb autopod. *Development* 122: 2349–2357
- Ganan Y, Macias D, Basco RD, Merino R, Hurlle J (1998) Morphological diversity of the avian foot is related with the pattern of *msx* gene expression in the developing autopod. *Dev Biol* 196: 33–41
- Gong SG, Kiba A (1999) The role of *Xmsx-2* in the anterior-posterior patterning of the mesoderm in *Xenopus laevis*. *Differentiation* 65: 131–140
- Gawantka V, Delius H, Hirschfeld K, Blumenstock C, Niehrs C (1995) Antagonizing the Spemann organizer: role of the homeobox gene *Xvent-1*. *EMBO J* 14: 6268–6279
- Graff JM, Thies SR, Song JJ, Celeste AJ, Melton DA (1994) Studies with a *Xenopus* BMP receptor suggest that ventral mesoderm-inducing signals override dorsal signals in vivo. *Cell* 79: 169–179
- Graham A, Heyman I, Lumsden A (1993) Even-numbered rhombomeres control the apoptotic elimination of neural crest cells from odd-numbered rhombomeres in the chick hindbrain. *Development* 119: 233–245
- Graham A, Francis-West P, Brickell P, Lumsden A (1994) The signaling molecule BMP4 mediates apoptosis in the rhombencephalic neural crest. *Nature* 372: 684–686
- Ishimura A, Maeda M, Takeda M, Kikkawa M, Daar IO, Maéno M (2000) Involvement of BMP-4/*msx-1* and FGF pathways in neural induction in *Xenopus* embryo. *Develop. Growth & Differ.* 42: 307–316
- Jones CM, Lyons KM, Lapan PM, Wright CVE, Hogan BLM (1992) *DVR-4* (bone morphogenetic protein-4) as a posterior-ventralizing factor in *Xenopus*. *Development* 115: 639–647
- Jowett AK, Vainio S, Ferguson MW, Sharpe PT, Thesleff I (1993) Epithelial-mesenchymal interactions are required for *msx1* and *msx2* gene expression in the developing murine molar tooth. *Development* 117: 461–470
- Kao KR and Elinson RP (1988) The entire mesodermal mantle behaves as Spemann's organizer in dorsoanterior enhanced *Xenopus laevis* embryos. *Dev Biol* 127: 64–77
- Ladher R, Mohun J, Smith JC, Snape A (1996) *Xom*: a *Xenopus* homeobox gene that mediates the early effects of BMP-4. *Development* 122: 2385–2394
- Mackenzie A, Ferguson MWJ, Sharpe PT (1992) Expression patterns of the homeobox gene, *Hox-8*, in the mouse embryo suggest a role in specifying tooth initiation and shape. *Development* 115: 403–420
- Maeda R, Kobayashi A, Sekine R, Lin JJ, Kung HF, Maéno M (1997) *Xmsx-1* modifies mesodermal tissue pattern along dorsoventral axis in *Xenopus laevis* embryo. *Development* 124: 2553–2560
- Maéno M, Ong RC, Suzuki A, Ueno N, Kung HF (1994) A truncated bone morphogenetic protein-4 receptor alters the fate of ventral mesoderm to dorsal mesoderm: Roles of animal pole tissue in the development of ventral mesoderm. *Proc Natl Acad Sci USA* 91: 10260–10264
- Mayor R, Morgan R, Sargent MG (1995) Induction of the prospective neural crest of *Xenopus*. *Development* 121: 767–777
- Monaghan AP, Davidson DD, Sime C, Graham E, Baldock R, Bhattacharya SS, Hill RE (1991) The *Msx*-like homeobox genes define domains in the developing vertebrate eye. *Development* 112: 1053–1061
- Moors JM, Wang S, Krinks M (1995) Anti-dorsalizing morphogenetic protein is a novel TGF- β homolog expressed in the Spemann organizer. *Development* 121: 4293–4301
- Muneoka K and Sassoon D (1992) Molecular aspects of regeneration in developing vertebrate limbs. *Dev Biol* 152: 37–49
- Nieuwkoop PD and Faber J (1967) *Normal Table of Xenopus laevis (Daudine)*. Amsterdam: North-Holland.
- Nohno T, Noji S, Koyama E, Nishikawa K, Myokai F, Saito T, Taniguchi S (1992) Differential expression of two *msh*-related homeobox genes *Chox-7* and *Chox-8* during chick limb development. *Biochem Biophys Res Commun* 182: 121–128
- Onichtchouk D, Gawantka V, Dosch R, Delius H, Hirschfeld K, Blumenstock C, Niehrs C (1996) The *Xvent-2* homeobox gene is part of BMP-4 signaling pathway controlling dorsoventral patterning of *Xenopus* mesoderm. *Development* 122: 3045–3053
- Papalopulu N and Kintner C (1996) A *Xenopus* gene, *Xbr-1*, defines a novel class of homeobox genes and is expressed in the dorsal ciliary margin of the eye. *Dev Biol* 174: 104–114
- Phippard DJ, Weber-Hall SJ, Sharpe PT, Naylor MS, Jayatalake H, Maas R, Hill RE, Dale TC (1996) Regulation of *Msx-1*, *Msx-2*, *Bmp-2* and *Bmp-4* during foetal and postnatal mammary gland development. *Development* 122: 2729–2733
- Schmidt JE, Suzuki A, Ueno N, Kimmel D (1995) Localized BMP-4 mediates dorsal/ventral patterning in early *Xenopus* embryo. *Dev Biol* 169: 37–50
- Semenza GL, Wang GL, Kundu R (1995) DNA binding and transcriptional properties of wild-type and mutant forms of the homeodomain protein *msx2*. *Biochem. Biophys Res Commun* 209: 257–262
- Shaine DH and Zuber MX (1996) Sodium dodecyl sulfate (SDS)-based whole-mount in situ hybridization of *Xenopus laevis* embryos. *Biochem Biophys Meth* 31: 185–188
- Shapira E, Marom K, Yelin R, Levy A, Fainsod A (1999) A role for the homeobox gene *Xvex-1* as part of the BMP-4 ventral signaling pathway. *Mech Dev* 86: 99–111
- Su MW, Suzuki HR, Solursh M, Ramirez F (1991) Progressively restricted expression of a new homeobox-containing gene during *Xenopus laevis* embryogenesis. *Development* 111: 1179–1187
- Suzuki A, Thies RS, Yamaji N, Song JJ, Wozney JM, Murakami K, Ueno N (1994) A truncated bone morphogenetic protein receptor affects dorsal-ventral patterning in the early *Xenopus* embryo. *Proc Natl Acad Sci USA* 91: 10255–10259
- Suzuki A, Ueno N, Hemmati-Brivanlou A (1997) *Xenopus msx-1* mediates epidermal induction and neural inhibition by BMP4. *Development* 124: 3037–3044
- Takahashi K, Nuckolls GH, Tanaka O, Semba I, Takahashi I, Dashner R, Shum L, Slavkin HC (1998) Adenovirus-mediated ectopic expression of *msx2* in even-numbered rhombomeres induces apoptotic elimination of cranial neural crest cells in ovo. *Development* 125: 1627–1635
- Takeda M, Saito Y, Sekine R, Onitsuka I, Maeda R, Maéno M (2000) *Xenopus msx-1* regulates the dorsoventral axis formation by suppressing the expression of organizer genes. *Comp Biochem Physiol* 126: 157–168
- Zhang H, Hu G, Wang H, Sciacivolino P, Iler N, Shen MM, Abate-Shen C (1997) Heterodimerization of *Msx* and *Dlx* homeoproteins results in functional antagonism. *Mol Cell Biol* 17: 2920–2932
- Zou H and Niswander L (1996) Requirement for BMP signaling in interdigital apoptosis and scale formation. *Science* 272: 738–741
- Wang Y and Sassoon D (1995) Ectoderm-mesenchyme and mesenchyme-mesenchyme interactions regulate *Msx-1* expression and cellular differentiation in the murine limb bud. *Dev Biol* 168: 374–382

(Received March 17, 2000 / Accepted May 20, 2000)