



Molecular Characterization of a cDNA Encoding Putative Vitellogenin from the Pacific Oyster *Crassostrea gigas*

Authors: Matsumoto, Toshie, Nakamura, Akihumi M., Mori, Katsuyoshi, and Kayano, Toshiaki

Source: Zoological Science, 20(1) : 37-42

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.20.37>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Molecular Characterization of a cDNA Encoding Putative Vitellogenin from the Pacific Oyster *Crassostrea gigas*

Toshie Matsumoto^{1*}, Akihumi M Nakamura², Katsuyoshi Mori² and Toshiaki Kayano³

¹National Research Institute of Aquaculture, Nansei, Mie 516-0193 Japan,

²Tohoku University, Sendai 981-8555 Japan,

³National Institute of Agrobiological Sciences,
Kannondai, Tsukuba 305-8602 Japan

ABSTRACT—To elucidate the molecular mechanisms involved in oogenesis, we applied a differential display method to identify genes whose expression was detected only in ovaries containing oocytes. One of the cDNA fragments isolated by mRNA differential display was similar in structure to vitellogenin. Using this fragment, a full-length cDNA encoding putative vitellogenin in the Pacific oyster *Crassostrea gigas* was cloned by RACE (rapid amplification of cDNA ends), and its amino acid sequence was deduced. The open reading frame predicted 1583 amino acid residues. The deduced primary structure of putative vitellogenin in *C. gigas* was shown to be similar to vitellogenins of various other mollusk, fish, crustacean and nematode species, especially in the N-terminal region. Reverse transcription-mediated PCR revealed that mRNA encoding putative vitellogenin was expressed only in the ovary. *In situ* hybridization analysis revealed that putative vitellogenin mRNA was expressed strongly in the follicle cells in the ovary. It is concluded that the follicle cells are the site of putative vitellogenin synthesis.

Key word: follicle cells, mRNA differential display, oogenesis, oyster, vitellogenin

INTRODUCTION

In marine bivalves as in most oviparous animals, a large amount of yolk protein (vitellin) is accumulated in oocytes during ovarian maturation. In vertebrates such as fish, amphibians and birds, yolk protein is synthesized from a precursor, vitellogenin (Vg), produced by the liver, and is transported to the oocytes via the blood circulation system. The relationship between yolk protein and Vg has been extensively studied in *Xenopus laevis* and in the chicken, *Gallus gallus*, and the amino acid sequences of these species have been determined (Gerber-Hunber *et al.*, 1987; Van het Schip *et al.*, 1987). Vgs have been shown to be present in almost all species of oviparous animals ranging from nematodes to vertebrates, and extensive sequence conservation is seen among these groups (Chen *et al.*, 1997). Recently, complete cDNA sequences of Vg have been reported from several fish and crustacean species (LaFleur *et al.*, 1995; Mouchel *et al.*, 1996; Tsutsui *et al.*, 2000; Okuno *et al.*, 2002).

In bivalve molluscs, a few biochemical studies of yolk proteins have been carried out (Osada *et al.*, 1992; Suzuki

et al., 1992). In the oyster, vitellin has been isolated and characterized (Suzuki *et al.*, 1992; Li *et al.*, 1998), but the characterization of Vg has not been reported, and information on bivalve vitellogenesis is still limited. To identify the genes associated with vitellogenesis, we cloned and sequenced more than 100 cDNA fragments using mRNA differential display, which is a method for studying differential gene expression from different sources (Liang and Pardee, 1992), and found that one of the isolated genes was similar to Vg. The objective of the present study was to clone the full-length of the Pacific oyster putative Vg cDNA on the basis of the cDNA fragment obtained by mRNA differential display, to deduce the complete primary amino acid sequence, and to examine the expression of its mRNA.

MATERIALS AND METHODS

Animals and isolation of RNA

Cultured Pacific oysters were collected monthly from March to September 1999 in Gokasho Bay, Mie Prefecture. The gonad was homogenized using ISOGEN (Nippon Gene, Toyama, Japan), and the total RNA was extracted according to the manufacturer's instructions (Chomzynski and Sacchi, 1987).

mRNA differential display

First strand cDNA was synthesized by annealing pd(N)₆Primer per 5 µg of total RNA using a First Strand cDNA Synthesis kit

* Corresponding author: Tel. +81-599-66-1830;
FAX. +81-599-66-1962.
E-mail: mtosie@fra.affrc.go.jp

(Amersham Biosciences Corp., NJ), according to the manufacturer's protocol. The resulting cDNAs were amplified with seven different 10-nucleotide random primers (RA1, 3, 5, 7, 13, 14 and 16) selected from the list of Monna *et al.* (1994). The cycling parameters were as follows: one cycle of 92°C for 1 min, and 40 cycles of 92°C for 1 min, 45°C for 1 min, 72°C for 1 min and lastly one cycle of 72°C for 5 min. The amplified cDNA was then separated on an 8% polyacrylamide gel and stained with ethidium bromide. Complementary DNA fragments whose expression was detected only in the spawning season (July and August) were recovered directly by cutting out the gel slices, and reamplified using the same primer and PCR conditions except an annealing temperature at 55°C. Reamplified PCR fragments were subcloned and sequenced.

Oligonucleotides

The oligonucleotides used as PCR primers are listed here and shown in Fig. 2. Primers for β -actin were based on the nucleotide sequences of sea scallop and oyster (Patwary *et al.*, 1996; Cadoret *et al.*, 1999).

- Primer 1. 5'- ACG GGT AAG TAC TGT TGG AC -3'
 Primer 2. 5'- GAC GAA TGT TCC GAC GTG GG -3'
 Primer 3. 5'- CCC GAT GTC AAT GGC TTA TGG -3'
 Primer 4. 5'- GCA GAT GGA AGG ATG TCC ATC AG -3'
 Primer 5. 5'- TTC ACA GTC ATG GAG CCC AGC AT -3'
 Primer 6. 5'- GAC CCG TCA GAA TTG TTG TCA GAC -3'
 Primer 7. 5'- GAC TTC GAA CAA GAG ATG -3'
 Primer 8. 5'- GAT ATC GAC ATC ACA TTT C -3'

Rapid amplification of cDNA ends (RACE)

5' RACE was performed using a 5' RACE System (Invitrogen Corp., Carlsbad, CA) according to the manufacturer's instructions (Frohman *et al.*, 1988). First-strand cDNA was synthesized in a 25 μ l solution containing 5 μ g of total RNA, 2.5 pM of primer 1, and 200 U of Superscript II. PCR with primer 2 and 5' RACE abridged anchor primer was initiated at 94°C for 2 min, followed by 30 cycles of 94°C for 0.5 min, 55°C for 0.5 min, 72°C for 2 min and lastly one cycle of 72°C for 7 min. For 3'RACE, first strand cDNA was synthesized by annealing Not I-d(T)₁₈Primer using a First strand cDNA Synthesis kit (Amersham Biosciences) as described above. PCR was conducted with primer 3 and adapter primer using the same conditions but 35 cycles of amplification. The PCR products were subcloned and sequenced. Since these fragments were too long to sequence, PCR walking was performed with the designed primers (primers 4, 5 and 6).

Nucleotide sequence analysis

The cDNA fragments amplified by PCR were subcloned into a pCR2.1 plasmid using a TA Cloning kit (Invitrogen). Both strands of the insert in the plasmid were sequenced on a model 377 DNA sequencer using a BigDye Terminator Cycle Sequencing kit (Applied Biosystems, Foster City, CA).

Reverse transcription-mediated PCR

Total RNA was prepared from ovary, gill, muscle, mantle tissues of female oysters collected in April, and testis and ovary of the oysters collected in March, May, July and September, and first strand cDNA was synthesized as described above. Twenty-five (Fig. 5A) and 20 (Fig. 5B) cycles of amplification for vitellogenin (primers 4 and 5) and β -actin (primers 7 and 8) using AmpliTaq Gold (Applied Biosystems) were carried out under the following conditions: denaturation at 94°C for 0.5 min, annealing at 54°C for 0.5 min, and extension at 72°C for 0.5 min. At completion of the PCR, fragments were separated on a 1.5% agarose gel and stained with ethidium bromide.

In situ hybridization

For *in situ* hybridization, the digoxigenin (DIG)-labeled sense

and antisense RNA probes were transcribed *in vitro* from 807 bp cDNA fragment of oyster putative Vg (region between primers 4 and 5) using a DIG RNA Labeling kit SP6/T7 (Roche Diagnostics, Mannheim, Germany). The ovarian fragments were fixed in 4% paraformaldehyde in 0.1 M phosphate buffer at pH 7.4 for 16 hr, embedded in paraffin, and cut into 5 μ m serial sections. Deparaffinized sections were treated with proteinase K (10 μ g/ml) in PBS at 37°C for 15 min, and then incubated at 42°C for 16 hr in a hybridization buffer (50% formamide, 2 x SSC, 1 μ g/ μ l tRNA, 1 μ g/ μ l Salmon sperm DNA, 1 μ g/ μ l BSA, 10% Dextran sulfate) and 1 mg/ml DIG-labeled probe. After hybridization, the sections were washed three times with 2 x SSC/50% formamide at 42°C for 1 hr and twice with 2 x SSC at room temperature for 15 min. The sections were incubated with the Fab fragment of an anti-DIG alkaline phosphatase-conjugated antibody (1:500 dilution, Roche Diagnostics), and visualized with nitroblue tetrazolium and 5-bromo-4-chloro-3-indolyl phosphate (Roche Diagnostics).

RESULTS AND DISCUSSION

Isolation and characterization of cDNA encoding putative Vg

In order to identify the expressed genes in the ovary, transcripts derived from the ovaries during the spawning season and the spent gonads containing no oocytes were examined by mRNA differential display using seven different primers. During this procedure, we identified a cDNA of approximately 1.2 kb in length, which was detected in the ovary during the spawning season (Fig. 1). Sequence analysis of the cDNA fragment followed by a database search revealed a similarity to Vg (Fig. 2: C1184).

Subsequently the 5' and 3' ends of this cDNA were obtained by 5' and 3' RACE with the gene-specific primers, respectively (Fig. 2). The deduced amino acid sequence of




Fig. 1. Differential display of mRNAs from ovary (July and August) and spent gonad (September) of oyster. RNA was reverse transcribed with random primer. Resulting cDNA was amplified with an arbitrary primer (RA16, CCGACAGCTT). A differentially expressed fragment (arrow) was recovered, eluted from the gel slice, and reamplified. Numbers to the right represent molecular size markers.




Fig. 2. Cloning strategy and schematic view of *C. gigas* putative vitellogenin cDNA. Location of the specific primers is shown corresponding to oligonucleotides listed in the Materials and Methods. The obtained cDNA is delineated, showing positions of the start codon (13 ATG) and stop codon (4762 TGA). C1184 indicates the PCR product obtained by mRNA differential display.

C. gigas putative Vg cDNA is shown in Fig. 3. The cDNA consisted of 5023 bp, comprising a 5' untranslated region (12 bp), and open reading frame (4749 bp), a stop codon (TGA), and a 3' untranslated region (259 bp). The 3' untranslated region contained a polyadenylation signal (AATAAA) (Fig. 3).

Characterization of the deduced amino acid sequence

The cDNA had an ORF (open reading frame) that encoded 1583 amino acid residues with a predicted molecular mass of 179,191 Da. The deduced amino acid sequence contained a consensus cleavage site, R-X-R-R (Arg⁸²⁸ to Arg⁸³¹), capable of undergoing processing by endoproteases of the subtilisin family (Barr, 1991) that have also been reported in Vgs of insects and crustaceans (Chen *et al.*, 1994; Yano *et al.*, 1994; Tsutsui *et al.*, 2000; Okuno *et al.*, 2002). If this processing site is cleaved by an endoprotease, the resultant two subunits would have calculated molecular masses of 94,000 Da and 85,209 Da. Previous studies have shown that in the Pacific oyster *C. gigas*, two major bands (179 and 110 kD) and several minor bands (Li *et al.*, 1998), and seven protein bands with relative molecular masses of 105, 85, 66, 64, 60, 45 and 41 kD (Suzuki *et al.*, 1992) were recognized in SDS-immunoblot analysis of the vitellin. The cDNA isolated in the present study may correspond to one or some of these proteins. The overall serine content of the oyster putative Vg (11.9%) was higher than in the mosquito (10.1%) and the silkworm Vg (9.8%), which possess polyserine clusters. The deduced amino acid sequence contained 56.2% of serine residues in the region between Ser¹¹⁴⁹ and Ser¹¹⁸⁰, and this region may be the polyserine domain in *C. gigas* putative Vg. This protein possessed six potential N-linked glycosylation sites, which is conserved in vertebrate and invertebrate Vgs.

The deduced amino acid sequence was compared to those of Vgs of the scallop *Patinopecten yessoensis* (partial-length, accession number AB055960), *Caenorhabditis elegans* (Spieth *et al.*, 1991), *Fundulus heteroclitus* (LaFleur *et al.*, 1995), rainbow trout (Mouchel *et al.*, 1996), and kuruma prawn (Tsutsui *et al.*, 2000). The BLAST algorithm indicated

homologies of 35% to scallop Vg (residues 24–519), 21% to *C. elegans* vit-5 (residues 24–954), 19% and 24% to *Fundulus* Vg1 (residues 22–285 and 454–953, respectively), 25% and 22% to rainbow trout Vg (residues 24–266 and 567–932, respectively) and 28% to kuruma prawn Vg (residues 559–940). As shown in Fig. 4, the region of *C. gigas* putative Vg which showed similarity to other Vg was limited to the N-terminal and the central region. The C-terminal region showed no significant homology to any of the other reported species. Thus this region appears to possess unique properties compared with Vg of the other known species. Best conserved amino acid motif between invertebrate and vertebrate Vgs (Mouchel *et al.*, 1996) is found in the sequence of *C. gigas* putative Vg (KTIGNAG: positions 603–609). This sequence is speculated to play a highly conserved role in vitellogenesis, such as specific recognition by oocytes (Spieth *et al.*, 1991).

Tissue distribution of oyster putative Vg mRNA

The levels of putative Vg mRNA in various tissues from female oyster and stage-specific expression were measured by reverse transcription-mediated PCR. In April, the oyster contained fully grown oocytes and putative Vg mRNA expression was detected only in the ovary (Fig. 5A). As Fig. 5B shows, putative Vg mRNA expression was detected only in the ovary, and indicated maximum level in March (the early stage of maturation). To determine the distribution of oyster putative Vg mRNA expression in ovary, we performed *in situ* hybridization using DIG-labeled sense and antisense RNA probes, respectively. A strong signal was detected in the follicle cells using an antisense RNA probe (Fig. 6A). In contrast, hybridization with a sense control probe produced no significant signal (Fig. 6B).

The synthesis, secretion and processing of Vgs differ among phyla. Vgs are synthesized by extraovarian tissues such as the liver in vertebrates and the fat body in insects (Sappington and Raikhel, 1998), secreted into the circulatory system, and transported into the ovary. In crustaceans, three tissues have been reported as sites of Vg synthesis:

```

1 TTACACGGCAGAAATGTTGCTTCTGCTTGC GGTTCAACAGCCCTGGTGGCGGCAGCCGCCAGACTCCCAGTCAAGCCTATGAAAGCAAT
      M L L L L A V S T A L V A A A A Q T P S Q A Y E S N      26
KEYLYEYETQALTI PMGSTIYSGMKMKS DVRIQFRSRS SATLKM DKLTF AKINDPIESVDPTQQQVPAEMFQPLTGRDAEQMLTDLSRP      116
INFRYVRGNVKDIHHEADDP EWSVNVK KGLLSMLEMNL EKRELKRS GVI PQVLRPQSSDEGSMFTVMEPSIVGECETLYRISPWTSTSA      206
NPWMHITKVRNYQHCLDRPKYFGSMFHLRQCAECVREQSEPLRSASQIRYTLRGNMRQFQIQSAIAESQHVFTFPYSAKGGHVATYLNQTL      296
NLVKEEDVKTSLSE PQSP TKVKSGLQYTSRELEMRDSSS ILQSSDSLSTDKSSPESPQHSSSSSNHHPSPQPPSPQPKTTSNQAADRIR      386
KLLKMLEVFMKPTIQPEAGP LLMSS LLEEIRTADES LRAVYRECSNGETAATVQDADGHPSICGTT SATEVLVDS ILNDQLPQSEAVVAL      476
GILSLSARP DVLI AKLLDL TRSSQVSKDRYLKRAAFCLG SVAGMLREEGWRRSREISRQEEIVKTLISQESRSQTRNELKSKKRELED      566
LKKREHS IKTKIKQEI VKELOMLMRSTVFDDKILSFKTIGNAGLWEMIPITRTYIQDKSQPQVLRQTQAIYSLRKLARHPDDIQTLLPL      656
YFDQSEKEEVRIGSYLVMTFTEPSRQLLEMVAQSLHRE RNPHVGT FVYTHLEQMSNSTYPCIMSWAKNASFAMRF AKKFS PMYHYSRFMH      746
LSGFNEMHKMGAAAELGLVTTPEEFI PRAGAVNLHTYVFGRSV NFAEIGFNT EQLQTLVSKLVG PLGELTKGKSLVDVLKQ EVRR SAESS      836
QPSDPI SQIHKQLKVPRTSPTPKGHMYMKMMG NELQYITLDGTLVD TLLREGKLLSGVSEQDLKTGLNVEVHRS TMPLEAEIMIPSECG      926
LPLRLKLHGTA AIKVTGKVGVTGMP SIFEINRPGKQAKELS FN FELRPSVLFQLRGEMEMDAEYFKMGV ALKTM AHMETPLSLTASANLP      1016
KAKFYTKFNIEKLSEKIARLEVS PYTYFKEDPSEITKYPLPRETQEISVAKNAKVFMSMAYGKQTLGLEIKLSGQAVVRDFEMDVPYYP      1106
MIGKQEMIVT LSPGTD PKYVEIQFQLMKRMSQSKQPTSEESSSSTGLLSWVSSLI GSDQSDQSSPSSSSSSSTSPPKDIRSKDTS LKD      1196
LLEHLKESQSVHPDGSVTRNNSMGLIFNVIGIDQSR SIKRHFHISMAAGMNP SKTTSI LLMNRSPIPTMETKPNMVDVAVNVNLP SR      1286
LADPSELLSDAYQRELEQQIEVYVRRHGESDYHRWIMLEEPRWESSVLQRLLSLSDTQEVIDEVMNSELSKSEESGTPEDRNIIDLIRKQ      1376
KQILKEARKIWQRSYDSL RVQDIKAKLSS LIKRSESVRKDLEKDSASVSKSEKSDALRKYCALLQQLIDQRLEAVRSSRRSLSSKERQD      1466
IAKSLESSKESLRKILQQHRSPSKVDSS IHHLALFKSILHRSVKVILK LKNLRTQNKKINPSNNSLNSPRT PCLAINLPKCLVVRL      1556

4681 TGTCAAAGTCAGTGT CAGACCTTTTGCAAAAACAGAAAGATGTGGTTCAGGATATTTCCAGGTCAAAGGAATCAGAGGGATTGAAATCTG
      C Q S Q C Q T F C K N R K M W F R I F P G Q R N Q R D *      1583
4771 TTGATGAGAAAAAATCAAGGAGGC ACTTCTTTTAGCAAAAAGATTACTGAAAAGATTCTGTC AAGAAAATTCTGGCCCATCTCTCAA
4861 ATATCGAGAGGGAAAGACAAAGACTCATGAAATCCGTCATTCAACAACAGCAAATAATGAAATGTTTGGAGCAGCGGGTAAAGAAACCAC
4951 AAACCGCCTCCGGCAAATCTGAAGACCA ATAAAA GAAATTTCAGAGAATTATGCAAAAAAAAAAAAAAAAAAAAAA      5023

```

Fig. 3. Deduced amino acid sequence and 5' and 3' untranslated regions of putative Vg cDNA. Positions of nucleotides and amino acids are indicated by numbers on the left and right sides of the figure, respectively. Putative *N*-glycosylation sites are underlined and consensus cleavage sequence for processing by endoproteases of the subtilisin family is framed. The polyadenylation signal is shown in white lettering with black background. The position of DIG-labeled RNA probe used for *in situ* hybridization is indicated by a dotted underline. The nucleotide sequence has been submitted to DDBJ/EMBL/GenBank with the accession number AB084783.

the ovary (Lee and Watson, 1995; Tsutsui *et al.*, 2000), adipose tissue (Okuno *et al.*, 2000) and hepatopancreas (Soroka *et al.*, 2000; Tsutsui *et al.*, 2000; Okuno *et al.*, 2002). Autosynthetic yolk formation is thought to be the main type of vitellogenesis in bivalves and gastropods (Jong-Brink *et al.*, 1983). In marine bivalves, autosynthesis of yolk proteins in the ovary has been postulated, based on the morphological evidence (Pipe, 1987; Dorange and Le Pennec, 1989; Eckelbarger and Davis, 1996). In oyster and other bivalve species, the ovarian acinus is a simple structure containing only developing oocytes and associated follicle cells within a thin germinal epithelium. The function(s) of follicle cells in the bivalve ovary are not well understood, though they are suspected of playing some role in oocyte nutrition. In the present study putative Vg mRNA expression

was in the follicle cells in the ovary of *C. gigas*. This suggests that yolk synthesis in *C. gigas* may take place in the follicle cells.

In teleosts, as in other oviparous vertebrates, it is clearly established that the Vg gene expression is regulated by estradiol-17 β (E_2), produced by the maturing female ovarian follicles, in the liver (Vaillant *et al.*, 1988; Pakdel *et al.*, 1991; Okumura *et al.*, 2002). In the Pacific oyster, E_2 has been detected in the ovary, and its content shows a synchronous profile with gonadal maturity and decreased in the spawning season (Matsumoto *et al.*, 1997). This is correspondent to the change of putative Vg mRNA levels from early vitellogenic stage to spawning stage, suggesting that E_2 is at least one of the factors which promotes vitellogenesis in oysters. However, further investigation is required in




Fig. 4. Alignment of *C. gigas* putative Vg with other animal Vgs: scallop *Patinopecten yessoensis* (partial-length, accession number AB055960); *Caenorhabditis elegans* (Spieth *et al.*, 1991); *Fundulus heteroclitus* (LaFleur *et al.*, 1995); rainbow trout (Mouchel *et al.*, 1996); and kuruma prawn (Tsutsui *et al.*, 2000). The open bar represents the deduced amino acid sequence of *C. gigas* putative Vg. The solid bars indicate the regions of Vg which have homology to those of other animals. Percentage of homology is also indicated.




Fig. 5. Distribution of putative Vg mRNA in tissues of oyster. RNA was prepared from ovary, gill, muscle, mantle of female oysters collected in April (A), testis and ovary in March, and ovary in May, July and September (B). β -actin was used as a positive control.




Fig. 6. Localization of putative Vg mRNA in the ovary by *in situ* hybridization. Strong expression of putative Vg was seen in the follicle cells with an antisense probe (A) in the ovary. Hybridization with a sense probe (B) produced no significant signal in the ovary. Bar=25 μ m.

order to clarify the estrogen regulatory mechanisms in the oyster.

ACKNOWLEDGMENTS

This work was supported by the Integrated Research Program for Effects of Endocrine Disrupters on Agriculture, Forestry and Fisheries and Their Action Mechanisms on Domestic Animals and Fishes (ED-02-II-3-2).

REFERENCES

- Barr PJ (1991) Mammalian subtilisins: the long-sought dibasic processing endoproteases. *Cell* 66: 1–3
- Cadoret JP, Debon R, Cornudella L, Lardans V, Morvan A, Roch P, Boulo V (1999) Transient expression assays with the proximal promoter of a newly characterized actin gene from the oyster *Crassostrea gigas*. *FEBS Lett* 460: 81–85
- Chen JS, Cho WL, Raikhel AS (1994) Analysis of mosquito vitellogenin cDNA. Similarity with vertebrate phosphatases and arthropod serum proteins. *J Mol Biol* 237: 641–647
- Chen JS, Sappington TW, Raikhel AS (1997) Extensive sequence conservation among insect, nematode, and vertebrate vitellogenins reveals ancient common ancestry. *J Mol Evol* 44: 440–451
- Chomzynski P and Sacchi N (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162: 156–159
- Dorange G and Le Pennec M (1989) Ultrastructural study of oogenesis and oocytic degeneration in *Pecten maximus* from the Bay of St. Brieuc. *Mar Biol* 103: 339–348
- Eckelbarger KJ and Davis CV (1996) Ultrastructure of the gonad and gametogenesis in the eastern oyster, *Crassostrea virginica*. I. Ovary and oogenesis. *Mar Biol* 127: 79–87
- Frohman MA, Dush MK, Martin GR (1988) Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer. *Proc Natl Acad Sci USA* 85: 8998–9002
- Gerber-Hunber S, Nardelli D, Haefliger JA, Cooper DN, Givel F, Germond JE, Engel J, Green NM, Wahli W (1987) Precursor-product relationship between vitellogenin and the yolk proteins as derived from the complete sequence of a *Xenopus* vitellogenin gene. *Nucleic Acids Res* 15: 4737–4753
- Jong-Brink M, Boer HH, Joosse J (1983) Mollusca. In "Reproductive biology of invertebrates. Vol 1." Eds by KG Adiyodi, RG Adiyodi, John Wiley & Sons Ltd, New York, pp 297–355
- LaFleur GJ Jr, Byrne BM, Kanungo J, Nelson LD, Greenberg RM, Wallace RA (1995) *Fundulus heteroclitus* vitellogenin: the deduced primary structure of a piscine precursor to noncrystalline, liquid-phase yolk protein. *J Mol Evol* 41: 505–521
- Lee CY and Watson RD (1995) *In vitro* study of vitellogenesis in the blue crab (*Callinectes sapidus*): site and control of vitellin synthesis. *J Exp Zool* 271: 364–372
- Li Q, Osada M, Suzuki T, Mori K (1998) Changes in vitellin during oogenesis and effect of estradiol-17 β on vitellogenesis in the Pacific oyster *Crassostrea gigas*. *Invert Reprod Develop* 33: 87–93
- Liang P and Pardee AB (1992) Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science* 257: 967–971
- Matsumoto T, Osada M, Osawa Y, Mori K (1997) Gonadal estrogen profile and immunohistochemical localization of steroidogenic enzymes in the oyster and scallop during sexual maturation. *Comp Biochem Physiol B* 118: 811–817
- Monna L, Miyao A, Inoue T, Fukuoka S, Yamazaki M, Zhong HS, Sasaki T, Minobe Y (1994) Determination of RAPD markers in rice and their conversion into sequence tagged sites (STSs) and STS-specific primers. *DNA Res* 1: 139–148
- Mouchel N, Trichet V, Betz A, Le Pennec JP, Wolff J (1996) Characterization of vitellogenin from rainbow trout (*Oncorhynchus mykiss*). *Gene* 174: 59–64
- Okumura H, Todo T, Adachi S, Yamauchi K (2002) Changes in hepatic vitellogenin mRNA levels during oocyte development in the Japanese eel, *Anguilla japonica*. *Gen Comp Endocrinol* 125: 9–16
- Okuno A, Katayama H, Nagasawa N (2000) Partial characterization of vitellin and localization of vitellogenin production in the terrestrial isopod, *Armadillidium vulgare*. *Comp Biochem Physiol B* 126: 397–407
- Okuno A, Yang WJ, Jayasankar V, Saido-Sakanaka H, Huong Do TT, Jasmani S, Atmomarsono M, Subramoniam T, Tsutsui N, Ohira T, Kawazoe I, Aida K, Wilder MN (2002) Deduced primary structure of vitellogenin in the giant freshwater prawn, *Macrobrachium rosenbergii*, and yolk processing during ovarian maturation. *J Exp Zool* 292: 417–429
- Osada M, Unuma T, Mori K (1992) Purification and characterization of a yolk protein from the scallop ovary. *Nippon Suisan Gakkai-shi* 58: 2283–2289
- Pakdel F, Feon S, Le Gac F, Le Menn F, Valotaire Y (1991) *In vivo* estrogen induction of hepatic estrogen receptor mRNA and correlation with vitellogenin mRNA in rainbow trout. *Mol Cell Endocrinol* 75: 205–212
- Patwary MU, Reith M, Kenchington EL (1996) Isolation and characterization of a cDNA encoding an actin gene from sea scallop (*Placopecten magellanicus*). *J Shellfish Res* 15: 265–270
- Pipe RK (1987) Oogenesis in the marine mussel *Mytilus edulis*: an ultrastructural study. *Mar Biol* 95: 405–414
- Sappington TW and Raikhel AS (1998) Molecular characteristics of insect vitellogenins and vitellogenin receptors. *Insect Biochem Mol Biol* 28: 277–300
- Soroka Y, Milner Y, Sagi A (2000) The hepatopancreas as a site of yolk protein synthesis in the prawn *Macrobrachium rosenbergii*. *Invert Reprod Develop* 37: 61–68
- Spieth J, Nettleton M, Zucker-Aprison E, Lea K, Blumenthal T (1991) Vitellogenin motifs conserved in nematodes and vertebrates. *J Mol Evol* 32: 429–438
- Suzuki T, Hara A, Yamaguchi K, Mori K (1992) Purification and immunolocalization of a vitellin-like protein from the Pacific oyster *Crassostrea gigas*. *Mar Biol* 113: 239–245
- Tsutsui N, Kawazoe I, Ohira T, Jasmani S, Yang WJ, Wilder MN, Aida K (2000) Molecular characterization of a cDNA encoding vitellogenin and its expression in the hepatopancreas and ovary during vitellogenesis in the kuruma prawn, *Penaeus japonicus*. *Zool Sci* 17: 651–660
- Vaillant C, Le Guellec C, Pakdel F, Valotaire Y (1988) Vitellogenin gene expression in primary culture of male rainbow trout hepatocytes. *Gen Comp Endocrinol* 70: 284–290
- Van Het Schip FD, Samallo J, Broos J, Ophuis J, Mojet M, Gruber M, Ab G (1987) Nucleotide sequence of a chicken vitellogenin gene and derived amino acid sequence of the encoded yolk precursor protein. *J Mol Biol* 196: 245–260
- Yano K, Sakurai MT, Watabe S, Izumi S, Tomino S (1994) Structure and expression of mRNA for vitellogenin in *Bombyx mori*. *Biochim Biophys Acta* 1218: 1–10

(Received June 17, 2002 / Accepted September 30, 2002)