

Campylium longicuspis (Lindb. & Arnell) Hedenäs (Bryophyta, Amblystegiaceae), another Arctic moss in the northern Scandinavian mountain range

Authors: Jacobson, Conny, and Hedenäs, Lars

Source: *Lindbergia*, 38(1) : 17-19

Published By: Dutch Bryological and Lichenological Society and Nordic
Bryological Society

URL: <https://doi.org/10.25227/linbg.01069>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Campylium longicuspis (Lindb. & Arnell) Hedenäs (Bryophyta, Amblystegiaceae), another Arctic moss in the northern Scandinavian mountain range

Conny Jacobson and Lars Hedenäs

C. Jacobson (conny.jacobson@naturvardsverket.se), Swedish Environmental Protection Agency, SE-106 48 Stockholm, Sweden. – *L. Hedenäs* (lars.hedenas@nrm.se), Swedish Museum of Natural History, Dept of Cryptogamic Botany, Box 50007, SE-104 05 Stockholm, Sweden

The conditions at the first Swedish and mainland European *Campylium longicuspis* locality are described and compared with the few so far reported finds of this species. The species appears to prefer chiefly calcium-rich wetland habitats. *Campylium longicuspis* is mainly Arctic, and a map of its known distribution is presented.

The occurrence of predominantly Arctic or even high-Arctic moss species in the northern portion of the Scandinavian mountain range has for a long time been of great interest to Scandinavian bryologists (Mårtensson 1956). Examples of such species include *Funaria arctica* (Berggr.) Kindb. (Rumsey 1990), *Psilopilum cavifolium* (Wilson) I.Hagen (Long 1985, Lübenau and Lübenau 1971), *Rhizomnium andrewsianum* (Steere) T.J.Kop. (Engelmark 1974), and *Timmia sibirica* Lindb. & Arnell (Carlsson 2003). The occurrences of this kind of species add to the bryological temptations offered by the Scandes, but nowadays these species are also of great interest for another reason. The ongoing gradual climate change will most likely affect these species, which presumably depend on relatively cold conditions, relatively soon. It is therefore important both to find localities of such species, while the species still grow there, and to keep track of their future destiny.

During the summer of 2007 CJ carried out an investigation in the alpine region of the Swedish mountain range to improve our knowledge about the habitat “Pioneer formations of *Caricion bicoloris-atrofuscae*”, a habitat protected by the EU habitats directive (EEC 1992). In a small, 0.25 m², field plot on Mt Pältsa in the northernmost portion of Sweden a few scattered, unbranched moss shoots were noticed. One shoot that was collected, and later identified by LH, turned out to belong to *Campylium longicuspis* (Lindb. & Arnell) Hedenäs (Hedenäs 1989).

This species reminds a bit about *C. laxifolium* Engelmark & Hedenäs, but has longer (2.1–4.6 versus 1.8–2.4 mm) and more strongly concave stem leaves. The leaves are more narrowly ovate in their basal portion and have frequently got a short differentiated acumen, constituting 18–33% of the leaf length. During a second visit to the locality in 2011 by N. Lönnell and T. Hallingbäck the species was again found.

Globally, *Campylium longicuspis* was overlooked for a long time, but its distribution is now known to include large portions of the Arctic, with outliers in the Altai Mountains (Fig. 1). The present find is the first from the European mainland. In addition, *C. longicuspis* was reported from both Serbia and Montenegro (Sabovljevic et al. 2004). However, this is likely due to a mistake in interpreting the synonymy of *C. longicuspis* and *C. protensum* (Brid.) Kindb. (Martinčić 2009), and in addition it seems very unlikely that the Arctic *C. longicuspis* should occur in the Balkan area.

In the literature, *C. longicuspis* was reported both together with species of calcium-rich wetlands, such as *Drepanocladus trifarius* (F.Weber & D.Mohr) Paris, *D. turgescens* (T.Jensen) Broth., and *Tomentypnum nitens* (Hedw.) Loeske (Arnell, H. W. 1913, Arnell, S. and Mårtensson 1959, Lindberg and Arnell 1890) and in one case together with *Blepharostoma trichophyllum* (L.) Dumort., *Scapania simmonsii* Bryhn & Kaal., and *Ceratodon* (Arnell, H. W. 1913). Since the habitat of *C. longicuspis* is still rather




Figure 1. Distribution of *Campylium longicuspis*, with the N Scandinavian locality indicated with a dot having a double line. The map is based on material present in S, and the literature (Afonina 2004, Czernyadjeva 2001, Fedosov, et al. 2011, Hedenäs 1989, 2003, Ignatov 1994, Ignatov, et al. 2006). The Canadian record (?) is based on information in (Afonina 2004) where this was published as from 'Arctic Canada', but with no more exact locality information.

poorly known (Hedenäs 2003), some observations from its newly found north Swedish site follow (Fig. 2). The site is located in a gentle slope, heading towards SW in a mountain pass. The vegetation is sparse with no trees or shrubs. The habitat of the species consists of very thin soil, most likely calcareous, often with trickling water or spring influence from the slopes nearby. Within the 0.25-m² field plot with *C. longicuspis*, the following bryophytes were recorded: *Aneura pinguis* (L.) Dumort., *Blepharostoma trichophyllum*, *Blindia acuta* (Hedw.) Bruch & Schimp., *Bryum* sp., *Campylium stellatum* (Hedw.) C.E.O.Jensen, *Cephalozia* sp., *Hylocomium splendens* (Hedw.) Schimp., and *Scorpidium cossonii* (Schimp.) Hedenäs. In the same

plot, 18 species of vascular plants were recorded, of which *Bistorta vivipara* (L.) Gray, *Carex bicolor* All., *Saxifraga aizoides* L., *S. oppositifolia* L., *Saussurea alpina* (L.) DC., *Silene acaulis* (L.) Jacq., and *Thalictrum alpinum* L. were most common. Of the other ones, *Carex capillaris* L., *C. fuliginosa* Schkuhr, *Juncus biglumis* L., *Salix reticulata* L., and *Tofieldia pusilla* (Michx.) Pers. can be mentioned.

Cited specimen: **Sweden:** Torne Lappmark, Kare-suando, Mt Pältsa, plot 1 (WP 669), 69°00'28.7"N, 20°23'68.6"E (WGS-84), 1090 m a.s.l., thin soil with trickling water, most likely calcareous and sometimes with spring influence, leg. C. Jacobson, 10 August 2007, det. L. Hedenäs (S; reg. No. B183239).



Figure 2. The 0.25-m² field plot with *Campylopus longicuspis* on Mt Pältsa in northernmost Sweden.

References

- Afonina, O. M. 2004. Konspekt flori mchov Tjukotki {Moss flora of Chukotka}. – Rossijskaja Akademia Nauk. Botanitjeskij Institut im. V. L. Komarov RAN, Sankt-Peterburg.
- Arnell, H. W. 1913. Zur Moosflora des Lena-Tales. – Ark. Bot. 13: 1–94.
- Arnell, S. and Mårtensson, O. 1959. A contribution to the knowledge of the bryophyte flora of W. Spitsbergen and Kongsfjorden (Kings Bay, 79° N) in particular. – Ark. Bot. 4: 105–164.
- Carlsson, P. 2003. *Timmia sibirica* och lite annat smått och gott från Padjelanta. – Myrinia 13: 61–68.
- Czernyadjeva, I. V. 2001. Moss flora of Yamal Peninsula (West Siberian Arctic). – Arctoa 10: 121–150.
- EEC 1992. Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. – Off. J. Eur. Communities L206: 7–50.
- Engelmark, T.-B. 1974. *Trichostomum cuspidatissimum* Card. & Thér. and *Rhizomnium andreusianum* (Steere) Kop. discovered in Northern Sweden. – Lindbergia 2: 218–221.
- Fedosov, V. E., Ignatova, E. A., Ignatov, M. S. et al. 2011. Rare species and preliminary list of mosses of the Anbar Plateau (Subarctic Siberia). – Arctoa 20: 153–174.
- Hedenäs, L. 1989. *Amblystegium longicuspis* Lindb. & H. Arn., its status and taxonomic position. – Lindbergia 14: 142–146.
- Hedenäs, L. 2003. The European species of the *Calliargon-Scorpidium-Drepanocladus* complex, including some related or similar species. – Meylania 28: 1–116.
- Ignatov, M. S. 1994. Bryophytes of Altai Mountains. I. Study area and history of its bryological exploration. – Arctoa 3: 13–27.
- Ignatov, M. S., Afonina, O. M. and Ignatova, E. A. 2006. Check-list of mosses of east Europe and north Asia. – Arctoa 15: 1–130.
- Lindberg, S. O. and Arnell, H. W. 1890. Musci Asiae borealis. – Kungl. Sven. Vetenskapsakad. Handl. 23: 1–163.
- Long, D. G. 1985. Polytrichaceae. – In: Mogensen, G. S. (ed.), Illustrated moss flora of arctic North America and Greenland. 1. – Medd. Grønland Bioscience 17: 9–57.
- Lübenau, R. and Lübenau, K. 1971. *Psilopilum cavifolium* (Wils.) Hagen in der Torne-Lappmark in Schweden. – Herzogia 2: 149–151.
- Martinčić, A. 2009. Contributions to the bryophyte flora of Republic of Macedonia. – Hacquetia 8: 97–114.
- Mårtensson, O. 1956. Bryophytes of the Torneträsk area, Northern Swedish Lappland. II. Musci. – Kungl. Sven. Vetenskapsakad. Avhandl. Naturskyddsärenden 14: 1–321.
- Rumsey, F. J. 1990. Additions to the bryophyte flora of the Torneträsk area, Swedish Lapland. – J. Bryol. 16: 199–208.
- Sabovljevic, M., Cvetic, T. and Stevanovic, V. 2004. Bryophyte Red List of Serbia and Montenegro. – Biodivers. Conserv. 13: 1781–1790.