

## **Redescription of the Central Asian spider *Pholcoides seclusa* comb. nov. (Araneae: Filistatidae)**

Authors: Marusik, Yuri M., and Zonstein, Sergei

Source: Arachnologische Mitteilungen: Arachnology Letters, 57(1) : 43-47

Published By: Arachnologische Gesellschaft e.V.

URL: <https://doi.org/10.30963/aramit5708>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Redescription of the Central Asian spider *Pholcoides seclusa* comb. nov. (Araneae: Filistatidae)

Yuri M. Marusik & Sergei Zonstein



doi: 10.30963/aramit5708

**Abstract.** *Filistata seclusa* O. Pickard-Cambridge, 1885, a species described from Himalayas was thought to be based on immature specimens. The study of one of the syntypes revealed that it is an adult female, which is selected as the lectotype and re-described. Somatic and endogyne morphology indicate that this species belongs to *Pholcoides* Roewer, 1960, a genus known only from two species from Afghanistan (*P. afghana* Roewer, 1960) and adjacent Tajikistan (*P. monticola* (Spassky, 1941)). Therefore, a new combination is proposed for the species: *Pholcoides seclusa* (O. Pickard-Cambridge, 1885) **comb. nov.** Comparative figures are given for all three species as well as a map of distribution of the genus.

**Keywords:** Afghanistan, Aranei, *Filistata*, India, Pakistan, Prithinae, Tajikistan, Xinjiang

**Zusammenfassung. Wiederbeschreibung der zentralasiatischen Spinne *Pholcoides seclusa* comb. nov. (Araneae: Filistatidae).** *Filistata seclusa* O. Pickard-Cambridge, 1885, eine Art die aus dem Himalaya beschrieben wurde, wurde vermeintlich nach immaturren Exemplaren beschrieben. Die Untersuchung eines Syntypus zeigte, dass es ein adultes Weibchen ist, das als Lectotyp ausgewählt und wiederbeschrieben wird. Habitus und Morphologie der Endogyne belegen, dass die Art zur Gattung *Pholcoides* Roewer, 1960 gehört. *Pholcoides* enthält bislang lediglich zwei Arten aus Afghanistan (*P. afghana* Roewer, 1960) und dem benachbarten Tadschikistan (*P. monticola* (Spassky, 1941)). Daher wird für die Art eine neue Kombination vorgeschlagen: *Pholcoides seclusa* (O. Pickard-Cambridge, 1885) **comb. nov.** Vergleichende Abbildungen werden für alle drei Arten vorgelegt sowie eine Verbreitungskarte für die Gattung.

*Pholcoides* Roewer, 1960 is a small genus with only two species (WSC 2018) belonging to Prithinae Gray, 1995 (Zonstein et al. 2013). Both species, *Pholcoides afghana* Roewer, 1960 and *P. monticola* (Spassky, 1941) are known from the mountains of Central Asia, Afghanistan and Tajikistan. While studying types of spiders described by Pickard-Cambridge (1885) based on material of the Second Yarkand Mission (deposited in Oxford University Museum of Natural History) we found one specimen belonging to Filistatidae. Although vials with the types from the Second Yarkand Mission lack any geographical or species name labels (Prószyński & Żochowska 1981, Marusik 1993, 2017, Marusik et al. 2018, Marusik & Nadolny 2018) it was easy to recognize that the filistatid specimen belongs to *Filistata seclusa* O. Pickard-Cambridge, 1885, the only species of the family considered in O. Pickard-Cambridge (1885). Because this species was never properly figured, and its endogyne was never described we decided to redescribe this species. Study of the somatic morphology reveals that the species is misplaced in the genus and even in the subfamily, and is very similar to *Pholcoides*. Examination of the endogyne of this species proved that it belongs to *Pholcoides*. The goals of this paper are as follows: providing a detailed redescription of *Filistata seclusa*, establishing a new combination, comparing *P. seclusa* with two other congeners and rediagnosing the genus.

### Material and methods

Specimens were photographed with a Canon EOS 7D camera attached to an Olympus SZX16 stereomicroscope and Pro-Microscan camera attached to an Olympus BH-2 stereomicroscope. Digital images were mounted using CombineZP and Helicon Focus 3.10 image stacking software. Endogynes were cleared in a KOH/water solution until the soft tissues were dissolved. Standard abbreviations are used for leg seg-

ments: Fe – femur, Pa – patella, Ti – tibia, Mt – metatarsus, Ta – tarsus. Further abbreviations: Pg – pore gland, Rc – receptacle proper, Rs – stalk of receptacle, Vs – vesicle. All measurements are in mm. The lectotype of *Pholcoides seclusa* is from the Oxford University Museum of Natural History, United Kingdom (OUMNH).

### Taxonomic survey

#### *Pholcoides* Roewer, 1960

*Pholcoides* Roewer, 1960: 40.

*Pholcoides*: Huber 2009: 68 (transferred from Pholcidae to Filistatidae).

*Pholcoides*: Zonstein et al. 2013: 65.

*Pholcoides*: Zonstein & Marusik 2016: 86.

**Type species.** *Pholcoides afghana* Roewer, 1960 from Afghanistan.


**Comment.** So far, the genus is known by females only.

**Diagnosis.** Within the Palearctic Filistatidae *Pholcoides* is unique in possessing the combination of a medium-sized compact body with carapace and sternum circular or even nearly wider than long, the spinnerets and anal tubercle located posteriorly and closer to the opisthosoma apex (Figs 3, 8), and rather long and slender female pedipalps and legs I–IV (Figs 1, 3, 7, unusual for the Palearctic Filistatidae). Females (males are not known) can be distinguished from other Palearctic genera by the long receptacles with long membranous bases (stems) and lobulated terminal parts. Long and lobulated receptacles are unique for Filistatidae.


**Description.** Small to medium-sized spider, with carapace 1.2–1.7 mm long. Carapace flattened circular, with or without dark (partially reticulate) pattern on clypeus, postocular area and margins. Eye tubercle low, blackened. Clypeus moderately long, subhorizontal. Sternum circular; labium much longer than wide; maxillar lobes long weakly converging. Legs long and slender. Calamistrum very short, formed by two rows of setae. Opisthosoma elongate with spinneret group located relatively close to opisthosoma apex. Male characters unknown. Endogynes in all known species are rather similar and composed by pair of undivided receptacles. Receptacle has

Yuri M. MARUSIK, Institute for Biological Problems of the North RAS, Portovaya Str. 18, Magadan, Russia; E-Mail: yurmar@mail.ru; Department of Zoology & Entomology, University of the Free State, Bloemfontein 9300, South Africa  
Sergei ZONSTEIN, Steinhardt Museum of Natural History, Tel-Aviv University, 69978 Tel-Aviv, Israel; E-mail: znn@post.tau.ac.il

submitted 16.8.2018, accepted 17.2.2019, online 25.3.2019



**Figs 1-10:** Habitus and somatic characters of females of *Pholcoides seclusa* (1-6), *P. monticola* (7-9) and *P. afghana* (10). **1, 7,** habitus, dorsal; **2, 8,** opisthosoma, ventral; **3,** habitus, lateral; **4-5,** prosoma, dorsal and frontal; **6, 9-10,** metatarsus IV showing calamistrum dorsal. 7-9 after Zonstein & Marusik (2016), 10 after Zonstein et al. (2013)



**Figs 11-17:** Endogyne of *Pholcoides seclusa* (11-12), *P. monticola* (13-14) and *P. afghana* (15-17). **11-13, 17,** dorsal; **14,** anterior; **15,** ventral; **16,** dorso-anterior. 13-14 after Zonstein & Marusik (2016), 15-17 after Zonstein et al. (2013). Abbreviations: Pg – pore gland, Rc – receptacle proper, Rs – stalk of receptacle, Vs – vesicle

wide weakly sclerotized stem (Rs), longer than wide, roundly bent, proximally converging and in mid part strongly or weakly diverging. Receptacles proper (Rc) sclerotized stronger than stems; in all three species complex, composed by several lobes (Vs) of different size; basal proximal nodules smaller than distal. Distinct pore glands absent. *Pholcoides seclusa* have pore glands in the basal part of the proper receptacle (cf. Fig. 11).

**Composition.** The genus was described as monotypic. Until recently it was known by two species, *P. afghana* Roewer, 1960 and *P. monticola* (Spassky, 1941). Together with *P. seclusa* (O. Pickard-Cambridge, 1885) it now encompasses three species.

***Pholcoides afghana* Roewer, 1960** (Figs 10, 15-18)

*Pholcoides afghana* Roewer, 1960: 40, figs 12 a-d (♀).

*Pholcoides afghana*: Zonstein et al. 2013: 65, figs 1-12 (♀).

**Comments.** The species was redescribed and the endogyne was figured in Zonstein et al. (2013). It is known from the type locality only – Kabul.

***Pholcoides monticola* (Spassky, 1941)** (Figs 7-9, 13-14, 18)

*Filistata monticola* Spassky, 1941: 12 (♀).

*Zaitunia monticola*: Zonstein 1990: 50 (transferred from *Filistata*).

*Pholcoides monticola*: Zonstein & Marusik 2016: 86, figs 45A-E (transferred from *Zaitunia*).

**Comments.** The species was illustrated in Zonstein & Marusik (2016). It is known from two localities in eastern Tajikistan (Zonstein & Marusik 2016).

***Pholcoides seclusa* (O. Pickard-Cambridge, 1885) comb. nov.** (Figs 1-6, 11-12, 18)

*Filistata seclusa* O. Pickard-Cambridge, 1885: 5, pl. 1, fig. 2 (immature ♀).

*Filistata seclusa*: Reimoser 1919: 189 (indicated that ♀ was described by O. Pickard-Cambridge).

*Filistata seclusa*: Caporiacco 1934: 116, pl. 1, fig. 2 (reported from several localities: ♀♀, juvenile ♀♀ and ♂♂ in Karakorum and provided figure of eyes in ♀, probably misidentified).


*Filistata reclusa*: Reimoser 1935: 170 (misspelling; reported ♀ from Kashgar, Xinjiang China, most likely misidentified).

*Filistata seclusa*: Roewer 1955: 1279 (distribution indicated as “Turkestan, Karakorum”, that ♀ was described by O. Pickard-Cambridge, and that Caporiacco (1934) had described both sexes).

**Material examined.** Lectotype ♀ designated here, label in the vial: “B[ottle] 378 v[ial] 4 1 sp”. Information in the text: “Leh, August or September 1873; Pankong-valley, 15th to 21st September 1873”. Paralectotype specimens, either subadult females or adult females not found.

**Note.** The World Spider Catalog (2018) states that the species was described based on a female, although O. Pickard-Cambridge (1885) mentioned in his description an “immature female” only. The WSC (2018) indicates that Caporiacco (1934) described a male of *P. seclusa*, although males (seem to be juvenile) are listed among the material examined, but not mentioned in the text and not figured.

**Diagnosis.** This species differs from congeners by its smaller size (carapace 1.26 long, vs. 1.74 and 1.6 in two other



**Fig. 18:** Distribution records of *Pholcoides* species: *P. afghana* (dot), *P. monticola* (triangle) and *P. seclusa* (black squares refer to localities of syntypes and blue diamonds refer to records by Caporiacco (1934) and Reimoser (1935) which are most likely based on misidentifications)

species), and in having a distinct carapace pattern with dark marginal stripes (vs. no marginal stripes). The endogyne of *P. seclusa* is most similar to those of *P. afghana* by having receptacles with two large lobes. *Pholcoides seclusa* can be distinguished from sibling species by the almost straight stalks of its receptacles (vs. roundly bent in *P. afghana*) separated by more than one radius (vs. separated by less than one radius). The endogyne of *P. seclusa* differs markedly from *P. monticola* by the almost straight and less spaced stalks (vs. bent stalks separated by more than their diameter) and larger vesicles.

### Description

**Measurements.** Body 3.63 long; carapace 1.26 long, 1.28 wide, 0.4 high; opisthosoma 2.5 long, 1.68 wide. Eye sizes and interdistances: AME 0.09, ALE 0.14, PME 0.1, PLE 0.14; ALE–ALE 0.14, PME–PME 0.1. Clypeus on ALE 0.17, on AME 0.27. Palp and leg length as in Tab. 1.

**Somatic characters.** Carapace brownish yellow with brown median patch and marginal stripes, anteriorly fused with yellow spots; eye tubercle dark brown. Opisthosoma dark greyish yellow with median brownish violet stripe margined with narrow brown outer stripes. Ventral surface and legs (concolorous) slightly paler than carapace. Ventral opisthosoma, otherwise, slightly darker. Spines completely lost but bases for 1 dorsobasal spine on all femora visible, as well as at least 3 retrolateral and 1 prolateral and several ventral spines on tibia IV. Other legs also with spines.

**Tab. 1:** Palp and leg lengths

	Fe	Pt	Ti	Mt	Ta	Total
<b>Palp</b>	0.85	0.45	0.6	–	0.73	2.63
<b>I</b>	1.55	0.53	1.45	1.28	0.88	5.69
<b>II</b>	1.38	0.55	1.2	1.13	0.7	4.96
<b>III</b>	1.33	0.55	1.2	1.25	0.7	5.03
<b>IV</b>	1.63	0.55	1.38	1.63	0.83	6.02

**Endogyne.** Stalks of receptacles almost straight, spaced by more than one radius; receptacles diverging, with two large lobes and a few smaller ones; basal part of receptacles with poorly distinct pore glands.

**Distribution and comments.** The World Spider Catalog (2018) indicated the distribution of this species as China (Yarkand), Karakorum, and the Himalayas. It was described based on specimens from eastern part of the Jammu and Kashmir State in India and one locality lying on the border between Jammu and Kashmir and Xizang (Tibet), China. The species was reported from SW Xinjiang (Reimoser 1935) and several localities in northern Jammu and Kashmir (Caporiacco 1934) now belonging to Pakistan. Specimens from Karakoram and Xinjiang were identified based on the brief description and habitus figures given by Pickard-Cambridge (1885). Given that most filistatid species have a very local distribution in the Palearctic (cf. Zonstein & Marusik 2016, Zamani & Marusik 2018) it is very likely that the record from Xinjiang refers to another species, and possibly even to another genus. Records by Caporiacco (1934) from Karakoram may also belong to other species, as the distance between the type locality and records by Caporiacco is over 200 km, and the two localities belong to different mountain ranges.

### Acknowledgements

We thank Zoë Simmons (Oxford, UK) and Dmitri V. Logunov (Manchester, UK) for their help in obtaining the type of *Pholcoides seclusa*. Special thanks go to Seppo Koponen (ZMUT) for providing us with museum facilities. This study was supported in part by the Ministry of Absorption, Israel.

### References

Caporiacco L di 1934 Aracnidi dell'Himalaia e del Karakoram raccolti dalla Missione Italiana al Karakoram (1929-VII). – Memorie della Società Entomologica Italiana, Genova 13: 113-160

- Huber BA 2009 Four new generic and 14 new specific synonymies in Pholcidae, and transfer of *Pholcoides* Roewer to Filistatidae (Araneae). – *Zootaxa* 1970: 64–68 – doi: [10.11646/zootaxa.3847.3.5](https://doi.org/10.11646/zootaxa.3847.3.5)
- Marusik YM 1993 Re-description of spiders of the families Heteropodidae and Thomisidae (Aranei), described by O. P.-Cambridge from the material of the second Yarkand mission. – *Entomologicheskoye Obozrenie* 72: 456–468 [in Russian, English abstract]
- Marusik YM 2017 Redescription of *Dysdera cylindrica* (Aranei: Dysderidae) a single known species of the family in Pakistan with notes on the distribution of the family. – *Arthropoda Selecta* 26: 310–314
- Marusik YM & Nadolny AA 2018 Redescription of enigmatic spider genus *Stoliczka* O. Pickard-Cambridge, 1885 (Araneae: Lycosoidea) based on the type species. – *Zootaxa* 4497: 295–300 – doi: [10.11646/zootaxa.4497.2.9](https://doi.org/10.11646/zootaxa.4497.2.9)
- Marusik YM, Omelko MM & Koponen S 2018 Redescription of Himalayan *Trachelas costatus* (Araneae: Trachelidae). – *Zootaxa* 4433: 390–392 – doi: [10.11646/zootaxa.4433.2.11](https://doi.org/10.11646/zootaxa.4433.2.11)
- Pickard-Cambridge O 1885 Araneidea. In: Scientific results of the second Yarkand mission; based upon the collections and notes of the late Ferdinand Stoliczka, Ph. D. Government of India, Calcutta. 115 pp., 2 pls – doi: [10.5962/bhl.title.119960](https://doi.org/10.5962/bhl.title.119960)
- Prószyński J & Źochowska K 1981 Redescriptions of the O. P.-Cambridge Salticidae (Araneae) types from Yarkand, China. – *Polskie Pismo Entomologiczne* 51: 13–35
- Reimoser E 1919 Katalog der echten Spinnen (Araneae) des Paläarktischen Gebietes. – *Abhandlungen der Zoologisch-Botanischen Gesellschaft in Wien* 10 (2): 1–280
- Reimoser E 1935 Araneida. In: *Wissenschaftliche Ergebnisse der niederländischen Expeditionen in den Karakorum*, Zoologie. Leipzig, pp. 169–176
- Roewer CF 1955 Katalog der Araneae von 1758 bis 1940, bzw. 1954. 2. Band, Abt. a (Lycosaeformia, Dionycha [excl. Salticiformia]). 2. Band, Abt. b (Salticiformia, Cribellata) (Synonyma-Verzeichnis, Gesamtindex). Institut royal des Sciences naturelles de Belgique, Bruxelles. 1751 pp.
- Roewer CF 1960 Solifugen und Opilioniden, Araneae, Orthognathae, Haplogynae und Entelegynae (Contribution à l'étude de la faune d'Afghanistan, 23). – *Göteborgs Kungliga Vetenskaps- och Vitterhets-Samhälles handlingar B* 8 (7): 1–53
- Spassky S 1941 Araneae palaearticae novae VI. – *Folia Zoologica et Hydrobiologica*, Riga 11: 12–26
- World Spider Catalog 2018 World Spider Catalog. Version 19.5. – Internet: <http://wsc.nmbe.ch> (8. Aug. 2018) – doi: [10.24436/2](https://doi.org/10.24436/2)
- Zamani A & Marusik YM 2018 New species and records of Filistatidae (Arachnida: Aranei) from Iran. – *Arthropoda Selecta* 27: 121–128
- Zonstein S & Marusik YM 2016 A revision of the spider genus *Zaitunia* (Araneae, Filistatidae). – *European Journal of Taxonomy* 214: 1–97 – doi: [10.5852/ejt.2016.214](https://doi.org/10.5852/ejt.2016.214)
- Zonstein SL 1990 [A synopsis of species of the spider family Filistatidae (Aranei) of the USSR with description of a new genus and a new species from western Tien-Shan]. – *Zoologicheskii Zhurnal* 69(10): 50–53 [in Russian, English abstract]