

A new Agyneta from Italy and Greece (Araneae: Linyphiidae)

Author: Tanasevitch, Andrei V.

Source: Arachnologische Mitteilungen: Arachnology Letters, 57(1) : 87-88

Published By: Arachnologische Gesellschaft e.V.

URL: <https://doi.org/10.30963/aramit5716>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A new *Agyneta* from Italy and Greece (Araneae: Linyphiidae)

Andrei V. Tanasevitch



doi: 10.30963/aramit5716

Abstract. A new species, *Agyneta inermis* **spec. nov.**, is described, based on two males from Italy and Greece. The new species is very similar to the Iranian *A. iranica* Tanasevitch, 2011 (first record for Turkmenistan), but differs in structural details of the male palp, primarily by the absence of a tooth at the base of the embolus.

Keywords: *Agyneta inermis*, *Agyneta iranica*, Europe, Iran, Mediterranean, Micronetinae, new species, spiders, taxonomy, Turkmenistan

Zusammenfassung. Eine neue *Agyneta* aus Italien und Griechenland (Araneae: Linyphiidae). Eine neue Art, *Agyneta inermis* **spec. nov.**, wird auf Basis zweier Männchen aus Italien und Griechenland beschrieben. Die neue Art ist der iranischen *A. iranica* Tanasevitch, 2011 (Erstnachweis für Turkmenistan) sehr ähnlich, unterscheidet sich aber in Details des männlichen Palpus, besonders durch das Fehlen eines Zahns an der Embolus-Basis.

Examination of the extensive spider material collected by the Swiss arachnologist Antoine Senglet in Italy and Greece revealed a new species of *Agyneta* Hull, 1911 described here in the present paper.

Material and methods

This paper is based on material kept at the Muséum d'histoire naturelle de Genève, Switzerland (MHNG). Specimens preserved in 70% ethanol were studied using a MBS-9 stereomicroscope. The sequence of leg segment measurements is as follows: femur + patella + tibia + metatarsus + tarsus. All measurements are given in mm. The terminology of the copulatory organs mainly follows that of Merrett (1963) and Saaristo (1971, 1973).

Abbreviations used in the text and figures

E = embolus, EP = embolus proper (after Saaristo 1971), L = lamella characteristica, Mt = metatarsus, PH = pit hook (after Saaristo 1973), R = radix, T = tooth, TmI = position of metatarsal trichobothrium on leg I, ZMMU = Zoological Museum of the Moscow State University, Moscow, Russia.

Results

Taxonomy

Family Linyphiidae Blackwall, 1859

Subfamily Micronetinae Hull, 1920

Genus *Agyneta* Hull, 1911

Type species. *Neriene subtilis* O. Pickard-Cambridge, 1863, by original designation.

Agyneta inermis **spec. nov.** (Figs 1-10)

Material. Holotype: male (MHNG), ITALY, Calabria, Province of Cosenza, Tarsia, valley of Crati River (ca. 39.609°N, 16.273°E), 50 m a.s.l., 4.–5. Aug. 1968, leg. A. Senglet. Paratype: GREECE: 1 male (MHNG), Halkidiki, Nea Kallikratis (ca. 40.313°N, 23.063°E), 10 m a.s.l., 15. Jun. 1970, leg. A. Senglet.

Comparative material examined. Male holotype of *Agyneta iranica* Tanasevitch, 2011 (MHNG), IRAN, Golestan, be-

tween Naharkhoran and Gorgan (ca. 36.766°N, 54.471°E), 500–600 m a.s.l., forest, sifted litter and moss, 20. Jul. 1973, leg. A. Senglet. TURKMENISTAN, 1 male (ZMMU), Kopetdagh Mts, Firjuza (ca. 37.915°N, 58.089°E), 600–700 m a.s.l., 6.–17. Feb. 1979, leg. S. Kuznetsov. New record for the country.

Diagnosis. *Agyneta inermis* **spec. nov.** is very similar to *A. iranica* Tanasevitch, 2011, known from Iran (Tanasevitch 2011) and Turkmenistan (new data). The new species is distinguished by certain details of the structure of the lamella characteristica (Figs 7-10 vs. Fig. 11), by the shallow hollow on the apex of the palpal tibia (Fig. 3 vs. Fig. 13), as well as by the absence of a basal tooth at the base of the embolus (Fig. 6 vs. Fig. 12, tooth marked as “T”).

Etymology. The specific epithet is a Latin adjective meaning “unarmed”, referring to the absence of a tooth at the base of the embolus.

Description. Male (holotype). Total length 1.80. Pro-soma 0.75 long, 0.55 wide, brown with dark margins. Chelicerae 0.33 long, weak. Legs pale brown. Leg I 3.07 long (0.80+0.23+0.78+0.73+0.53); IV 2.88 long (0.78+0.20+0.75+0.70+0.45). Opisthosoma 0.93 long, 0.60 wide, dark grey. Chaetotaxy: each tibia with two dorsal spines; Mt I–IV spineless. Metatarsus IV without a trichobothrium. TmI 0.31. Palp (Figs 1-10): tibia short, with a small, pointed, dorso-prolateral tubercle and a short, wide, keel-shaped, retrolateral outgrowth. Cymbium without posterodorsal apophysis. Paracymbium with well-developed pockets, except for apical one (see Saaristo 1973). Lamella characteristica a narrow band expanded at base, uncinately apically. Embolus without teeth at its base.

Female. Unknown.

Variation. As the palps in both type specimens seem to be identical, no variation in this structure has been found.


Distribution. The species is known from only two localities, one each in Italy and Greece.

Acknowledgements

I am deeply grateful to Peter Schwendinger (MHNG) and Kirill Mikhailov (ZMMU) for the opportunity to work on the spider collections under their care, as well as to Sergei Golovatch (Moscow, Russia) for checking the English of an advanced draft. I also thank Nadine Dupérré (Hamburg, Germany) and Robert Bosmans (Gent, Belgium) who kindly reviewed the manuscript.

Andrei V. TANASEVITCH, A. N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Leninsky prospekt 33, Moscow 119071, Russia; E-mail: tanasevitch@gmail.com; ORCID: 0000-0002-9116-606X

submitted 29.1.2019, accepted 8.3.2019, online 25.3.2019



Figs 1-13: Palpal structure of *Agyneta inermis* sp. nov., holotype (1-10), and *A. iranica* Tanasevitch, 2011, holotype (11-13) after Tanasevitch (2011). **1.** right palp, retrolateral view; **2.** cymbium, prolateral view; **3-4, 13.** palpal tibia, (3, 13) dorsal view, (4) dorso-prolateral view; **5.** palpal tibia and paracymbium, retrolateral view; **6.** embolus and lamella characteristic, ventral view; **7-11.** lamella characteristic, different angles from ventro-retrolateral to dorso-retrolateral view, respectively, (10) and (11) nearly at the same position; **12.** embolic division, ventral view. Scale bars: 0.1 mm

References

- Merrett P 1963 The palpus of male spiders of the family Linyphiidae. – Proceedings of the Zoological Society of London 140: 347-467 – doi: [10.1111/j.1469-7998.1963.tb01867.x](https://doi.org/10.1111/j.1469-7998.1963.tb01867.x)
 Saaristo MI 1971 Revision of the genus *Maro* O. P.-Cambridge (Araneae, Linyphiidae). – Annales Zoologici Fennici 8: 463-482

- Saaristo MI 1973 Taxonomical analysis of the type-species of *Agyneta*, *Anomalaria*, *Meioneta*, *Aprolagus*, and *Syedrella* (Araneae, Linyphiidae). – Annales Zoologici Fennici 10: 451-466
 Tanasevitch AV 2011 A new *Agyneta* Hull, 1911 from Iran (Araneae, Linyphiidae, Micronetinae). – Revue suisse de Zoologie 118: 45-48 – doi: [10.5962/bhl.part.117798](https://doi.org/10.5962/bhl.part.117798)