

## **First record of the genus *Thyreosthenius* (Araneae: Linyphiidae) from the Iberian Peninsula with notes on the host preference of the myrmecophilic *T. biovatus***

Authors: Castellucci, Filippo, Schifani, Enrico, García, Fede, Luchetti, Andrea, and Scharff, Nikolaj

Source: Arachnologische Mitteilungen: Arachnology Letters, 65(1) : 27-30

Published By: Arachnologische Gesellschaft e.V.

URL: <https://doi.org/10.30963/aramit6506>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# First record of the genus *Thyreosthenius* (Araneae: Linyphiidae) from the Iberian Peninsula with notes on the host preference of the myrmecophilic *T. biovatus*

Filippo Castellucci, Enrico Schifani, Fede García, Andrea Luchetti & Nikolaj Scharff



doi: 10.30963/aramit6506

**Abstract.** The presence of the spider genus *Thyreosthenius* Simon, 1884 (Linyphiidae) is for the first time confirmed in the Iberian Peninsula with an observation of the myrmecophile species *T. biovatus* (O. Pickard-Cambridge, 1875). Four adult females and two sub-adult male specimens were collected inside mound nests belonging to the red wood ant species *Formica lugubris* Zetterstedt, 1838 in the Aragon and Catalonia regions of Spain. This represents the second known record of an occurrence of *T. biovatus* in association with *F. lugubris*, and the first for Continental Europe. In addition, the existing information on the host preference of this species is reviewed and discussed in relation to the new data from the Spanish Pyrenees.

**Keywords:** ant association, *Formica lugubris*, myrmecophily, new record, RWA, Spain, spider

**Zusammenfassung. Erstnachweis der Gattung *Thyreosthenius* (Araneae: Linyphiidae) von der Iberischen Halbinsel mit Bemerkungen zur Wirtspräferenz der myrmekophilen Art *T. biovatus*.** Die Präsenz der Spinnengattung *Thyreosthenius* Simon, 1884 (Linyphiidae) auf der Iberischen Halbinsel wird durch eine Beobachtung der myrmekophilen Art *T. biovatus* (O. Pickard-Cambridge, 1875) bestätigt. Vier adulte Weibchen und zwei subadulte männliche Exemplare wurden in Hügelnestern der Waldameisenart *Formica lugubris* Zetterstedt, 1838 im spanischen Aragonien und Katalonien gesammelt. Dies stellt den zweiten bekannten Nachweis einer Vergesellschaftung von *T. biovatus* mit *F. lugubris* dar, sowie den ersten für Kontinentaleuropa. Zusätzlich werden die vorhandenen Informationen zur Wirtspräferenz dieser Art gesichtet sowie in Zusammenhang mit den neuen Daten aus den spanischen Pyrenäen diskutiert.

The Iberian Peninsula hosts an extremely diverse spider fauna, with 1493 recorded species, 282 of which are endemic (Branco et al. 2019). Continental Spain, with 1386 species, represents one of the most spider-rich areas in Europe, ranking third after Italy and France (Nentwig et al. 2023).

Myrmecophily is defined as the ability to live in a close association with ants, from foraging alongside them to spending the whole life cycle inside ant nests (Wasmann 1894, Donisthorpe 1927, Hölldobler & Wilson 1990). Cushing (1997, 2012) reviewed this phenomenon in spiders and reported this lifestyle in 13 different spider families. Linyphiidae, the second largest family in spiders (World Spider Catalog 2023), holds the highest number of myrmecophilic taxa, with ten species from nine genera (Cushing 1997, 2012) (Tab. 1). The linyphiid genus *Thyreosthenius* Simon, 1884 includes two species, the free-living Holarctic *Thyreosthenius parasiticus* (Westring, 1851) and the myrmecophilic Palearctic *Thyreosthenius biovatus* (O. Pickard-Cambridge, 1875) (World Spider Catalog 2023). In Europe, *T. biovatus* is widespread in northern and central countries, while it has never been recorded from Iberia, the Balkans (excluding Bulgaria), Belarus, and Lithuania (Nentwig et al. 2023). This species lives inside the nests of ants belonging to the genus *Formica* Linnaeus, 1758, including mound building red wood ant species (RWA) belonging to the *Formica rufa* group, such as *Formica aquilonia* Yarrow, 1955, *Formica lugubris* Zetterstedt, 1838, *For-*

*mica polyclena* Foerster, 1850, *Formica pratensis* Retzius, 1783 and *Formica rufa* Linnaeus, 1761, and the non-RWA species *Formica fusca* Linnaeus, 1758 and *Formica sanguinea* Latreille, 1798 (Bösenberg 1899, Simon 1926, Bristowe 1939, Wiehle 1960, Palmgren 1976, Robinson 1998, Parmentier et al. 2014, 2015, 2020, Castellucci et al. 2022). Due to the low amount of cuticular hydrocarbons (CHCs) registered for *T. biovatus* (Parmentier et al. 2017), it is likely that its integration within the colonies happens by means of chemical insignificance (Lenoir et al. 2001, 2013, Witte et al. 2008).

Most of the host association data available for *T. biovatus* originate from Central and Northern European countries, while little is known about its ecology at the southern limits of its distribution range. Only recently, Castellucci et al. (2022) shed light on its host preferences in the Italian Alps, recording associations with three different RWA species, including the first observation of *F. aquilonia* as a host. The only available information about this species in the Pyrenees comes from Simon (1926), who reported it living in the nests of *Formica fusca* on the French side of the mountain range. In this study, we report about the first record of the genus *Thyreosthenius* for the Iberian Peninsula, based on new data from the Spanish Pyrenees.

## Materials and Methods

Fieldwork was conducted in 2017 and 2021 in the surroundings of Benasque, Huesca, and Toses, Girona, in the Spanish Pyrenees. A couple of liters of material were collected from the central part of RWA mound nests and sifted using an entomological sieve with an 8 × 8 mm mesh on a white fabric sheet, to facilitate the observation of myrmecophile spiders. These, and RWA workers, were then hand collected and stored in 70% or 96% ethanol. After the sifting phase, residual nest material and worker ants were carefully placed back on the mound to minimize disturbance. Spiders were examined with a stereoscopic microscope and photographed using a BK+ Imaging System from Visionary Digital equipped with a Canon EOS 7D camera. Identification was carried out following the keys by Roberts (1987). Ants were examined with a stereoscopic microscope and photographed with a Canon

Filippo CASTELLUCCI, Department of Biological, Geological and Environmental Sciences, University of Bologna, via Selmi 3, 40126 Bologna, Italy; Zoology Section, Natural History Museum of Denmark, University of Copenhagen, Universitetsparken 15, 2100, Copenhagen, Denmark; E-mail: filippo.castellucci2@unibo.it, ORCID: <https://orcid.org/0000-0002-9944-2196>

Enrico SCHIFANI, Department of Chemistry, Life Sciences & Environmental Sustainability, University of Parma, parco Area delle Scienze 11/a, 43124 Parma, Italy; E-mail: enrico.schifani@unipr.it, ORCID: <https://orcid.org/0000-0003-0684-6229>

Fede GARCÍA, Blesa, 45, 08004 Barcelona, Spain; E-mail: chousas2@gmail.com

Andrea LUCHETTI, Department of Biological, Geological and Environmental Sciences, Alma Mater Studiorum Università di Bologna, via Selmi 3, 40126 Bologna, Italy; E-mail: andrea.luchetti@unibo.it, ORCID: <https://orcid.org/0000-0002-2986-721X>

Nikolaj SCHARFF, Zoology Section, Natural History Museum of Denmark, University of Copenhagen, Universitetsparken 15, 2100 Copenhagen, Denmark; E-mail: nscharff@snm.ku.dk, <https://orcid.org/0000-0001-6809-2878>

Academic editor: Konrad Wiśniewski

submitted 18.2.2023, accepted 19.6.2023, online 4.8.2023

**Tab. 1:** List of myrmecophile Linyphiidae species, with information on their known host ant species

| Species                                                         | Host ant species                                                                                                                                                                                                                                             | References                                                                  |
|-----------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|
| <i>Acartauchenius scurrilis</i><br>(O. Pickard-Cambridge, 1873) | <i>Formica rufa</i> Linnaeus, 1761, <i>Lasius flavus</i> (Fabricius, 1782),<br><i>Tetramorium caespitum</i> (Linnaeus, 1758)                                                                                                                                 | Cushing (1997)                                                              |
| <i>Diastanillus pecuarius</i> (Simon, 1884)                     | <i>Formica</i> cf. <i>fusca</i> (Linnaeus, 1758), <i>F. lemani</i> Bondroit, 1917                                                                                                                                                                            | Cushing (2012)                                                              |
| <i>Evansia merens</i><br>O. Pickard-Cambridge, 1901             | <i>Formica cunicularia</i> Latreille, 1798, <i>F. fusca</i> (Linnaeus, 1758),<br><i>F. sanguinea</i> Latreille, 1798, <i>Lasius niger</i> (Linnaeus, 1758)                                                                                                   | Cushing (1997)                                                              |
| <i>Grammonota pictilis</i><br>(O. Pickard-Cambridge, 1875)      | <i>Atta texana</i> (Buckley, 1860)                                                                                                                                                                                                                           | Cushing (1997)                                                              |
| <i>Masoncus pogonophilus</i><br>Cushing, 1995                   | <i>Pogonomyrmex badius</i> (Latreille, 1802)                                                                                                                                                                                                                 | Cushing (1997)                                                              |
| <i>Masoncus</i> sp.                                             | <i>Atta texana</i> (Buckley, 1860)                                                                                                                                                                                                                           | Cushing (1997)                                                              |
| <i>Pseudomaro aenigmaticus</i> Denis, 1966                      | <i>Lasius flavus</i> (Fabricius, 1782)                                                                                                                                                                                                                       | Cushing (2012)                                                              |
| <i>Thyreosthenius biovatus</i><br>(O. Pickard-Cambridge, 1875)  | <i>Formica aquilonia</i> Yarrow, 1955, <i>F. fusca</i> (Linnaeus, 1758), <i>F. lugubris</i><br>Zetterstedt, 1838, <i>F. polycтена</i> Foerster 1850, <i>F. pratensis</i> Retzius 1783,<br><i>F. rufa</i> Linnaeus, 1761, <i>F. sanguinea</i> Latreille, 1798 | Cushing (1997), Par-<br>mentier et al. (2020),<br>Castellucci et al. (2022) |
| <i>Scotinotylus formicarius</i><br>(Dondale & Redner, 1972)     | <i>Formica obscuripes</i> Forel, 1886                                                                                                                                                                                                                        | Cushing (1997)                                                              |
| <i>Syedra myrmicarum</i> (Kulczyński, 1882)                     | <i>Manica rubida</i> (Latreille, 1802), <i>Formica</i> sp.                                                                                                                                                                                                   | Cushing (2012)                                                              |

MP-E 65mm f/2.8 1–5 × macro lens mounted on a Canon 1300D camera. Measurements were acquired from photos using the software ImageJ (Schneider et al. 2012). Identification followed the key provided by Seifert (2021).

Results

New record for Spain

*Thyreosthenius biovatus* (O. Pickard-Cambridge, 1875)  
(Figs 1, 2)

For a complete list of taxonomic references see the World Spider Catalog (2023).

**Examined material.** SPAIN: 1 ♀, Benasque, Huesca, Aragon, 42.6126°N, 0.5572°E, 1927 m a.s.l., 14. Jul. 2021, inside *F. lugubris* mound nest, F. Castellucci leg.; 3 ♀♀, 1 sub-adult ♂, Benasque, Huesca, Aragon, 42.6135°N, 0.5570°E, 1912 m a.s.l., 14. Jul. 2021, inside *F. lugubris* mound nest, F. Castellucci leg; 1 sub-adult ♂, Plans de Querol, Toses, Girona, Cataluña,


42.3123°N, 2.0310°E, 1680 m a.s.l., 13. Oct. 2017, inside *F. lugubris* mound nest, F. García leg.

**Distribution.** Europe, Russia (Europe to Far North East) (World Spider Catalog 2023). First record for Spain and the Iberian Peninsula.

**Ecology.** Spiders were collected inside mound nests belonging to the RWA species *Formica lugubris*, between 1680 and 1927 m a.s.l. in a montane pine forest (Fig. 3).


Discussion

*Thyreosthenius biovatus* was already known to occur in the French Pyrenees (Simon 1926). Its presence in Spain probably went unnoticed due to its myrmecophile lifestyle and the lack of focused searches carried out with the specific methods needed for inventorying ant-associates. These peculiar taxa are often neglected during classic biodiversity inventories based on the use of traditional sampling techniques such as pitfall



**Fig. 1:** *Thyreosthenius biovatus* female, habitus. **a.** dorsal view; **b.** ventral view





**Fig. 2:** *Thyreosthenius biovatus* female, epigyne

trapping, sweep netting, or the use of beating trays. Moreover, the number of studies focusing on myrmecophile spiders in Southern Europe is extremely low when compared to Central and Northern countries (Castellucci et al. 2022, Lenzini et al. 2022). Given the known presence of some of its host ant species in other regions of Spain and Portugal (Seifert 2021), it is likely that the actual distribution of *T. biovatus* in the Iberian Peninsula extends further away from the Pyrenees, and that this species is more widespread and abundant than previously thought, as observed in other countries like Belgium, Britain, Denmark and Italy after targeted searches were conducted (Donisthorpe 1927, Scharff & Gudik-Sørensen 2006, Parmentier et al. 2014, Castellucci et al. 2022). Our findings of *T. biovatus* in association with *F. lugubris* represent the only observations of co-occurrence between these two species outside the United Kingdom, where two adult males were collected in two different *F. lugubris* nests by Robinson (1998). Little is known about the host preference of this spider in the Pyrenees, with only Simon (1926) reporting its association with the non-RWA species *F. fusca* on the French side of the range. The first record of *T. biovatus* from the Iberian Peninsula suggests that the study of myrmecophile taxa in Southern Europe

has long been neglected and that studies of the cryptic, but fascinating, microhabitat that ant nests represent for spiders and other invertebrate taxa may reveal many new discoveries.

### Acknowledgements

The authors are grateful to Miquel Arnedo for his suggestions in identifying potential sampling areas. The authors are also grateful to Marc Doménec and to the second anonymous reviewer for their comments and suggestions that helped improve the quality of the manuscript. This work has been supported by Canziani funding to AL; the PhD grant to FC was co-funded by Canziani and by the Natural History Museum of Denmark.

### References

- Bösenberg W 1899 Die Spinnen der Rheinprovinz. – Verhandlungen des Naturhistorischen Vereins der Preussischen Rheinlande und Westfalens 56: 69–131
- Branco VV, Morán E & Cardoso P 2019 An update to the Iberian spider checklist (Araneae). – Zootaxa 4614: 201–254 – doi: [10.11646/zootaxa.4614.2.1](https://doi.org/10.11646/zootaxa.4614.2.1)
- Bristowe WS 1939 The comity of spiders vol. 1. Ray Society, London. 228 pp.
- Castellucci F, Schifani E, Luchetti A & Scharff N 2022 New association between red wood ant species (*Formica rufa* group) and the myrmecophilic spiders *Mastigusa arietina* and *Thyreosthenius biovatus*. – Bulletin of Insectology 75: 231–238
- Cushing PE 1997 Myrmecomorphy and myrmecophily in spiders: a review. – Florida Entomologist 80: 165–193 – doi: [10.2307/3495552](https://doi.org/10.2307/3495552)
- Cushing PE 2012 Spider-ant associations: an updated review of myrmecomorphy, myrmecophily, and myrmecophagy in spiders. – Psyche 2012 (151989): 1–23 – doi: [10.1155/2012/151989](https://doi.org/10.1155/2012/151989)
- Donisthorpe H 1927 The guests of British ants, their habits and life histories. Routledge and Sons, London. 244 pp.
- Hölldobler B & Wilson EO 1990 The ants. Harvard University Press, Cambridge, Massachusetts. 746 pp.
- Lenoir A, D'Ettorre P & Errard C 2001 Chemical ecology and social parasitism in ants. – Annual Review of Entomology 46: 573–599 – doi: [10.1146/annurev.ento.46.1.573](https://doi.org/10.1146/annurev.ento.46.1.573)
- Lenoir A, Háva J, Hefetz A, Dahbi A, Cerdá X & Boulay R 2013 Chemical integration of *Thorictus* myrmecophilous beetles into *Cataglyphis* ant nests. – Biochemical Systematics and Ecology 51: 335–342 – doi: [10.1016/j.bse.2013.10.002](https://doi.org/10.1016/j.bse.2013.10.002)



**Fig. 3:** *Formica lugubris* mound nest in the Spanish Pyrenees in which the myrmecophilic linyphiid *Thyreosthenius biovatus* was found

- Lenzini L, Castellucci F, Poso M, Kulczycki A, Simeon E, Greco G, Piccinini A & Legittimo CM 2022 First records of *Anagraphis ochracea* (Araneae, Gnaphosidae) for continental Italy and Sicily with new observations about its myrmecophilous lifestyle. – Arachnologische Mitteilungen 64: 83–92 – doi: [10.30963/aramit6410](https://doi.org/10.30963/aramit6410)
- Nentwig W, Blick T, Bosmans R, Gloor D, Hänggi A & Kropf C 2023 araneae – Spiders of Europe, version 06.2023. – Internet: <https://araneae.nmbe.ch> (4. Jun. 2023) – doi: [10.24436/1](https://doi.org/10.24436/1)
- Palmgren P 1976 Die Spinnenfauna Finnlands und Ostfennoskandien. VII. Linyphiidae 2 (Micryphantinae, mit Ausnahme der Linyphiidae-ähnlichen). – Fauna Fennica 29: 1–126
- Parmentier T, Dekoninck W & Wenseleers T 2014 A highly diverse microcosm in a hostile world: a review on the associates of red wood ants (*Formica rufa* group). – Insectes Sociaux 61: 229–237 – doi: [10.1007/s00040-014-0357-3](https://doi.org/10.1007/s00040-014-0357-3)
- Parmentier T, Dekoninck W & Wenseleers T 2015 Meta-population processes affecting diversity and distribution of myrmecophiles associated with red wood ants. – Basic and Applied Ecology 16: 553–562 – doi: [10.1016/j.baae.2015.04.008](https://doi.org/10.1016/j.baae.2015.04.008)
- Parmentier T, Dekoninck W & Wenseleers T 2017 Arthropods associate with their red wood ant host without matching nestmate recognition cues. – Journal of Chemical Ecology 43: 644–661 – doi: [10.1007/s10886-017-0868-2](https://doi.org/10.1007/s10886-017-0868-2)
- Parmentier T, De Laender F & Bonte D 2020 The topology and drivers of ant-symbiont networks across Europe. – Biological Reviews 95: 1664–1688 – doi: [10.1111/brv.12634](https://doi.org/10.1111/brv.12634)
- Roberts MJ 1987 The spiders of Great Britain and Ireland, Volume 2: Linyphiidae and check list. Harley Books, Colchester. 204 pp.
- Robinson N 1998 First records of the myrmecophile spider *Thyreosthenius biovatus* Cambridge in nests of the wood ant *Formica lugubris* Zetterstedt. – The Carlisle Naturalist 6 (2): 33.
- Scharff N & Gudik-Sørensen O 2006 Katalog over Danske edderkopper [Catalogue of Danish spiders (Araneae)]. – Entomologiske Meddelelser 74: 3–71 [in Danish]
- Schneider CA, Rasband WS & Eliceiri KW 2012 NIH Image to ImageJ: 25 years of image analysis. – Nature Methods 9: 671–675 – doi: [10.1038/nmeth.2089](https://doi.org/10.1038/nmeth.2089)
- Seifert B 2021 A taxonomic revision of the Palearctic members of the *Formica rufa* group (Hymenoptera: Formicidae) – the famous mound-building red wood ants. – Myrmecological News 31: 133–179 – doi: [10.25849/myrmecol.news\\_031:133](https://doi.org/10.25849/myrmecol.news_031:133)
- Simon E 1926 Les arachnides de France. Synopsis générale et catalogue des espèces françaises de l'ordre des Araneae. Tome VI. 2e partie. Roret, Paris. 309–532
- Wasmann E 1894 Kritisches Verzeichniss der myrmecophilin und termitophilen Arthropoden. Felix Dames, Berlin. 231 pp.
- Witte V, Leingärtner A, Sabass L, Hashim R & Foitzik S 2008 Symbiont microcosm in an ant society and the diversity of interspecific interactions. – Animal Behavior 76: 1477–1486 – doi: [10.1016/j.anbehav.2008.05.010](https://doi.org/10.1016/j.anbehav.2008.05.010)
- Wiehle H 1960 Spinnentiere oder Arachnoidea (Araneae). XI. Micryphantidae - Zwergspinnen. – Die Tierwelt Deutschlands 47: 1–620
- World Spider Catalog 2023 World spider catalog, version 24.0. Natural History Museum, Bern. – Internet: <https://wsc.nmbe.ch> (4. Jun. 2023) – doi: [10.24436](https://doi.org/10.24436)