

INDEX TO VOLUME 68

Source: Madroño, 68(4) : 515-516

Published By: California Botanical Society

URL: https://doi.org/10.3120/0024-9637-68.4.515

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

MADROÑO, Vol. 68, No. 4, pp. 515-516, 2021

INDEX TO VOLUME 68

Classified entries: Major subjects, key words, and results; botanical names (new names are in **boldface**); geographical areas; reviews, commentaries. Incidental references to taxa (including most lists and tables) are not indexed separately. Species appearing in Noteworthy Collections are indexed under name, family, and state or country. Authors and titles are listed alphabetically by author in the Table of Contents to the volume.

- Alaska: *Corallorhiza maculata*, and its relationship to *C. mertensiana*, 75; fruit retention in four berry species, 487.
- Arctostaphylos and Ceanothus, absence of flowering shifts over the past century of climate warming, 461.
- Arizona (see Sonoran Desert)
- Asteraceae: Cirsium hydrophyllum var. hydrophylum, limiting life history stages, 39; Lasthenia californica, late planting shortens flowering period and reduces fecundity, 377; L. gracilis, growth response to simulated drought, 366.
 - New taxon: *Chaenactis kyhosii*, from Baja Calif., Mexico, 122; *Helianthus devernii* from NV, 52.
 - Noteworthy collections: *Dittrichia viscosa*, from CA, 6; *Taraxacum alaskanum* and *T. scopulorum* from WA, 7.

Berries, winter retention in four AK spp., 487.

Bird Haven Ranch flora, CA, 23.

Boechera cascadensis, noteworthy collection from WA, 7.

Boraginaceae (see Cryptantha and Eriodictyon)

Botrychium hesperium, noteworthy collection from ID, 71. Brassicaceae (see *Boechera*)

- Bryofloras, contributions toward for CA, OR, and WY, 139-335.
- Bryophytes: Cascade-Siskiyou Natl. Monument, 257; Contra Costa Co., Mt. Diablo, 163; Napa Co., 191; San Luis Obispo Co., 141; Siskiyou Co., 209.

Cactaceae (see Carnegia and chollas)

- California: Arctostaphylos and Ceanothus, absence of flowering shifts over the past century of climate warming, 461; Bird Haven Ranch flora, 23; Calochortus plummerae, interannual variation in flowering, 360; Cirsium hydrophilum var. hydrophilum, limiting life history stages, 39; Clarkia cylindrica and C. unguiculata have similar phenological sensitivities, 388; Dirca occidentalis, association of annual dormancy release and fruit set, 416; habitat shifts and flowering phenology overlap, 406; Lasthenia: L. caifornica, late planting shortens flowering period and reduces fecundity, 377; L. gracilis, growth response to simulated drought, 366; Phytophthora ramorum infection effect on Notholithocarpus densiflorus and Umbellularia californica, 99; Quercus lobata, budburst timing, 434; winter precipitation as climatic driver of spring phenology in five Quercus spp., 450; woody vegetation encroachment as a driver of herbaceous spp. loss in a coastal fen, 9.
 - Noteworthy collections: Carnegia gigantia, 61; Castilleja angustifolia var. flavescens, 4; Clinopodium menthifolium subsp. ascendens, 64; Cyrtomium falcatum, 67; Sphenopholis interrupta subsp. californica, 69.
- Calochortus plummerae, interannual variation in flowering, 360.

Carnegia gigantia, noteworthy collection from CA, 61.

- *Castilleja angustifolia* var. *flavescens*, noteworthy collection from CA, 4.
- Ceanothus and Arctostaphylos, absence of flowering shifts over the past century of climate warming, 461

Chaenactis kyhosii, new sp. From Baja Calif., Mexico, 122. Chollas, evolution of, 109.

- Cirsium hydrophilum var. hydrophilum, limiting life history stages, 39.
- *Clarkia*: Phenological sensitivities similar in *C. cylindrica* and *C. unguiculata*, 388.
- Climate change and phenology: several articles, 339-510.
- *Clinopodium menthifolium* subsp. *ascendens*, noteworthy collection from CA, 64.
- Corallorhiza maculata, in AK, and its relationship to C. mertensiana, 75.
- Cryptantha arenophila, new sp. From Baja Calif., Mexico, 127.

Cylindropuntia, (see cholla)

Cyrtomium falcatum, noteworthy collection from CA, 67.

Diphasiastrum alpinum, noteworthy collection from ID, 71. *Dirca occidentalis*, association of annual dormancy release and fruit set, 416.

Dittrichia viscosa, noteworthy collection from CA, 6.

Drought (see Lasthenia)

Dryopteridaceae (see Cyrtomium)

Editor's report, 512.

- Elymus hirsutus, noteworthy collection from ID, 71.
- Encroachment: Woody vegetation as a driver of herbaceous spp. diversity loss in a CA coastal fen, 9.

Ericaceae (see Arctostaphylos)

- Eriodictyon capitatum, smoke-induced germination, 87.
- *Eschscholzia california*, intraspecific variation in flowering date in relation to climate change, 343.

Euphrasia nemorosa, noteworthy collection from ID, 71.

Fagaceae (see *Quercus*)

- Fen, coastal (see Encroachment)
- Fire response (see *Calochortus*)

Flowering phenology overlap and habitat shifts, 406.

Germination, smoke-induced (see *Eriodictyon*) *Grusonia* (see chollas)

Habitat shifts and flowering phenology overlap, 406. *Helianthus devernii*, new sp. from NV, 52.

- Idaho: Noteworthy collections: *Botrychium hesperium*, *Diphasiastrum alpinum*, *Elymus hirsutus*, *Euphrasia nemorosa*, *Utricularia ochroleuca*, 71.
- Keys: Cryptantha arenophila, C. intermedia var. intermedia, and C. patula, 136; Helianthus spp. of NV, 56.

Lamiaceae (see *Clinopodium* and *Trichostema*)

Lasthenia: L. caifornica, late planting shortens flowering period and reduces fecundity, 377; L. gracilis, growth response to simulated drought, 366.

Lentibulareaceae (see Utricularia)

Lauraceae (see Umbellularia)

MADROÑO

Liliaceae (see *Calochortus*) Lycopodiaceae (see *Diphasiastrum*)

Marsh, brackish (see *Cirsium*) MEXICO (see *Chaenactis* and *Cryptantha*) *Micropuntia* (see chollas)

Namaceae (see Eriodictyon) Nevada (see Helianthus) Notholithocarpus (see Phytopthora)

Onagraceae (see *Clarkia*) Ophioglossaceae (see *Botyrchium*) Orchidaceae (see *Corallorhiza*) Orobanchaceae (see *Castilleja* and *Euphrasia*) Oregon: Bryophytes, Cascade-Siskiyou Natl. Monument, 257.

Papaveraceae (see *Eschscholzia*)Phenology: Introduction to patterns in flora of western No. Am., 339; several articles, 339-510.

Phytophthora ramorum infection, effect on Notholithocarpus densiflorus and Umbellularia californica, 99.
Poaceae (see Elymus and Sphenopholis)
President's report, 511.

Quercus: Q. lobata, budburst timing, 434; winter precipitation as climatic driver of spring phenology in five Calif. *Quercus* spp., 450. Rain shadow (see Corallorhiza)
Redwood Natl. Park, CA (see Phytopthora)
Restoration, wetlands (see Bird Haven Ranch)
Reviewers of 2021 manuscropts, 514.
Reviewers; Botanical revelation: European encounters with Australian plants before Darwin by D. J. Mabberly, 59; Driven by nature: A personal journey from Shanghai to botany and global sustainability by Peter H. Raven, 57.
Rhamnaceae (see Ceanothus)

Sonoran Desert: Dominant plant spp. have divergent phenological responses to climate change, 473.Sphenopholis interrupta subsp. californica, noteworthy collection from CA, 69.Sudden oak death (see Phytopthora)

Taraxacum alaskanum and T. scopulorum, noteworthy collections from WA, 7.
Taylor, Dean William, Vol.68 dedicated to, 1.
Thymelaeaceae (see Dirca)
Trichostema, nomencatural notes, 20.

Umbellularia (see *Phytopthora*) *Utricularia ochroleuca*, noteworthy collection from ID, 71.

Washington: Noteworthy collections, *Boechera cascadensis*, *Taraxacum alaskanum and T. scopulorum*, 7.Wyoming: Bryophytes of Shoshone Natl. Forest, 283.

516