

Genetic Similarity Amongst Phenotypically Diverse Little Free-Tailed Bats, *Chaerephon pumilus*

Authors: Jacobs, David S., Eick, Geeta N., Richardson, Eleanor J., and Taylor, Peter J.

Source: *Acta Chiropterologica*, 6(1) : 13-21

Published By: Museum and Institute of Zoology, Polish Academy of Sciences

URL: <https://doi.org/10.3161/001.006.0102>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Genetic similarity amongst phenotypically diverse little free-tailed bats, *Chaerephon pumilus*

DAVID S. JACOBS¹, GEETA N. EICK², ELEANOR J. RICHARDSON³, and PETER J. TAYLOR⁴

¹Department of Zoology, University of Cape Town, Private Bag, Rondebosch 7701, South Africa
E-mail: djacobs@botzoo.uct.ac.za

²Evolutionary Genomics Group, University of Stellenbosch, Private Bag XI, Matieland, South Africa
³52 Bowen Ave, Glenmore, Durban, 4001, South Africa

⁴Durban Natural Science Museum, P.O. Box 4085, Durban, South Africa

The African molossid *Chaerephon pumilus* shows extensive variation in colour, size and echolocation across its wide distributional range with a light-winged form in north-eastern Africa and a dark form in southern Africa. There is also much variation in supposedly diagnostic characters (e.g., degree of palatal emargination) amongst the dark form of this species in southern Africa. These differences suggest that there may be cryptic species within *C. pumilus*. We used phylogenetic and phenetic analyses of sequence data from the mitochondrial cytochrome-*b* gene of a number of *C. pumilus* individuals to investigate the status of the light and dark-winged forms of this species and to evaluate the possibility of cryptic species within the dark-winged form of *C. pumilus* in southern Africa. We evaluated species status by comparing the level of sequence divergence amongst *C. pumilus* with the level of sequence divergence between known species in the genus. These included *C. ansorgei*, *C. chapini*, *C. nigeriae* and *C. jobensis*. Intrageneric sequence divergences among the *Chaerephon* spp. included here ranged from 6.51 to 11.18%, whereas the average sequence divergence between the light and dark forms was 0.9%. This suggests that these two forms are not distinct species. Individuals of the dark form of *C. pumilus* were genetically indistinguishable from each other having the same cytochrome *b* haplotype. We thus found no evidence of cryptic species in southern African *C. pumilus*.

Key words: *Chaerephon pumilus*, cytochrome *b*, genetics species concept

INTRODUCTION

There is much evidence for intraspecific variation in bat morphology (Bogdanowicz, 1990; Jacobs, 1996, 1999a; Aspetsberger *et al.*, 2003; Miller-Butterworth *et al.*, 2003), diet (Jacobs, 1999b; Brack and Whitaker, 2001; Aspetsberger *et al.*, 2003) and echolocation calls (Buchler, 1980; Heller and von Helversen, 1989; Jones *et al.*, 1992; Obrist, 1995; Betts, 1998; Jacobs, 1999a; Aspetsberger *et al.*, 2003). Such variation can

be small in magnitude and in some cases might mask cryptic species. For example, *Pipistrellus pipistrellus*, has been shown, on the basis of differences in echolocation calls (Jones and van Parijs, 1993), to consist of two distinct species despite a marked similarity in their morphology. Genetic divergence supports the reclassification of *P. pipistrellus* as two different species (Barratt *et al.*, 1997; Mayer and von Helversen, 2001a). Similarly, large genetic differences in morphologically similar species of

Plecotus spp. (Mayer and von Helversen, 2001b; Kiefer *et al.*, 2002; Mucedda *et al.*, 2002) and *Myotis* spp. (Mayer and von Helversen, 2001b; von Helversen *et al.*, 2001) indicate the existence of cryptic species. Therefore, a widely distributed species that displays much morphological divergence might in fact consist of more than one species. However, very little genetic differences have been found between *Eptesicus serotinus* and *E. nilssonii*, two well-recognized species that differ in many morphological characteristics (Mayer and von Helversen, 2001b).

The African molossid *Chaerephon pumilus* (Cretzschmar, 1830; see Bouchard, 1998 for review of the species taxonomy) is a 10–14g insectivorous bat that is widespread in Africa south of the Sahara (Hayman and Hill, 1971; Skinner and Smithers, 1990; Bouchard, 1998; Jacobs and Fenton, 2001). This wide distribution is accompanied by considerable variation in colour and size. Populations of *C. pumilus* vary in the colour of wings and venter (Bouchard, 1998), with dark or light wings being the primary character to divide the forms. At Amani in Tanzania, the population consists of the light form which has pale almost transparent wings, white abdomens and white to yellow bands of fur on the wings running along the flank of the bats from armpit to leg, characters that had been associated with the subspecies *C. p. limbata* (Koopman, 1965). However, this subspecies has been dropped because sub-specific separation based on colour is not possible. Bats of different colour sometimes occupy the same roost (Hayman and Hill, 1971). The form, which is found in southern Africa, is dark in colour with dark wings and body. However, there is some variation within this group in the white colouring on the venter. Some do not have the white undersides or flank bands (Hayman and Hill, 1971; Aspetsberger *et al.*, 2003) while others have

the white flank bands with the white on the abdomen reduced, ranging from a tiny spot near the genitalia to a wide white stripe the whole length of the abdomen. Skinner and Smithers (1990) include both the dark and light forms in *Chaerephon pumilus*.

Chaerephon pumilus is larger in Kenya (mean forearm = 42.8 mm; mass = 13.8 g; Smart and Clark, 1991) than elsewhere in Africa (mean forearm range = 36–38 mm; mean mass = 10.8 g; Marshall and Corbet, 1959; Kingdon, 1974; McWilliam, 1988; Aspetsberger *et al.*, 2003). In Kenya *C. pumilus* has two breeding seasons (Harrison, 1958), but in Uganda and West Africa it breeds continuously (Marshall and Corbet, 1959; Bouchard, 1998). Similarly, although *C. pumilus* uses the same foraging habitat (i.e., open areas) in different parts of its range (Aldridge and Rautenbach, 1987; McWilliam, 1988), there is considerable variation in its diet. In the Kruger National Park, South Africa, this species eats mainly Coleoptera and Hemiptera (Aldridge and Rautenbach, 1987). In Kenya the diet varied greatly. Whitaker and Mumford (1978) found that *C. pumilus* eats mainly Hemiptera and Coleoptera with some Lepidoptera. However, at Lake Naivasha, also in Kenya, the diet consists of more than 50% Diptera with Coleoptera and Hemiptera eaten to a lesser extent (Clark and Smart, 1991). At Amani, Tanzania, Blattodea form a major component of the diet of *C. pumilus* with Hemiptera being the only other order to make up more than 10% of its diet (Aspetsberger *et al.*, 2003). Variation in the echolocation of *C. pumilus* has not been as widely studied, but Aspetsberger *et al.* (2003) report that *C. pumilus* at Amani, Tanzania (the light form) used calls of lower frequency, shorter duration and longer interpulse interval (IPI) than *C. pumilus* in South Africa (the dark form). These morphological, dietary and echolocation differences between the dark and light forms of

C. pumilus suggest cryptic species within *C. pumilus*. We investigated this possibility using mitochondrial cytochrome-*b* sequence data.

Furthermore, on the basis of morphological ambiguities (e.g., pronounced palatal emargination across a wide range of groups previously recognized as separate species), Taylor (1999) suggested a revision of Koopman's (1975) key for the genus. Using the revised key Taylor (1999) identified a number of *Chaerephon* spp. as *C. pumilus* that would have been classified as *C. nigeriae* or *C. ansorgei* using Koopman's (1975) key. We thus used tissue samples from some of the specimens used by Taylor (1999) for the revision of Koopman's (1975) key to determine if genetic data were consistent with Taylor's (1999) revised species key.

MATERIALS AND METHODS

Sampling

We collected tail punches or heart tissue from *C. pumilus* from a variety of localities in South Africa, Zimbabwe, Zambia and Tanzania (Table 1). Tail punches were taken using a 3 mm diameter skin biopsy punch. Individuals of the dark form of *C. pumilus* were collected from the Durban metropolitan area (29°52'S, 30°59'E; 12 bats), Dalton (29°22'S, 30°37'E; one bat) and the 121 SA Infantry Battalion Training Base at Hell's Gate on the shores of Lake St. Lucia (28°03'S, 32°25'E; two bats) both sites in the KwaZulu-Natal Province of South Africa.

We collected tail punches from light forms of *C. pumilus* from Amani Nature Reserve (5°10'S, 38°47'E; one bat), Tanzania and from near Kitwe, Zambia (12°56'S, 28°18'E; five bats – DSJZM101, 107, 109, 113, 114, University of Cape Town). We also used heart tissue from five specimens [Durban Natural Science Museum (DNSM): DM6283–DM6287] identified as *C. ansorgei* and four specimens (DM 6299–DM6302) identified as *C. pumilus* using Koopman's (1975) key. Taylor (1999) later identified all nine specimens as *C. pumilus* using his revised key. Tissue samples of known *C. ansorgei*, *C. nigeriae*, *C. chapini* and *C. jobensis* were used to determine the degree of sequence divergence between valid species in the genus *Chaerephon*. Vouchers of the first three species from which the tissue samples were

taken reside in the Natural History Museum (NHM), Bulawayo, Zimbabwe. These were taken at the confluence of the Lutope and Ngolangola Rivers at Sebungwe (18°17'S, 28°5'E). FPD Cotterill of the NHM supplied the tissue samples (Table 1). The tissue sample from *C. jobensis* was obtained from the Australian Museum (EBU23959). *Cistugo seabrai* was included in this study as an outgroup taxon for *Chaerephon* given the sister-taxa status of Molossididae and Vespertilionidae (Teeling *et al.*, 2002).

Genetic Analysis

Total genomic DNA was extracted from tail punches or tissue preserved in 95% ethanol using a standard phenol-chloroform extraction procedure (Sambrook *et al.*, 1989). A 604 bp fragment of the mitochondrial cytochrome *b* gene was amplified using primers L14724 and H15275 (Irwin *et al.*, 1991). Polymerase chain reactions (PCRs) were performed in 25 l reaction volumes containing the following reagents: 0.2 U BIOTAQ DNA polymerase (Bioline, Whitehead Scientific), 1 X Bioline reaction buffer, 0.2 mM dNTPs, 3 mM MgCl₂, 0.4 M forward and reverse primer and approximately 100 ng of template DNA. PCR reaction mixtures were covered with a layer of mineral oil before thermal cycling on an MJ Research Inc. PTC-100™ thermal cycler. The cycling profile consisted of an initial denaturation step at 94°C for 5 minutes, followed by 35 cycles of denaturation at 94°C for 30 seconds, primer annealing at 50°C for 45 seconds and extension at 72°C for 45 seconds, followed by a final extension step at 72°C for 10 minutes. PCR products were sequenced directly following automated sequencing protocols (ABI Prism® BigDye™ Cycle Sequencing Ready Reaction Protocol manual). Cycle sequencing products were purified by ethanol precipitation and analyzed on an ABI Prism 3100 Genetic Analyzer (Applied Biosystems, U.S.A.). Sequences were trimmed to 423 bases to facilitate comparisons amongst all individuals sequenced, and aligned using Clustal X (Thompson *et al.*, 1997). All sequences generated in this study have been deposited in Genbank (Table 1). Sequence data was analysed using both phylogenetic (parsimony) and phenetic (genetic distance) methods in PAUP* (Swofford, 2002). In the parsimony analysis, all nucleotide characters were given equal weight. Nodal support was evaluated using bootstrap values (Felsenstein, 1985) generated from 1000 pseudoreplicate data sets. For purposes of comparison with previously published papers, genetic distances were calculated using the Kimura 2-parameter model of nucleotide substitution (Kimura, 1980) under the criterion of minimum evolution.

RESULTS

Five haplotypes (A–E) were identified among the 30 *C. pumilus* sampled in southern Africa (Table 1). When included in a parsimony analysis, these five haplotypes formed a highly supported monophyletic group that was also recovered in a phenetic analysis of the data (Fig. 1). The *Chaerephon* spp. from Hell's Gate identified as *C. ansorgei* using Koopman's (1975) key, had

the same haplotype as *C. pumilus* individuals sampled at the same site (Table 1) but differed from the known *C. ansorgei* by 10.62% (Table 1). The dark form of *C. pumilus* in South Africa displayed slightly more variability represented by three haplotypes (A, B and E; Table 1) with the sequence divergence among these haplotypes ranging from 0.47 to 1.68% (Table 2). Intrageneric species divergences in *Chaerephon* based on the species included in this

TABLE 1. List of *Chaerephon* specimens. Dark (D) or Light (L) winged forms of *C. pumilus* are indicated in parentheses after the species name. * — bats identified using Koopman's (1975) key

Species	Accession No.	Collection Locality	Haplotype	Genbank No.
* <i>C. ansorgei</i> (D)	DM6287	Hell's Gate, South Africa	B	AY377940
* <i>C. ansorgei</i> (D)	DM6285	Hell's Gate, South Africa	B	AY377941
* <i>C. ansorgei</i> (D)	DM6284	Hell's Gate, South Africa	B	AY377942
* <i>C. ansorgei</i> (D)	DM6286	Hell's Gate, South Africa	B	AY377947
* <i>C. ansorgei</i> (D)	DM6283	Hell's Gate, South Africa	B	AY377966
<i>C. ansorgei</i>	FWC4783	Sebungwe, Zimbabwe	n.a.	AY377967
<i>C. chapini</i>	FWC3664	Zambia	n.a.	AY591329
<i>C. jobensis</i>	EBU23959	Solomon Islands	n.a.	AY591331
<i>C. nigeriae</i>	FWC3824	Zambia	n.a.	AY591330
<i>C. pumilus</i> (D)	—	Durban, Amanzimtoti	B	AY377939
* <i>C. pumilus</i> (D)	DM6301	Hell's Gate	B	AY377943
* <i>C. pumilus</i> (D)	DM6299	Hell's Gate	B	AY377944
* <i>C. pumilus</i> (D)	DM6302	Hell's Gate	B	AY377945
* <i>C. pumilus</i> (D)	DM6300	Hell's Gate	B	AY377946
<i>C. pumilus</i> (D)	—	Durban, Yellowwood Park	A	AY377948
<i>C. pumilus</i> (D)	—	Hell's Gate	B	AY377949
<i>C. pumilus</i> (D)	—	Hell's Gate	B	AY377950
<i>C. pumilus</i> (D)	—	Durban, Amanzimtoti	B	AY377951
<i>C. pumilus</i> (D)	—	Durban, Springfield	E	AY377952
<i>C. pumilus</i> (D)	—	Durban, Carrington Heights	E	AY377953
<i>C. pumilus</i> (D)	—	Durban, Westville	E	AY377954
<i>C. pumilus</i> (L)	—	Tanzania	C	AY377955
<i>C. pumilus</i> (D)	—	Durban, Glenwood	E	AY377957
<i>C. pumilus</i> (D)	—	Durban, Pinetown	E	AY377958
<i>C. pumilus</i> (D)	—	Durban, Waterfall	E	AY377959
<i>C. pumilus</i> (D)	—	Durban, Gillitts	E	AY377960
<i>C. pumilus</i> (D)	—	Durban, Glenmore	E	AY377962
<i>C. pumilus</i> (D)	—	Dalton	E	AY377964
<i>C. pumilus</i> (D)	—	Durban, Newlands	E	AY377965
<i>C. pumilus</i> (L)	DSJZM114	Zambia	C	AY500285
<i>C. pumilus</i> (L)	DSJZM107	Zambia	C	AY500286
<i>C. pumilus</i> (L)	DSJZM109	Zambia	C	AY500287
<i>C. pumilus</i> (L)	DSJZM113	Zambia	D	AY500288
<i>C. pumilus</i> (L)	DSJZM101	Zambia	C	AY500289
<i>Cistugo seabrai</i>	M977	Northern Cape, S. Africa	n.a.	AY591332

analysis range from 6.51–11.18%, with *C. ansorgei* the most basal and genetically divergent from the other *Chaerephon* species (Table 2, Fig. 1). Sequence divergence between the genus *Chaerephon* and *Cistugo* ranged from 22.86% to 25.31% (Table 2). The divergence between the five *pumilus* haplotypes ranged from 0.47% to 1.68% (Table 2). The light-winged form (haplotypes C and D) differed by an average of 0.9% from the dark-winged *C. pumilus* (haplotypes A, B and E).

DISCUSSION

Relationships within *C. pumilus* identified using genetic distances were

identical to those recovered under the parsimony criterion. This suggests that our sequence divergence values are likely to be good indicators of species status (Bradley and Baker, 2001) in the genus *Chaerephon*. Despite the marked morphological, ecological and echolocation differences between the light and dark forms of *C. pumilus* (Aspetsberger *et al.*, 2003) the low average genetic divergence (0.9%) between them suggests that they are all one species. This is also true for the dark form in South African. Not only do all the *C. pumilus* fall within a strongly supported monophyletic clade (Fig. 1) but the genetic divergence between them (including dark southern African forms and the light form from




FIG. 1. Minimum evolution tree constructed using Kimura 2-parameter genetic distances for members of the bat genus *Chaerephon*. Maximum parsimony (MP) bootstrap values are indicated above nodes, and minimum evolution (ME) bootstrap values below nodes. Branch lengths are proportional to the number of substitutions per site as indicated by the scale bar

Tanzania and Zambia) was much lower than that between known species in the genus (0.47–1.68% cf. 6.51–11.18%, Table 2). Furthermore, the range of divergence among the different forms of *C. pumilus* falls below the 2% criterion identified by Bradley and Baker (2001) for intraspecific variation.

The four Hell's Gate individuals identified as *C. pumilus* and the five individuals from the same site identified as *C. ansorgei* using Koopman's (1975) key were genetically indistinguishable. In addition, sequence divergence between the Hell's Gate individuals and other *C. pumilus* ranged from 0.47–1.19% (Table 2), which is well within the range of intraspecific variation characterizing other bat species (Johns and Avise, 1998; Ditchfield, 2000; Bradley and Baker, 2001; Ruedi and Mayer, 2001). This suggests that all the Hell's Gate individuals are in fact *C. pumilus*. This is supported by the large sequence divergence between known *C. ansorgei* and the '*C. ansorgei*' from Hell's Gate (10.62%, Table 1) and the basal relationship of *C. ansorgei* to the highly supported *C. pumilus* clade (Fig. 1). This finding is consistent with Taylor's (1999) identification of the Hell's Gate specimens as *C. pumilus* based on his revision of Koopman's (1975) key. However, the genetic similarity between the different forms of *C. pumilus* does not support Taylor's (1999) suggestion of the existence of cryptic species.

The divergence we found between known *Chaerephon* species (6.51–11.18%) overlaps the range (7–11%) suggested by Bradley and Baker (2001) to be that of valid species and where the burden of proof should shift to the documentation that only a single species is involved. We are therefore confident that the low sequence divergence

TABLE 2. Percentage pairwise sequence divergence values for the taxa included in the study under a Kimura 2-parameter model of evolution. The species and haplotype names are abbreviated in the column titles as Cp A = *C. pumilus* haplotype A (for example), Ca = *C. ansorgei*, Cn = *C. nigeriae*, Cc = *C. chapini*, Cj = *C. jobensis* and Ciss = *Cistugo seabrai*

Species/Haplotype	Cp B	Cp A	Cp E	Cp C	Cp D	Ca	Cn	Cc	Cj	Ciss
<i>Chaerephon pumilus</i> B	–									
<i>C. pumilus</i> A	0.47	–								
<i>C. pumilus</i> E	1.19	1.68	–							
<i>C. pumilus</i> C	0.71	1.19	0.48	–						
<i>C. pumilus</i> D	0.95	1.43	0.71	0.24	–					
<i>C. ansorgei</i>	10.62	11.18	10.31	10.31	10.60	–				
<i>C. nigeriae</i>	7.10	7.67	6.51	6.79	7.09	10.94	–			
<i>C. chapini</i>	8.14	8.68	8.13	7.85	8.13	11.13	9.11	–		
<i>C. jobensis</i>	9.81	9.79	9.50	10.07	10.36	8.16	8.23	10.60	–	
<i>Cistugo seabrai</i>	24.19	24.86	22.86	23.20	22.86	20.89	24.86	25.31	23.81	–

between light and dark forms of *C. pumilus* is indicative of intraspecific variation despite the remote possibility that in some cases speciation may be completed at < 2% genetic differentiation (Bradley and Baker 2001; Mayer and von Helversen, 2001b). However, genetic sequence divergence for some taxa can be overestimated as a result of poor sampling. It is possible that our sequence divergence between known *Chaerephon* species may not be an accurate reflection of sequence divergence within this genus because we included only five of the 18 recognised species of *Chaerephon* (Simmons, In press). The possibility exists that some valid species within this genus could be separated by lower levels of sequence divergence. However, Bradley and Baker (2001) did not find < 2% sequence divergence in mitochondrial cytochrome *b* between recognized species in any of the 21 species pairs they tested, ten of which were bat species pairs. We therefore consider it unlikely that our interpretation of the low sequence divergence between the different forms of *C. pumilus* is incorrect.

In conclusion, the existence of cryptic species where there is little morphological variation (Barratt *et al.*, 1997; Mayer and von Helversen, 2001a, 2001b; von Helversen *et al.*, 2001; Kiefer *et al.*, 2002) and the absence of cryptic species (this study) or genetic divergence (Mayer and von Helversen, 2001b) where there is much morphological variation is indicative of the complexity and unpredictability of the process of speciation.

ACKNOWLEDGEMENTS

This study was funded by grants from the University Research Committee of the University of Cape Town and the National Research Foundation to

DSJ. We thank the Australian Museum and FPD Cotterill for tissue loans.

LITERATURE CITED

- ALDRIDGE, H. D. J. N., and I. L. RAUTENBACH. 1987. Morphology, echolocation and resource partitioning in insectivorous bats. *Journal of Animal Ecology*, 56: 763–778.
- ASPETSBERGER, F., D. BRANDSEN, and D. S. JACOBS. 2003. Geographical variation in the morphology, echolocation and diet of the little free-tailed bat, *Chaerephon pumilus* (Molossidae). *African Zoology*, 38: 245–254.
- BARRATT, E. M., R. DEAVILLE, T. M. BURLAND, M. W. BRUFORD, G. JONES, P. A. RACEY, and R. K. WAYNE. 1997. DNA answers the call of pipistrelle bat species. *Nature*, 387: 138–139.
- BETTS, B. J. 1998. Effects of interindividual variation in echolocation calls on identification of big brown and silver-haired bats. *Journal of Wildlife Management*, 62: 1003–1010.
- BOGDANOWICZ, W. 1990. Geographic variation and taxonomy of Daubenton's bat, *Myotis daubentoni*, in Europe. *Journal of Mammalogy*, 71: 205–218.
- BOUCHARD, S. 1998. *Chaerephon pumilus*. *Mammalian Species*, 547: 1–6.
- BRACK, V., JR., and J. O. WHITAKER, JR. 2001. Foods of the northern myotis, *Myotis septentrionalis*, from Missouri and Indiana, with notes on foraging. *Acta Chiropterologica*, 3: 203–210.
- BRADLEY, R. D., and R. J. BAKER. 2001. A test of the genetic species concept: cytochrome-*b* sequences and mammals. *Journal of Mammalogy*, 82: 960–973.
- BUCHLER, E. R. 1980. The development of flight, foraging, and echolocation in the little brown bat (*Myotis lucifugus*). *Behavioural Ecology and Sociobiology*, 6: 211–218.
- CLARK, F. L., and A. C. SMART. 1991. The diet of *Tadarida (Chaerephon) pumila* (Cretzschmar), Molossidae, at Lake Naivasha, Kenya. *Journal of African Zoology*, 105: 493–496.
- DITCHFIELD, A. D. 2000. The comparative phylogeography of Neotropical mammals: Patterns of intraspecific mitochondrial DNA variation among bats contrasted to nonvolant small mammals. *Molecular Ecology*, 9: 1307–1318.
- FELSENSTEIN, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*, 39: 783–791.
- HARRISON, D. L. 1958. A note on successive pregnancies in an African bat. *Mammalia*, 22: 592–595.

- HAYMAN, R. W., and J. E. HILL. 1971. Order Chiroptera. Pp. 1–73, in *The mammals of Africa: an identification manual. Part 2* (J. MEESTER and H. W. SETZER, eds.). Smithsonian Institution Press, Washington D.C.
- HELLER, K.-G., and O. VON HELVERSEN. 1989. Resource partitioning of sonar frequency bands in rhinolophoid bats. *Oecologia*, 80: 178–186.
- IRWIN, D. M., T. D. KOCHER, and A. C. WILSON. 1991. Evolution of the cytochrome *b* gene of mammals. *Journal of Molecular Evolution*, 32: 128–44.
- JACOBS, D. S. 1996. Morphological divergence in an insular bat, *Lasiurus cinereus semotus*. *Functional Ecology*, 10: 622–630.
- JACOBS, D. S. 1999a. Intraspecific variation in wingspan and echolocation call flexibility might explain the use of different habitats by the insectivorous bat, *Miniopterus schreibersii* (Vespertilionidae: Miniopterinae). *Acta Chiropterologica*, 1: 93–103.
- JACOBS, D. S. 1999b. The diet of the insectivorous Hawaiian hoary bat (*Lasiurus cinereus semotus*) in an open and a cluttered habitat. *Canadian Journal of Zoology*, 77: 1603–1608.
- JACOBS, D. S., and M. B. FENTON. 2001. The status of *Sauromys petrophilus* and *Chaerephon pumila* (Chiroptera: Molossidae) in the Western Cape province of South Africa. *African Zoology*, 36: 129–136.
- JOHNS, G. C., and J. C. AVISE. 1998. A comparative summary of genetic distances in the vertebrates from the mitochondrial cytochrome *b* gene. *Molecular Biology and Evolution*, 15: 1481–1490.
- JONES, G., and S. M. VAN PARIJS. 1993. Bimodal echolocation in pipistrelle bats: are cryptic species present? *Proceedings of the Royal Society of London B*, 251: 119–125.
- JONES, G., T. GORDON, and J. NIGHTINGALE. 1992. Sex and age variation in echolocation calls of the lesser horseshoe bat, *Rhinolophus hipposideros*. *Mammalia*, 56: 189–193.
- KIEFER, A., F. MAYER, J. KOSUCH, O. VON HELVERSEN, and M. VEITH. 2002. Conflicting molecular phylogenies of European long-eared bats (*Plecotus*) can be explained by cryptic diversity. *Molecular Phylogenetics and Evolution*, 25: 557–566.
- KIMURA, M. 1980. A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16: 111–120.
- KINGDON, J. 1974. *East African mammals: an atlas of evolution in Africa. Volume 2, Part A* (Insectivores and bats). Academic Press, London, 340 pp.
- KOOPMAN, K. F. 1965. Status of forms described or recorded by J. A. Allen in the American Museum Congo Expedition Collection of bats. *American Museum Novitates*, 2219: 1–34.
- KOOPMAN, K. F. 1975. Bats of the Sudan. *Bulletin of the American Museum of Natural History*, 154: 353–443.
- MARSHALL, A. J., and P. S. CORBET. 1959. The breeding biology of equatorial vertebrates: reproduction of the bat *Chaerephon hindei* Thomas at latitude 0°26'N. *Proceedings of the Zoological Society of London*, 132: 607–616.
- MAYER, F., and O. VON HELVERSEN. 2001a. Sympatric distribution of two cryptic bat species across Europe. *Biological Journal of the Linnean Society*, 74: 365–374.
- MAYER, F., and O. VON HELVERSEN. 2001b. Cryptic diversity in European bats. *Proceedings of the Royal Society of London, Series B — Biological Sciences*, 268: 1825–1832.
- MCWILLIAM, A. N. 1988. Social organisation of the Bat *Tadarida (Chaerephon) pumila* (Chiroptera: Molossidae) in Ghana, West Africa. *Ethology*, 77: 115–124.
- MILLER-BUTTERWORTH, C. M., D. S. JACOBS, and E. H. HARLEY. 2003. Strong population substructure is correlated with morphology and ecology in a migratory bat. *Nature*, 424: 187–191.
- MUCEDDA, M., A. KIEFER, E. PIDINCHEDDA, and M. VEITH. 2002. A new species of long-eared bat (Chiroptera, Vespertilionidae) from Sardinia (Italy). *Acta Chiropterologica*, 4: 121–135.
- OBRIST, M. K. 1995. Flexible bat echolocation: the influence of individual, habitat and conspecifics on sonar signal design. *Behavioural Ecology and Sociobiology*, 36: 207–219.
- RUEDI, M., and F. MAYER. 2001. Molecular systematics of the genus *Myotis* (Vespertilionidae) suggests deterministic ecomorphological convergences. *Molecular Phylogenetics and Evolution*, 21: 436–448.
- SAMBROOK, J., E. F. FRITSCH, and T. MANIATIS. 1989. *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory Press, New York.
- SIMMONS, N. In press. Order Chiroptera. In *Mammal species of the World: a taxonomic and geographic reference*, 3rd edition (D. E. WILSON and D. M. REEDER, eds.).
- SKINNER, J. D., and R. H. N. SMITHERS. 1990. *The mammals of the southern African subregion*. University of Pretoria, Pretoria, xxxii + 771 pp.
- SMART, A. C., and F. L. CLARK. 1991. The emergence behaviour of *Tadarida (Chaerephon) pumila* (Cretzchmar), Molossidae, at Lake Naivasha, Kenya. *Journal of African Zoology*, 105: 497–501.

- SWOFFORD, D. L. 2002. PAUP*: Phylogenetic analyses using parsimony (and other methods), version 4.10. Illinois Natural History Survey, Champaign, Illinois.
- TAYLOR, P. J. 1999. Problems with the identification of southern African *Chaerephon* (Molossidae), and the possibility of a cryptic species from South Africa and Swaziland. *Acta Chiropterologica*, 1: 191–200.
- TEELING, E. C., O. MADSEN, R. A. VAN DEN BUSSCHE, W. W. DE JONG, M. J. STANHOPE, and M. S. SPRINGER. 2002. Microbat paraphyly and the convergent evolution of a key innovation in Old World rhinolophoid microbats. *PNAS*, 99: 1431–1436.
- THOMPSON, J. D., T. J. GIBSON, F. PLEWNIAK, F. JEANMOUGIN, and D. G. HIGGINS. 1997. The CLUSTAL X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research*, 25: 4876–4882.
- VON HELVERSEN, O., K.-G. HELLER, F. MAYER, and A. NEMETH. 2001. Cryptic mammalian species: a new species of whiskered bat (*Myotis alca-thoe* n.sp.) in Europe. *Naturwissenschaften*, 88: 217–223.
- WHITAKER, J. O. JR., and R. E. MUMFORD. 1978. Food and ectoparasites of bats from Kenya, East Africa. *Journal of Mammalogy*, 59: 632–634.

Received 19 December 2003, accepted 14 April 2004